



FOR THE PEOPLE  
FOR EDUCATION  
FOR SCIENCE

LIBRARY  
OF  
THE AMERICAN MUSEUM  
OF  
NATURAL HISTORY







Occasional Papers  
OF THE  
Boston Society of Natural History  
VII.

---

FAUNA OF NEW ENGLAND.

11.

LIST OF THE AVES.

BY GLOVER M. ALLEN.

---

BOSTON:  
PRINTED FOR THE SOCIETY FROM THE  
GURDON SALTONSTALL FUND.  
JUNE, 1909.



## FAUNA OF NEW ENGLAND.

The Library and Publishing Committee will issue, as volume seven of the Occasional Papers, a Catalogue of the animals of New England.

Parts will be published at irregular intervals, and though the details of the several lists will vary somewhat in the different groups, each list will include, 1st, the accepted name (scientific and vernacular); 2d, reference to the original description, with record of locality; 3d, reference to an authentic description and illustration, and 4th, habitat and occurrence.

The Committee considers these lists a necessary preliminary to a series of comprehensive, illustrated monographs, the publication of which it is hoped the Society will at some later time take upon itself.

The Committee also believes that these lists will aid in the accumulation of valuable material for the New England collection of the Society. With this in view a separate check list of names will be issued with each part. In this check list the + indicates a specimen in the Society's Museum; the — that a trustworthy record exists.

Toward the accomplishment of its aims the Committee invites the coöperation of all interested in the Fauna of New England.

Communications concerning specimens should be addressed to the CURATOR of the Society (Chas. W. Johnson), notes and records of occurrence to the EDITOR of the "Fauna of New England" (Samuel Henshaw).

Boston Society Natural History,  
234 Berkeley St., Boston, Mass.



## AVES.

The following list aims to include all species of birds known to have occurred in New England within historic times. Those only whose occurrence is believed to be natural are given a serial number, while introduced, escaped, or hypothetical species are given in a separate list. The nomenclature is that adopted by the American ornithologists' union, including the changes of the 14th supplement to the Check-List, July, 1908. Of the vernacular names, the first given is in each case that of the Check-List, and those following are local names current for the most part in New England. Unless otherwise stated, the term "migrant" means that the species occurs in both the spring and the fall migrations; "summer resident" implies that the species breeds; "resident," that individual birds remain the entire year in the State. The migration and nesting dates given, are almost wholly those available from published sources and are hence in many cases, somewhat meager. Unusual dates are enclosed within parentheses. If the records for the presence of a given species within any State do not exceed three, the place and date of each are given if known.

For brevity of citation, references to figures of eggs in the works of Bendire, Capen, and Seebohm are given by the name of the author followed by the date, plate, and figure. The titles of these works follow: —

Bendire, C. Life histories of North American birds with special reference to their breeding habits and eggs. Special bull. no. 1, U. S. nat. mus., 1892, viii+446 p., 12 pls.; special bull. no. 3, U. S. nat. mus., 1895, ix+518 pp., 7 pls.

Capen, Elwin A. Oology of New England: a description of the eggs, nests, and breeding habits of the birds known to breed in New England, with colored illustrations of their eggs. Boston, 1886, 4to, 116 pp., 25 pls.

Seebohm, Henry. Coloured figures of the eggs of British birds, with descriptive notices. Sheffield, xxiv+304 pp., pls.



	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Colymbus holboellii</i>	—	—	—	+	—	—
<i>auritus</i>	—	—	—	+	—	—
<i>Podilymbus podiceps</i>	—	—	—	+	—	—
<i>Gavia immer</i>	—	—	—	+	—	—
<i>arctica</i>	—	—	—	?	—	?
<i>stellata</i>	—	—	—	+	—	—
<i>Lunda cirrhata</i>	—	—	—	—	—	—
<i>Fratercula arctica</i>	—	+	—	—	?	—
<i>Cephalus grylle</i>	+	—	—	+	—	—
<i>Uria troile</i>	—	—	?	+	?	?
<i>lomvia</i>	—	—	—	+	—	—
<i>Alca torda</i>	—	—	—	+	—	—
<i>Plautus impennis</i>	—	—	—	—	—	—
<i>Alle alle</i>	—	—	—	+	—	—
<i>Megalestris skua</i>	—	—	—	—	—	—
<i>Stercorarius pomarinus</i>	+	—	—	+	—	—
<i>parasiticus</i>	—	—	—	+	—	—
<i>longicaudus</i>	—	—	—	—	—	—
<i>Pagophila alba</i>	—	—	—	—	—	—
<i>Rissa tridactyla</i>	—	—	—	+	—	—
<i>Larus hyperboreus</i>	—	—	?	+	—	—
<i>leucopterus</i>	+	—	—	—	—	—
<i>kumlieni</i>	—	—	—	—	—	—
<i>marinus</i>	—	—	—	+	—	—
<i>argentatus</i>	—	—	—	+	—	—
<i>delawarensis</i>	—	—	—	+	—	—
<i>atricilla</i>	—	—	—	—	—	—
<i>philadelphia</i>	—	—	—	+	—	—
<i>Xema sabinii</i>	—	—	—	—	?	—
<i>Gelochelidon nilotica</i>	+	—	—	—	?	—
<i>Sterna caspia</i>	—	—	—	+	—	—
<i>maxima</i>	—	—	—	—	—	—
<i>sandvicensis acufflava</i>	—	—	—	—	—	—
<i>forsteri</i>	—	—	—	—	—	—
<i>hirundo</i>	—	—	—	+	—	—
<i>paradisaea</i>	—	—	—	—	—	—
<i>dougalli</i>	—	—	—	+	—	—
<i>antillarum</i>	—	—	—	+	—	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Sterna fuscata</i>	—	—	+	—	—	—
<i>Hydrochelidon nigra surinamensis</i>	—	—	—	+	—	—
<i>Rynchops nigra</i>	—	—	—	+	—	—
<i>Fulmarus glacialis</i>	+	—	—	—	—	—
<i>Puffinus borealis</i>	—	—	—	+	—	—
<i>gravis</i>	—	—	—	—	—	—
<i>grisens</i>	—	—	—	+	—	—
<i>Aestrelata hasitata</i>	—	—	—	—	—	—
<i>Daption capensis</i>	—	—	—	—	—	—
<i>Oceanodroma leucorhoa</i>	—	—	—	+	—	—
<i>Oceanites oceanicus</i>	—	—	?	+	—	—
<i>Sula leucogastra</i>	—	—	—	—	—	?
<i>bassana</i>	+	—	—	+	—	—
<i>Phalacrocorax carbo</i>	+	—	—	—	—	—
<i>auritus</i>	—	—	—	+	—	—
<i>Pelecanus erythrorhynchos</i>	—	—	—	—	—	—
<i>occidentalis</i>	—	—	—	—	—	—
<i>Fregata aquila</i>	?	—	—	?	—	—
<i>Mergus americanus</i>	—	—	—	+	+	—
<i>serrator</i>	—	—	—	+	—	—
<i>Lophodytes cucullatus</i>	—	—	—	—	+	—
<i>Anas platyrhynchos</i>	—	—	—	+	—	—
<i>rubripes</i>	—	—	—	+	—	—
<i>Chaulelasmus streperus</i>	—	—	—	—	+	—
<i>Mareca penelope</i>	—	—	?	—	—	—
<i>americana</i>	—	—	—	+	—	—
<i>Nettion crecca</i>	—	—	—	—	—	—
<i>carolinensis</i>	—	—	—	+	—	—
<i>Querquedula discors</i>	—	+	—	+	—	—
<i>Spatula clypeata</i>	—	—	—	+	—	—
<i>Dafila acuta</i>	—	—	—	+	—	—
<i>Aix sponsa</i>	—	—	—	+	—	—
<i>Marila americana</i>	—	—	—	+	—	—
<i>vallisneria</i>	—	—	—	+	—	—
<i>marila</i>	—	—	—	+	+	—
<i>affinis</i>	—	—	—	+	—	—
<i>collaris</i>	—	—	—	+	—	—
<i>Clangula clangula americana</i>	—	—	—	+	—	—
<i>islandica</i>	+	—	—	+	—	—
<i>Charitonetta albeola</i>	—	—	—	+	—	—
<i>Harelda hyemalis</i>	—	—	—	+	—	—
<i>Histrionicus histrionicus</i>	+	—	—	—	—	—
<i>Camptorhynchus labradorius</i>	—	—	—	+	—	—



	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Somateria mollissima borealis</i>	+				?	
<i>dresseri</i>	+	—		+	—	—
<i>spectabilis</i>	+	—		—	—	—
<i>Oidemia americana</i>	—	—	—	+	—	—
<i>deglandi</i>	—	—	—	+	—	—
<i>perspicillata</i>	—	—	—	+	—	—
<i>Erisimatura jamaicensis</i>	—	—	—	+	—	—
<i>Nomonyx dominicus</i>			+	—		
<i>Chen hyperborea</i>	—	—	—	+	—	—
<i>nivalis</i>	—			—		—
<i>caerulescens</i>		—		+	—	
<i>Anser albifrons gambeli</i>			?	—		
<i>Branta canadensis</i>	—	—	—	+	—	—
<i>hutchinsii</i>	—			?		?
<i>bernicla glaucogastra</i>	—	—	—	+	—	—
<i>nigricans</i>				—		
<i>leucopsis</i>	?		—	—		—
<i>Olor cygnus</i>	—					
<i>columbianus</i>	?	—	—	+	—	—
<i>buccinator</i>		—		—		?
<i>Guara alba</i>			—			—
<i>Plegadis autumnnalis</i>		—		+		—
<i>Mycteria americana</i>	—	—	—	—	—	—
<i>Botaurus lentiginosus</i>	—	—	—	+	—	—
<i>Ixobrychus exilis</i>	—	+	—	+	—	—
<i>Ardea herodias</i>	—	—	—	+	—	—
<i>Herodias egretta</i>	—	—	—	—	+	—
<i>Egretta candidissima</i>			—	—		—
<i>Florida caerulea</i>	—	—	?	+	—	—
<i>Butorides virescens</i>	—	—	—	+	—	—
<i>Nycticorax nycticorax naevius</i>	—	—	—	+	—	—
<i>Nyctanassa violacea</i>	—			+	—	
<i>Grus americana</i>			—	—		
<i>canadensis</i>					—	
<i>mexicana</i>		—	—	—		—
<i>Rallus elegans</i>	—			+	—	—
<i>crepitans</i>	—	—	—	+	?	—
<i>virginianus</i>	—	—	—	+	—	—
<i>Porzana carolina</i>	—	—	—	+	—	—
<i>Coturnicops noveboracensis</i>	—	—	—	+	—	—
<i>Creciscus jamaicensis</i>	?			—		—
<i>Crex crex</i>	—				—	—
<i>Ionornis martinica</i>	—	—	?	—	+	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Gallinula galeata</i> . . . . .	—	—	—	+	—	—
<i>Fulica americana</i> . . . . .	—	—	—	+	—	—
<i>Phalaropus fulicarius</i> . . . . .	—	—	—	+	—	—
<i>Lobipes lobatus</i> . . . . .	—	—	—	+	—	—
<i>Steganopus tricolor</i> . . . . .	—	—	—	+	+	—
<i>Recurvirostra americana</i> . . . . .	—	—	—	—	—	—
<i>Himantopus mexicanus</i> . . . . .	—	—	—	—	—	—
<i>Philohela minor</i> . . . . .	—	—	—	+	—	—
<i>Gallinago delicata</i> . . . . .	—	+	—	+	—	—
<i>Macrorhamphus griseus</i> . . . . .	—	—	—	+	—	—
<i>scolopaceus</i> . . . . .				+	—	—
<i>Micropalama himantopus</i> . . . . .	—	+	—	+	—	—
<i>Tringa canutus</i> . . . . .	—	—	—	+	—	—
<i>Arquatella maritima</i> . . . . .	—	—	?	+	—	—
<i>Pisobia maculata</i> . . . . .	+	+	—	+	—	—
<i>fuscicollis</i> . . . . .	—	—	—	+	—	—
<i>bairdii</i> . . . . .	—	—	—	+	—	—
<i>minutilla</i> . . . . .	—	—	—	+	—	—
<i>Pelidna alpina</i> . . . . .	—	—	—	—	—	—
<i>sakhalina</i> . . . . .	—	—	—	+	—	—
<i>Erolia ferruginea</i> . . . . .	+	—	—	—	?	—
<i>Ereunetes pusillus</i> . . . . .	—	—	—	+	—	—
<i>mauri</i> . . . . .	—	—	—	+	—	—
<i>Calidris leucophaea</i> . . . . .	—	—	—	+	—	—
<i>Limosa fedoa</i> . . . . .	—	—	—	+	—	—
<i>haemastica</i> . . . . .	—	+	—	+	—	—
<i>Totanus melanoleucus</i> . . . . .	—	—	—	+	—	—
<i>flavipes</i> . . . . .	—	—	—	+	—	—
<i>Helodromas solitarius</i> . . . . .	—	—	—	+	—	—
<i>Catoptrophorus semipalmatus</i> . . . . .	—	—	—	+	—	—
<i>inornatus</i> . . . . .				+	—	—
<i>Pavoncella pugnax</i> . . . . .	—	—	—	—	—	—
<i>Bartramia longicauda</i> . . . . .	—	—	—	+	—	—
<i>Tryngites subruficollis</i> . . . . .	—	—	?	+	—	—
<i>Actitis macularia</i> . . . . .	—	—	—	+	—	—
<i>Numenius americanus</i> . . . . .	—	—	—	+	—	—
<i>hudsonicus</i> . . . . .	—	—	—	+	—	—
<i>borealis</i> . . . . .	—	—	—	+	—	—
<i>Squatarola squatarola</i> . . . . .	—	—	—	+	—	—
<i>Charadrius dominicus</i> . . . . .	—	+	—	+	—	—
<i>Oxyechus vociferus</i> . . . . .	—	—	—	+	—	—
<i>Aegialitis semipalmata</i> . . . . .	—	—	—	+	—	—
<i>meloda</i> . . . . .	—	—	?	+	—	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Ochthodromus wilsonius</i> . . . . .		?	?	—	?	—
<i>Arenaria interpres</i> . . . . .				—		
<i>morinella</i> . . . . .	—	—	—	+	—	—
<i>Haematopus palliatus</i> . . . . .	—			+		—
<i>Colinus virginianus</i> . . . . .	—	—	—	+	—	—
<i>Canachites canadensis canadensis</i> . . . . .	+	—	—	—		
<i>Bonasa umbellus</i> . . . . .			—	+	—	—
<i>togata</i> . . . . .	—	—	—	—		
<i>Lagopus lagopus</i> . . . . .	—			—	?	
<i>Tympanuchus cupido</i> . . . . .	?			+	—	—
<i>Meleagris gallopavo silvestris</i> . . . . .	—	—	—	—	—	—
<i>Ectopistes migratorius</i> . . . . .	+	—	—	+	—	—
<i>Zenaidura macroura carolinensis</i> . . . . .	—	—	—	+	—	—
<i>Sturnoenas cyanocephala</i> . . . . .				+		
<i>Cathartes aura septentrionalis</i> . . . . .	—	+	—	—	—	—
<i>Catharista uruba</i> . . . . .	—			—		—
<i>Elanoides forficatus</i> . . . . .		?	?	—		—
<i>Circus hudsonius</i> . . . . .	—	—	—	+	—	—
<i>Accipiter velox</i> . . . . .	—	—	—	+	—	—
<i>cooperii</i> . . . . .	—	—	—	+	—	—
<i>Astur atricapillus</i> . . . . .	—	—	—	+	—	—
<i>Buteo borealis</i> . . . . .	—	+	—	+	—	—
<i>lineatus</i> . . . . .	—	—	—	+	—	—
<i>swainsoni</i> . . . . .	—			—		
<i>platypterus</i> . . . . .	—	—	—	+	—	—
<i>Archibuteo lagopus sancti-johannis</i> . . . . .	—	—	—	+	—	—
<i>Aquila chrysaetos</i> . . . . .	+	—	—	+	—	—
<i>Haliaeetus leucocephalus</i> . . . . .	+	—	—	+	—	—
<i>Falco islandus</i> . . . . .	—	?		?		
<i>rusticolus</i> . . . . .	—					
<i>gyrfalco</i> . . . . .	—		?	—	—	
<i>rusticolus obsoletus</i> . . . . .	+	—		—	+	
<i>peregrinus anatum</i> . . . . .	—	—	—	+	—	—
<i>columbarius</i> . . . . .	—	—	—	+	—	—
<i>timunculus</i> . . . . .	—			—		
<i>sparverius</i> . . . . .	—	—	—	+	—	—
<i>Pandion haliaetus carolinensis</i> . . . . .	—	—	—	+	—	—
<i>Aluco pratincola</i> . . . . .			—	+	—	—
<i>Asio wilsonianus</i> . . . . .	—	—	—	+	+	—
<i>flammeus</i> . . . . .	—	—	—	+	—	—
<i>Strix varia</i> . . . . .	+	—	—	+	—	—
<i>Scotiaptex nebulosa</i> . . . . .	+	—	+	—	—	—
<i>Glaux funerea richardsoni</i> . . . . .	+	—	—	—	—	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Glaux acadicus</i> . . . . .	—	—	—	+	—	—
<i>Otus asio</i> . . . . .	—	—	—	+	—	—
<i>Bubo virginianus</i> . . . . .	—	—	—	+	—	—
<i>subarcticus</i> . . . . .	—			—		
<i>heteroenemis</i> . . . . .	—					
<i>Nyctea nyctea</i> . . . . .	—	—	—	+	—	—
<i>Surnia ulula caparoch</i> . . . . .	+	+	—	—	—	—
<i>Speotyto cunicularia hypogaea</i> . . . . .				—		
<i>Coccyzus americanus</i> . . . . .	—	—	—	+	—	—
<i>erythrophthalmus</i> . . . . .	—	—	—	+	—	—
<i>Ceryle alcyon</i> . . . . .	—	—	—	+	—	—
<i>Dryobates villosus</i> . . . . .	+	—	—	+	—	—
<i>pubescens medianus</i> . . . . .	—	—	—	+	—	—
<i>Picoides arcticus</i> . . . . .	+	—	—	—		—
<i>americanus</i> . . . . .	+	—	—	—		—
<i>Sphyrapicus varius</i> . . . . .	—	—	—	+	—	+
<i>Phloeotomus pileatus abieticola</i> . . . . .	+	+	—	—		—
<i>Melanerpes erythrocephalus</i> . . . . .	—	—	—	+	—	—
<i>Centurus carolinus</i> . . . . .				—	+	—
<i>Colaptes auratus luteus</i> . . . . .	—	—	—	+	—	—
<i>Antrostomus carolinensis</i> . . . . .				—		—
<i>vociferus</i> . . . . .	—	—	—	+	—	—
<i>Chordeiles virginianus</i> . . . . .	—	—	—	+	—	—
<i>Chaetura pelagica</i> . . . . .	—	—	—	+	—	—
<i>Trochilus colubris</i> . . . . .	—	—	—	+	—	—
<i>Muscivora forficata</i> . . . . .			—			—
<i>Tyrannus tyrannus</i> . . . . .	—	—	—	+	—	+
<i>dominicensis</i> . . . . .				—		
<i>verticalis</i> . . . . .	—					
<i>Myiarchus crinitus</i> . . . . .	—	—	—	+	—	—
<i>Sayornis phoebe</i> . . . . .	—	—	—	+	—	—
<i>saya</i> . . . . .				—		
<i>Nuttallornis borealis</i> . . . . .	—	—	—	+	—	—
<i>Myiochanes virens</i> . . . . .	—	—	—	+	—	—
<i>Empidonax flaviventris</i> . . . . .	+	—	—	+	—	—
<i>virescens</i> . . . . .			—	—		—
<i>traillii alnorum</i> . . . . .	—	+	—	+		—
<i>minimus</i> . . . . .	—	—	—	+	—	—
<i>Otocoris alpestris</i> . . . . .	—	—	—	+	—	—
<i>praticola</i> . . . . .	—	—	—	+	—	—
<i>hoyti</i> . . . . .						—
<i>Cyanocitta cristata</i> . . . . .	—	—	—	+	—	—
<i>Perisoreus canadensis</i> . . . . .	+	—	—	—		—
<i>Corvus corax principalis</i> . . . . .	+	—	—	—		—

<i>Corvus brachyrhynchos</i>	—	—	—	+	—	—
<i>ossifragus</i>			?	—		—
<i>Sturnus vulgaris</i>	—	—	—	—	—	—
<i>Dolichonyx oryzivorus</i>	—	—	—	+	—	+
<i>Molothrus ater</i>	—	—	—	+	—	—
<i>Xanthocephalus xanthocephalus</i>	—	—	—	—	—	—
<i>Agelaius phoeniceus</i>	—	—	—	—	+	—
<i>fortis</i>						—
<i>Sturnella magna</i>	—	—	—	+	—	—
<i>Icterus spurius</i>	—	—	—	—	+	+
<i>galbula</i>	—	—	—	+	—	—
<i>bullocki</i>	—					
<i>Euphagus carolinus</i>	—	—	—	+	—	—
<i>Quiscalus quiscula</i>			?	—	—	—
<i>aeneus</i>	—	—	—	+	—	—
<i>Hesperiphona vespertina</i>	—	—	—	+		—
<i>Pipilo erythrophthalmus</i>	—	—	—	+	—	—
<i>Carpodacus purpureus</i>	—	—	—	+	—	—
<i>Loxia curvirostra minor</i>	—	—	—	+	—	—
<i>leucoptera</i>	—	+	—	+	—	—
<i>Acanthis hornemannii exilipes</i>	—	—	?	+		
<i>linaria</i>	—	—	—	+	—	—
<i>holboellii</i>	—			—		
<i>rostrata</i>	—	?		+	—	—
“ <i>brewsteri</i> ”				—		
<i>Astragalinus tristis</i>	—	—	—	+	—	—
<i>Spinus pinus</i>	—	—	—	+	—	—
<i>pinus</i>				—		
<i>Plectrophenax nivalis</i>	—	—	—	+	—	—
<i>Calcarius lapponicus</i>	—	—	—	+	—	—
<i>ornatus</i>	—			+		
<i>Poocetes gramineus</i>	—	—	—	+	—	—
<i>Passerculus princeps</i>	—	—		+	—	—
<i>sandwichensis savanna</i>	—	—	—	+	—	—
<i>Coturniculus savannarum australis</i>	—	—	—	+	—	—
<i>Ammodramus henslowii</i>				—	—	—
<i>caudacutus</i>	—	—	?	+	—	—
<i>nelsoni</i>	—			—		—
<i>subvirgatus</i>	—			+	—	—
<i>maritimus</i>	—			+	+	—
<i>Chondestes grammacus</i>				—		
<i>Zonotrichia leucophrys</i>	—	—	—	+	—	—
<i>albicollis</i>	—	—	—	+	—	—
<i>Spizella monticola</i>	—	—	—	+	—	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Spizella passerina</i> . . . . .	—	—	—	+	—	—
<i>breweri</i> . . . . .				—		
<i>pusilla</i> . . . . .	—	—	—	+	—	—
<i>Junco hyemalis</i> . . . . .	+	—	—	+	—	—
<i>montanus</i> . . . . .				—		
<i>Melospiza melodia</i> . . . . .	—	—	—	+	—	—
<i>lincolni</i> . . . . .	—	—	—	+	+	+
<i>georgiana</i> . . . . .	—	—	—	+	—	+
<i>Passerella iliaca</i> . . . . .	—	—	—	+	—	—
<i>Pipilo erythrophthalmus</i> . . . . .	—	—	—	+	—	—
<i>Cardinalis cardinalis</i> . . . . .	—	—	—	—	—	—
<i>Zamelodia ludoviciana</i> . . . . .	+	—	—	+	—	—
<i>Guiraca caerulea</i> . . . . .		?	?	—	—	—
<i>Passerina cyanea</i> . . . . .	+	—	—	+	—	—
<i>Spiza americana</i> . . . . .	—			—	—	—
<i>Calamospiza melanocorys</i> . . . . .				—		
<i>Piranga ludoviciana</i> . . . . .	—			—		—
<i>erythromelas</i> . . . . .	—	—	—	+	—	—
<i>rubra</i> . . . . .	—	?	?	—	+	—
<i>Progne subis</i> . . . . .	—	—	—	+	—	—
<i>Petrochelidon lunifrons</i> . . . . .	—	—	—	+	—	—
<i>Hirundo erythrogaster</i> . . . . .	—	—	—	+	—	—
<i>Iridoprocne bicolor</i> . . . . .	—	—	—	+	—	—
<i>Riparia riparia</i> . . . . .	—	—	—	+	—	—
<i>Stelgidopteryx serripennis</i> . . . . .				—		+
<i>Bombycilla garrula</i> . . . . .	—	—	—	—		—
<i>cedrorum</i> . . . . .	—	—	—	+	—	—
<i>Lanius borealis</i> . . . . .	—	—	—	+	—	—
<i>ludovicianus</i> . . . . .				+		—
<i>migrans</i> . . . . .	+	—	+	+	—	—
<i>Vireosylva olivacea</i> . . . . .	—	—	—	+	—	—
<i>philadelphica</i> . . . . .	—	—	—	—	—	—
<i>gilva</i> . . . . .	—	—	—	+	—	—
<i>Lanivireo flavifrons</i> . . . . .	—	—	—	+	—	—
<i>solitarius</i> . . . . .	+	—	—	+	—	—
<i>Vireo griseus</i> . . . . .	?	?	—	+	—	—
<i>bellii</i> . . . . .				—		—
<i>Mniotilta varia</i> . . . . .	—	—	—	+	—	—
<i>Protonotaria citrea</i> . . . . .	—		?	+	—	
<i>Helminthophila vermivorus</i> . . . . .	?	?	—	—		+
<i>Helminthophila pinus</i> . . . . .			?	+	—	+
<i>chrysoptera</i> . . . . .			—	+	—	—
<i>rubricapilla</i> . . . . .	—	—	—	+	—	—
<i>celata</i> . . . . .	?	—		+	—	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Helminthophila peregrina</i> . . . . .	+	—	—	+	—	—
<i>Compsothlypis americana usneae</i> . . . . .	—	—	—	+	—	—
<i>Dendroica tigrina</i> . . . . .	+	—	—	+	—	—
<i>aestiva</i> . . . . .	—	—	—	+	—	—
<i>caerulescens</i> . . . . .	—	—	—	+	—	—
<i>coronata</i> . . . . .	—	—	—	+	—	—
<i>auduboni</i> . . . . .	—	—	—	—	—	?
<i>magnolia</i> . . . . .	—	—	—	+	—	—
<i>cerulea</i> . . . . .	—	—	—	—	—	—
<i>pennsylvanica</i> . . . . .	—	—	—	+	—	+
<i>striata</i> . . . . .	—	—	—	+	—	—
<i>blackburniae</i> . . . . .	+	—	—	+	—	—
<i>castanea</i> . . . . .	—	—	—	+	—	—
<i>dominica</i> . . . . .	—	—	—	—	—	—
<i>virens</i> . . . . .	—	—	—	+	—	+
<i>vigorsii</i> . . . . .	—	—	—	+	—	—
<i>palmarum</i> . . . . .	—	—	—	—	—	—
<i>hypochrysea</i> . . . . .	+	—	—	+	—	—
<i>discolor</i> . . . . .	—	—	—	+	—	—
<i>Seiurus aurocapillus</i> . . . . .	—	—	—	+	—	—
<i>Seiurus noveboracensis</i> . . . . .	—	—	—	+	—	—
<i>notabilis</i> . . . . .	—	—	—	—	—	—
<i>motacilla</i> . . . . .	—	?	?	+	—	—
<i>Oporornis formosa</i> . . . . .	—	—	—	—	—	—
<i>agilis</i> . . . . .	—	—	—	+	—	—
<i>philadelphia</i> . . . . .	+	+	—	—	—	—
<i>Geothlypis trichas</i> . . . . .	—	—	—	+	—	—
<i>Icteria virens</i> . . . . .	—	?	—	+	—	—
<i>Wilsonia citrina</i> . . . . .	—	—	?	—	+	+
<i>pusilla</i> . . . . .	—	—	—	+	—	—
<i>canadensis</i> . . . . .	—	—	—	+	—	+
<i>Setophaga ruticilla</i> . . . . .	—	—	—	+	—	—
<i>Anthus rubescens</i> . . . . .	—	—	—	+	—	—
<i>Mimus polyglottos</i> . . . . .	—	—	—	+	—	—
<i>Dumetella carolinensis</i> . . . . .	—	—	—	+	—	—
<i>Toxostoma rufum</i> . . . . .	—	—	—	+	—	—
<i>Thryothorus ludovicianus</i> . . . . .	—	—	?	+	—	—
<i>Thryomanes bewickii</i> . . . . .	?	—	—	—	—	—
<i>Troglodytes aedon</i> . . . . .	—	—	—	+	—	—
<i>Namus hiemalis</i> . . . . .	—	—	—	+	—	—
<i>Cistothorus stellaris</i> . . . . .	—	—	—	+	—	—
<i>Telmatodytes palustris</i> . . . . .	—	—	—	+	—	—
<i>Certhia familiaris americana</i> . . . . .	—	—	—	+	—	—
<i>Sitta carolinensis</i> . . . . .	—	—	—	+	—	—

	Me.	N.H.	Vt.	Mass.	R. I.	Conn.
<i>Sitta canadensis</i> . . . . .	—	—	—	+	—	—
<i>Baeolophus bicolor</i> . . . . .	—	?	?			—
<i>Penthestes atricapillus</i> . . . . .	—	—	—	+	—	—
<i>hudsonicus</i> . . . . .	+	—	—	—	—	—
<i>Regulus satrapa</i> . . . . .	—	—	—	+	—	—
<i>calendula</i> . . . . .	—	—	—	+	—	—
<i>Polioptila caerulea</i> . . . . .	—	?	?	+	—	—
<i>Hyalocichla mustelina</i> . . . . .	—	—	—	+	—	—
<i>fuscescens</i> . . . . .	—	—	—	+	—	—
<i>salicicola</i> . . . . .				—	—	—
<i>aliciae</i> . . . . .	—	—	—	+	—	—
<i>bicknelli</i> . . . . .	—	—	—	—	—	—
<i>ustulata swainsonii</i> . . . . .	—	+	—	+	—	—
<i>guttata pallasii</i> . . . . .	—	—	—	+	—	—
<i>Planesticus migratorius</i> . . . . .	—	—	—	+	—	—
<i>Ixoreus naevius</i> . . . . .				+		
<i>Sialia sialis</i> . . . . .	—	—	—	+	—	—







## AVES.

PYGOPODES.

COLYMBI

COLYMBIDAE.

### 1. COLYMBUS HOLBOELLII (Reinhardt) Ridgway.

**Holboell's grebe; Red-necked grebe.**

*Podiceps holböllii* Reinh., Vidensk. meddel. naturh. forh. Kjöbenhavn for 1853, 1854, p. 76. "GRÖNLAND. . . . JULIANEHAABS DISTRIKT."

*Podiceps rubricollis* Lath. Audubon, Birds of Amer., 1844, vol. 7, p. 312 pl. 480.

Fresh and salt water; nests among rushes in sloughs.

ME.—Uncommon winter resident coastwise. Sept.—April 24.

N. H.—Uncommon migrant and coastwise winter resident. Oct.—May (June).

Vt.—Rare migrant: Middlebury; Wells River; Windsor, spring.

MASS.—Uncommon migrant and winter resident, mainly coastwise. (Sept.) Oct. 10—May 24.

R. I.—Uncommon winter resident coastwise. Oct. 23—May 2.

CONN.—Uncommon winter resident coastwise. Oct. 19—spring.

### 2. COLYMBUS AURITUS Linné.

**Horned grebe; Devil diver; Hell diver; Little diver; Tinker; Tinker loon.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 135. "IN SUMMIS EUROPAE & AMERICA LACUBUS."

*Podiceps cornutus* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 316, pl. 481. Egg, Capen, 1886, pl. 24, fig. 3.

Salt and fresh water; nests among rushes in marshes.

## PYGOPODES.

## COLYMBI.

## COLYMBIDAE.

ME.—Common migrant, mainly coastwise; rare winter, and in northeastern part, rare summer resident.

N. H.—Uncommon migrant and winter resident, mainly coastwise. Oct.—April.

VT.—Rare migrant.

MASS.—Common migrant and winter resident, mainly coastwise. Oct. 1—May 6 (June 1, Springfield).

R. I.—Common winter resident coastwise. Oct. 23—April 24 (May).

CONN.—Common winter resident coastwise; accidental in summer (Melrose, July 26, 1882). Sept.—May (July 26).

3. *PODILYMBUS PODICEPS* (Linné) Lawrence.

**Pied-billed grebe; Dabchick; Devil diver; Diedapper; Dipper; Frog-in-throat; Hell diver; Water witch.**

Chapman, Handb. birds east. No. Amer., 1895, p. 57; Bird-life, 1898, pl. 2. Egg, Capen, 1886, pl. 24, fig. 5.

*Colymbus podiceps* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 136. "IN AMERICA SEPTENTRIONALI."

Freshwater rivers, ponds, swamps; nests among reeds in swamps.

ME.—Common migrant, mainly coastwise; less common summer resident. April—May 19; summer; Sept. 1—Nov. 13 (late Nov.). Eggs, June 6.

N. H.—Uncommon migrant and rare summer resident. Mar.—Nov.

VT.—Rare summer resident. Spring—Oct. 19.

MASS.—Rare spring, common fall migrant, and uncommon local summer resident; accidental in winter. (Feb.) Mar. 1—Dec. 10. Eggs, April 23—June 28.

R. I.—Uncommon migrant and summer resident. April—Oct. 22. Eggs, June 6.

CONN.—Common migrant, mainly in fall, and rare summer resident. April—summer; Sept.—Nov. (Dec. 25, 1908, South Norwalk).

PYGOPODES.

CEPHAL.

## GAVIIDAE.

4. *GAVIA IMMER* (Brünnich) A. O. U. comm.

**Loon; Big loon; Call-up-a-storm; Great northern diver.**

*Colymbus immer* Brünn., Ornith. borealis, 1764, p. 38. "E FAEROA."

*Urinator iuber* (Gummerus). Chapman, Handb. birds east. No. Amer., 1895, p. 58; Bird-life, 1898, pl. 3. Egg, Capen, 1886, pl. 24, fig. 2.

Salt and fresh water; nests on shores of lakes.

ME., N. H.—Common migrant, less common winter and summer resident. Eggs, June 11–19 (July).

VT.—Uncommon summer resident.

MASS.—Common migrant and winter resident, mainly coastwise; rare summer resident inland (formerly Springfield); a few non-breeding birds summer offshore.

R. I.—Common migrant and winter resident coastwise; a few non-breeding birds summer offshore. Sept. 15–May 28; summer.

CONN.—Common winter resident coastwise; formerly a rare summer resident (Easthampton). Oct.–April 29.

5. *GAVIA ARCTICA* (Linné) Allen.

**Black-throated loon or diver.**

*Colymbus arcticus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 135. "IN EUROPA & AMERICA BOREALI." Audubon, Birds of Amer., 1844, vol. 7, p. 295, pl. 477. Egg, Seebohm, 1896, p. 78, pl. 21, fig. 1.

Seacoasts; nests on the ground.

ME.—Rare winter visitor: China, Kennebec Co., April 19, 1899; Cutler, Dec. 8, 1881.

MASS.—Doubtfully recorded.

CONN.—Doubtfully recorded (? near Southbridge, Mass.).

PYGOPODES.

CEPPHIL.

GAVIIDAE.

6. *GAVIA STELLATA* (Pontoppidan) A. O. U. comm.

**Red-throated loon; Cape-race; Little loon; Red-throated diver;  
Scape-grace.**

*Colymbus stellatus* Pontopp., Danske atlas, 1763, vol. 1, p. 621.  
(Based on Willoughby.)

*Colymbus septentrionalis* Linné. Audubon, Birds of Amer., 1844,  
vol. 7, p. 299, pl. 478. Egg, Seebohm, 1896, p. 78, pl. 21, fig. 2.

Salt water mainly, sometimes on lakes; nests on the ground.

ME., N. H.—Common migrant and winter resident, mainly  
coastwise. (Aug. 16) Sept. 22—(June 22).

VT—Rare migrant. Nov. 1.

MASS.—Uncommon spring, common fall migrant and winter  
resident, mainly coastwise; casual in summer. Aug. 27—April 8  
(July 2).

R. I.—Common migrant and winter resident coastwise; a few  
non-breeding birds summer offshore. Oct. 1—April 27 (summer).

CONN.—Common winter resident, coastwise. Oct.—May.

ALCIDAE.

7. *LUNDA CIRRHATA* (Pallas) Pallas.

**Tufted puffin.**

*Alca cirrhata* Pallas, Spic. zool., 1769, fasc. 5, p. 7, pl. 1; pl. 2,  
figs. 1-3. "IN MARI INTER KAMTSCHATKAM ET AMERICAM ARCHI-  
PELAGUMGUE KURILUM."

*Mormon cirrhatus* Lath. Audubon, Birds of Amer., 1844, vol. 7,  
p. 234, pl. 462.

Rocky sea coasts; nests in burrows.

ME.—Accidental visitor: mouth of Kennebec River, winter of  
1831-32.

PYGPODES.

CEPPIII

ALCIDAE

## 8. FRATERCULA ARCTICA (Linné) Schäffer.

**Puffin; Paroquet; Sea parrot.**

*Alca arctica* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 130. "IN EUROPAE BOREALIS OCEANO."

*Mormon arcticus* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 238, pl. 464. Egg, Capen, 1886, pl. 24, fig. 6.

Rocky seacoasts; nests in burrows.

ME.—Common winter resident coastwise, and rare summer resident (Matineus Rock and Machias Seal Island).

N. H.—Uncommon winter visitor (at Isles of Shoals).

MASS.—Uncommon winter visitor coastwise. Oct. 16–March.

R. I.—Doubtfully reported.

## 9. CEPHUS GRYLLE (Linné) Brehm.

**Black guillemot; Pigeon murre; Sea pigeon.**

*Alca grylle* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 130. "IN EUROPAE BOREALIS OCEANO."

*Uria grylle* (Linné). Audubon, Birds of Amer., 1844, vol. 7, p. 272, pl. 474. Egg, Capen, 1886, pl. 24, figs. 7, 8.

Rocky seacoasts; nests in deep crevices of rocks.

ME.—Common winter resident coastwise; less common summer resident from Knox Co. eastward. Eggs, June 15–23.

N. H.—Uncommon winter visitor coastwise.

MASS.—Uncommon winter visitor coastwise. Sept.–April 19 (May).

R. I.—Rare winter visitor. Nov. 18–April.

CONN.—Rare winter visitor coastwise: Stony Creek, Dec., 1887.

PYGOPODES.

CEPPHI.

ALCIDAE.

10. *URIA TROILE* (Linné) Latham.**Murre; Foolish guillemot; Guillemot.**

COUES, Key No. Amer. birds, ed. 5, 1903, vol. 2, p. 1082, figs. 742, 743.

*Colymbus troille* (sic) Linné, Fauna Suec., ed. 2, 1761, p. 52. "IN ALTO MARI SEPTENTRIONALI."

*Alca troile* (Linné). Dresser, Birds of Europe, 1871-81, vol. 8, pl. [621]. Egg, Capen, 1886, pl. 24, fig. 9; pl. 25, fig. 1-3.

Rocky seacoasts; nests on sea cliffs.

ME.—Rare winter visitor: Eastport, Dec. 27, 1886; Scarborough.

VT.—Doubtfully recorded.

MASS.—Rare winter visitor: "Massachusetts" (Society collection).

R. I.—Doubtfully recorded.

CONN.—? Rare winter visitor (Merriam, probably in error for *U. lomvia*).

11. *URIA LOMVIA* (Linné) Bryant.**Brünnich's murre; Brünnich's guillemot; Ice-bird; Large-billed guillemot; Sea crow.**

*Alca lomvia* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 130. "IN EUROPA BOREALI."

*Uria brunnichii* Sabine. Audubon, Birds of Amer., 1844, vol. 7, p. 265, pl. 472. Egg, Seebohm, 1896, p. 94, pl. 26, figs. 5, 6.

Rocky seacoasts; nests on sea cliffs.

ME.—Common winter visitor coastwise, casual inland; a doubtful breeding record (Penobscot Bay, M. Hardy). Nov. 23-Mar.

N. H.—Irregular winter visitor coastwise; occasional inland. Nov. 25-Mar.



PYGOPODES.

CEPPII.

## ALCIDAE.

Vt.—Irregular winter visitor. Nov. Jan.

MASS.—Uncommon, irregular winter visitor coastwise; occasional inland. Nov. 21–Mar. 14 (April 15).

R. I.—Uncommon and irregular winter visitor. Nov. 15–Feb. 23.

CONN.—Irregular winter visitor, sometimes common. Nov. 18–Feb. 16.

12. *ALCA TORDA* Linné.

**Razor-billed auk; Razor-bill; Tinker murre; Turre.**

Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 130. “IN EUROPAE BOREALIS OCEANO.” Audubon, *Birds of Amer.*, 1844, vol. 7, p. 247, pl. 466. Egg, Capen, 1886, pl. 25, fig. 4–6.

Rocky seacoasts; nests in rock crevices.

ME.—Common winter visitor coastwise; said to have nested formerly.

N. H.—Uncommon winter visitor coastwise. Nov.–Mar.

MASS.—Uncommon winter visitor coastwise. Nov. 1–April 15.

R. I.—Rare winter visitor coastwise. Nov. 15–Feb. 4.

CONN.—Rare winter visitor coastwise.

13. *PLAUTUS IMPENNIS* (Linné) Steenstrup.

**Great auk; Garefowl.**

*Alca impennis* Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 130. “IN EUROPA ARCTICA.” Audubon, *Birds of Amer.*, 1844, vol. 7, p. 245, pl. 465. Egg, S. Grieve, *Great Auk, or Garefowl*, London, 1885, pls. 1, 2 (opp. pp. 108, 109).

Formerly a coastwise species, breeding at Funk Id., near Newfoundland, and at a few places on the European coast; last specimen killed in 1844.

## PYGOPODES.

## CEPHI.

## ALCIDAE.

ME., N. H., MASS.—Formerly a winter visitor coastwise.

R. I., CONN.—Probably occurred formerly as a winter visitor coastwise (bones in shell heaps at Block Id., R. I.).

## 14. ALLE ALLE (Linné) Stejneger.

**Dovekie; Ice-bird; Knotty; Little auk; Pine-knotty; Rotche; Sea dove.**

*Alca alle* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 131. "IN EUROPAE, AMERICAЕ ARCTICAЕ OCEANO."

*Mergulus alle* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 257, pl. 469. Egg, Seebohm, 1896, p. 95, pl. 26, fig. 3.

Rocky coasts; nests on ledges or in crevices of sea cliffs.

ME.—Uncommon winter visitor coastwise, casual inland; Late Nov.—Mar. 15.

N. H.—Uncommon winter visitor coastwise; casual inland.

VT.—Accidental winter visitor.

MASS.—Uncommon winter visitor, coastwise; casual inland. (Sept.) Nov. 4—May 1.

R. I.—Rare and irregular winter visitor coastwise; casual inland. Nov. 14—April 27.

CONN.—Rare and irregular winter visitor coastwise; casual inland. (Sept.) Nov. 10.

## LONGIPENNES.

## STERCORARIIDAE.

## 15. MEGALESTRIS SKUA (Brünnich) Ridgway.

**Skua; Great Skua; Sea-hen.**

*Catharacta skua* Brünn., Ornith. borealis, 1764, p. 33. "E FEROA & ISLANDIA."

*Megalestris catarrhactes* (Linné). Saunders, Cat. birds British mus., 1896, vol. 25, p. 315.

## LONGIPENNES.

## STERCORARIIDAE.

*Stercorarius catarrhactes* (Linné). Dresser, Birds of Europe, 1871-81, vol. 8, pl. [609]. Egg, Seebohm, 1896, p. 118, pl. 37, figs. 3, 4.

Pelagic; nests on rocky cliffs or tundras.

MASS.—Rare visitor offshore: Georges Banks, July, 1878; Nantucket Shoals, Oct. 11, 1883; Polluck Rip, Sept. 10, 1884.

## 16. STERCORARIUS POMARINUS (Temminck) Vieillot.

**Pomarine jaeger.**

*Lestris pomarinus* Temm., Man. d'ornith., 1815, p. 514. "LES REGIONS DU CERCLE ARCTIQUE." Audubon, Birds of Amer., 1844, vol. 7, p. 186, pl. 451. Egg, Seebohm, 1896, p. 116, pl. 37, fig. 6.

Largely marine; nests on the ground.

ME.—Uncommon migrant, casual in summer. May; (June 23, 24); July 14-late Oct.

MASS.—Uncommon migrant, occasional in summer. May 23; July 5-Sept. 28.

R. I.—Uncommon migrant offshore, mainly in fall. Oct.

## 17. STERCORARIUS PARASITICUS (Linné) Schäffer.

**Parasitic jaeger; Gull chaser; Marlinspike; Richardson's jaeger; Robber gull; Sea hen.**

*Larus parasiticus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 136. "INTRA TROPICUM CANCRI, EUROPAE, AMERICAE, ASIAE."

*Lestris richardsonii* Swains. Audubon, Birds of Amer., 1844, vol. 7, p. 190, pl. 452. Egg, Seebohm, 1896, p. 116, pl. 37, figs. 1, 2.

Largely marine; nests on the ground.

ME.—Uncommon migrant offshore; casual in summer. May-June; (July) Aug.-Oct.

## LONGIPENNES.

## STERCORARIIDAE.

N. H.— Uncommon migrant offshore. Sept.

MASS.— Uncommon migrant and summer visitor offshore; rare in winter. May 31–Oct. 14 (Jan., Feb.).

R. I.— Rare migrant offshore: Charlestown Beach, three, Sept. 2, 1897.

CONN.— Rare migrant offshore: Bridgeport; Portland, fall, 1875.

## 18. STERCORARIUS LONGICAUDUS Vieillot.

**Long-tailed jaeger; Buffon's jaeger or skua; Marlinspike; Whiptail.**

Vieill., Nouv. dict. d'hist. nat., 1819, vol. 32, p. 157. "LE NORD DE L'EUROPE, DE L'ASIE ET DE L'AMÉRIQUE."

*Lestrís parasiticus* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 192, pl. 453. Egg, Seebohm, 1896, p. 117, pl. 37, fig. 5.

Largely marine; nests on the ground.

ME.— Rare migrant offshore.

VT.— Accidental migrant: West Castleton, Sept. 7, 1877.

MASS.— Rare migrant. Spring; Aug. 24–Sept. 29.

CONN.— Rare or accidental visitor: Community Lake, Wallingford, Aug. 30, 1873.

## LARIDAE.

## 19. PAGOPHILA ALBA (Gunnerus) Coues.

**Ivory gull.**

*Larus albus* Gumm., in Leem's Beskriv. Finmark. Lapp., 1767, p. 285 (note). NORTHERN NORWAY.

*Larus eburneus* Gmel. Audubon, Birds of Amer., 1844, vol. 7 p. 150, pl. 445. Egg, Seebohm, 1896, p. 115, pl. 32, fig. 3.

Arctic seas, following the pack ice; nests on rocky cliffs.

## LONGIPENNES.

## LARIDAE.

ME.—The Boardman record refers to a New Brunswick specimen.

MASS.—Accidental winter visitor: Monomoy Id., Dec. 1, 1886.

20. *RISSA TRIDACTYLA* (Linné) Bonaparte.

**Kittiwake; Frost-bird; Jack gull; Meterick; Pinny owl or Pinyole; Snow gull; Winter gull.**

*Larus tridactylus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 136. "IN EUROPA SEPTENTRIONALI." Audubon, Birds of Amer., 1844, vol. 7, p. 146, pl. 444. Egg, Samuels, Ornith. and ool. N. E., 1867, pl. 4, fig. 3.

Maritime; nests on sea cliffs.

ME.—Common fall and winter visitor offshore; casual inland, and in summer. Nov.—April (June; July 11, 1903, Jericho Bay, barren bird; July 14, 1907, off Portland).

N. H.—Common fall and winter visitor offshore. Nov. 1—spring.

VT.—Rare winter visitor: Enosburg Falls, Nov. 12, 1906; St. Albans; St. Johnsbury.

MASS.—Common fall and winter visitor offshore, casual inland. Sept. 6—Mar. 10.

R. I.—Common fall and winter visitor offshore. Nov.—Mar. 23.

CONN.—Uncommon fall and winter visitor offshore. Fall—April 14.

21. *LARUS HYPERBOREUS* Gummerus.

**Glaucous gull; Burgomaster; Owl gull; White minister.**

Gummerus, in Leem's Beskriv. Finnmark. Lapp., 1767, p. 226 (note), p. 283 (note). No locality.

*Larus glaucus* Brünn. Audubon, Birds of Amer., 1844, vol. 7, p. 170, pl. 449. Egg, Seebohm, 1896, p. 112, pl. 32, figs. 4, 5.

Seacoasts; nests on cliffs.

## LONGIPENNES.

## LARIDAE.

ME.—Rare winter visitor. Late Nov.—April 27.

N. H.—Rare winter visitor: Hampton, about Feb. 20, 1886, and in May.

VT.—Doubtfully recorded.

MASS.—Rare winter visitor. Nov. 29—April 23 (May 26, 1907, Ipswich).

R. I.—Rare winter visitor: Newport, Mar. 12, 1900 (three); Dec. 1901.

## 22. LARUS LEUCOPTERUS Faber.

**Iceland gull; White-winged gull.**

Faber, Prodr. Isl. ornith., 1822, p. 91. "IM WINTER. . . NACH ISLAND." Saunders, Cat. birds British mus., 1896, vol. 25, p. 296. Audubon, Birds of Amer., 1844, vol. 7, pl. 447. Egg, Seebohm 1896, p. 113, pl. 33, fig. 3.

Seacoasts; nests on the ground.

ME.—Rare winter visitor. Nov.—April.

MASS.—Rare winter visitor. Nov. 4—Mar. 20.

CONN.—Rare winter visitor: Stamford, Feb. 16, 1894.

## 23. LARUS KUMLIENI Brewster.

**Kumlien's gull; Lesser glaucous-winged gull.**

Brewst., Bull. Nuttall ornith. club, 1883, vol. 8, p. 216. "CUMBERLAND SOUND, ARCTIC AMERICA." Dwight, Auk, 1906, vol. 23, p. 36—41, pl. 1, figs.

Coastwise; breeds on Arctic coasts.

ME.—Probably occasional winter visitor: near Eastport (may have been N. B.); "seen" in Portland Harbor and at Scarborough.

MASS.—Rare winter visitor: Boston Harbor, Feb. 22, 1905;

## LONGIPENNES.

## LARIDAE.

Plymouth, Jan. 5, 1888 (Dwight, immature); Boston and Swampscott, two or more, Dec. 29, 1907–Mar. 13, 1908.

CONN.—Rare winter visitor: Stamford, Feb. 16, 1894 (Dwight, immature).

## 24. LARUS MARINUS Linné.

**Great black-backed gull; Black-back; Black minister; Coffin carrier; Ghost-bird; Minister; Saddle-back.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 136. "IN EUROPA." Audubon, Birds of Amer., 1844, vol. 7, p. 172, pl. 450. Egg, Capen, 1886, pl. 22, figs. 2, 3.

Seacoasts; nests on the ground.

ME.—Common migrant and winter resident; casual summer visitor. Sept. 21–April (summer).

N. H.—Common winter resident coastwise. Oct. 15–spring.

MASS.—Common winter resident coastwise; a few non-breeding birds summer. Late Aug.–May 1 (summer).

R. I.—Common winter resident coastwise. Aug. 26–April 23.

CONN.—Uncommon winter resident coastwise. Fall–Mar.

## 25. LARUS ARGENTATUS Pontoppidan.

**Herring gull; Gray gull; Sea gull; Winter gull.**

Pontopp., Danske atlas, 1763, vol. 1, p. 622. No locality. Audubon, Birds of Amer., 1844, vol. 7, p. 163, pl. 448. Egg, Capen, 1886, pl. 22, fig. 4–6.

Seacoasts, large rivers and lakes; nests on the ground or rarely in trees.

ME.—Abundant resident, breeding mainly on outer islands, rarely on inland lakes. Eggs, (May 14) June 12–July 18 (Aug. 19).

## LONGIPENNES.

## LARIDAE.

N. H.—Common migrant and winter resident; less common summer visitor coastwise.

Vt.—Uncommon migrant and rare summer resident (Lake Champlain).

MASS.—Common migrant and winter resident, less common summer visitor coastwise; not now known to breed (nested at Wepecket Id., 1882, Mackay).

R. I.—Common migrant and winter resident, mainly coastwise; a few non-breeding birds summer. Sept. 10–May 22; summer.

CONN.—Common migrant and winter resident, mainly coastwise; a few non-breeding birds summer. Oct.–May; summer.

## 26. LARUS DELAWARENSIS Ord.

**Ring-billed gull.**

Ord, Guthrie's Geogr., hist., and commercial grammar, 2d Amer. ed., 1815, vol. 2, p. 319. "ON THE DELAWARE BELOW PHILADELPHIA." Chapman, Handb. birds east. No. Amer., 1895, p. 73.

*Larus zonorhynchus* Richardson. Audubon, Birds of Amer., 1844, vol. 7, p. 152, pl. 446.

Seacoasts; nests on the ground.

ME., N. H.—Uncommon migrant and rare winter resident coastwise. Mar.–April; Oct.–Nov. (winter).

MASS.—Uncommon migrant coastwise. Spring; July 17–Oct. 30.

R. I.—Uncommon migrant and winter resident. Nov.–April.

CONN.—Uncommon migrant and winter resident.



## LONGIPENNES.

## LARIDAE.

27. *LARUS ATRICILLA* Linné.**Laughing gull; Black-headed gull; Black-poll.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 136. "AMERICA."  
Audubon, Birds of Amer., 1844, vol. 7, p. 136, pl. 413. Egg,  
Capen, 1886, pl. 23, fig. 1-3.

Seacoasts; nests on the ground.

ME.—Rare summer resident (Metinic Green Id., Western Egg  
Rock). (Late April)–Sept. Eggs, June 30–July 5.

VT.—Rare visitor: Bristol, May, 1883; St. Albans.

MASS.—Uncommon summer resident south of Cape Cod (Mus-  
keget Id.; formerly at Egg Rock, Nahant); elsewhere a rare visitor.  
May 7–Oct. 1 (late Nov.). Eggs, June 7–July 10.

R. I.—Rare summer visitor: Seakonnet Point, Sept., 1884; Sea-  
konnet Point, Aug. 24, 1903.

CONN.—Rare summer visitor; may breed rarely.

28. *LARUS PHILADELPHIA* (Ord) Gray.**Bonaparte's gull.**

*Sterna philadelphia* Ord, Guthrie's Geogr., hist., and commer-  
cial grammar, 2d Amer. ed., 1815, vol. 2, p. 319. No locality;  
probably PHILADELPHIA.

*Larus bonapartii* Richardson. Audubon, Birds of Amer., 1844,  
vol. 7, p. 131, pl. 442. Egg, Seebohm, 1896, p. 106, pl. 36, fig. 7.

Salt and fresh water; nests on stumps or in low trees.

ME.—Common migrant, rare in winter. April–June 20; July  
25–Nov. 20 (winter).

N. H.—Common migrant and rare winter resident coastwise.  
May; Aug. 3–Oct. 20 (winter).

VT.—Rare migrant; ? "summer resident."

MASS.—Common migrant, mainly coastwise, and rare winter  
resident. Mar. 10–June 3; July 24–Nov. 25 (winter).

## LONGIPENNES.

## LARIDAE.

R. I.—Common migrant and uncommon winter resident. Aug. 5–May 22.

CONN.—Common migrant, mainly in fall. Aug. 1–Nov.; (winter ?)—April 21.

## 29. NEMA SABINI (Sabine) Leach.

**Sabine's gull; Fork-tailed gull.**

*Larus sabini* J. Sab., Trans. Linn. soc. London, 1818, vol. 12, p. 520, pl. 29. "WEST COAST OF GREENLAND. . . . LATITUDE 75° 29' N., AND LONGITUDE 60° 9' W." Audubon, Birds of Amer., 1844, vol. 7, p. 127, pl. 441. Egg, Seebohm, 1896, p. 105, pl. 36, fig. 8.

Arctic seas and coasts; nests on the ground.

ME.—Accidental visitor: Brothers Id., near Portland, Sept. 22, 1899; Scarborough, May 31, 1877; (*nee* Calais, 1878).

MASS.—Accidental visitor: Boston Harbor, Sept. 27, 1874; Cape Cod, 1888; North Truro, Aug. 21, 1889.

R. I.—Doubtfully recorded.

## 30. GELOCHELIDON NILOTICA (Hasselquist) Stejneger.

**Gull-billed tern; Marsh tern.**

*Sterna nilotica* Hasselq., Reise nach Pal., deutsch. ausg., 1762, p. 325. [NILE RIVER, EGYPT].

*Sterna anglica* Montagu. Audubon, Birds of Amer., 1844, vol. 7, p. 81, pl. 430. Egg, Seebohm, 1896, p. 99, pl. 31, figs. 6, 8.

Seacoasts; nests on the ground.

ME.—Accidental visitor, several instances. March; May 21; Sept.

MASS.—Accidental visitor: Ipswich, Sept. 1871.

R. I.—Doubtfully recorded.

## LONGIPENNES.

## LARIDAE.

## 31. STERNA CASPIA Pallas.

**Caspian tern; Squawker.**

Pallas, Nov. comm. petrop., 1770, vol. 14, pt. 1, p. 582, pl. 22, fig. 2. "MARE CASPIUM ET CIRCA OSTIUM TAICL." Dresser, Birds of Europe, 1871-81, vol. 8, pl. [584]. Egg, Seebohm, 1896, p. 100, pl. 31, fig. 2.

*Sterna tschegrava* Lepech. Chapman, Handb. birds east. No. Amer., 1895, p. 79, fig. 61, *a*.

Seacoasts and lakes; nests on the ground.

ME.—Rare migrant. May 11; Aug.—Sept. (Oct. 16).

N. H.—Rare migrant: Hampton, 1899.

MASS.—Rare spring and uncommon fall migrant. Early May; Aug. 8—Sept. 26.

R. I.—Rare migrant. May 10; July 27—Oct.

## 32. STERNA MAXIMA Boddaert.

**Royal tern; Cayenne tern.**

Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 58. Based on Buffon: "CAYENNE." Chapman, Handb. birds east. No. Amer., 1895, p. 79, fig. 61, *b*.

*Sterna cayana* Lath. Audubon, Birds of Amer., 1844, vol. 7, p. 76, pl. 429.

Seacoasts and lagoons; nests on the ground.

MASS.—Accidental visitor: Chatham, July 29, 1889; Ipswich, July 17, 1904; Nantucket, July 1, 1874.

## 33. STERNA SANDVICENSIS ACUFLAVIDA (Cabot) Ridgway.

**Cabot's tern.**

*Sterna acuflavida* Cabot, Proc. Boston soc. nat. hist., 1847, vol. 2, p. 257. "TANCAH, ON THE COAST OF YUCATAN."

## LONGIPENNES.

## LARIDAE.

*Sterna cantiaea* Gmel. Audubon, Birds of Amer., 1844, vol. 7, p. 87, pl. 431.

Seacoasts; nests on the ground.

MASS.—Accidental visitor: Chatham, Aug., 1865; Monomoy Id., Oct. 2, 1888.

## 34. STERNA FORSTERI Nuttall.

**Forster's tern.**

Nutt., Manual ornith. U. S. and Canada, 1834, vol. 2, p. 274. Based on Forster: "CHURCHILL RIVER," HUDSON BAY. Saunders, Cat. birds British mus., 1896, vol. 25, p. 47.

*Sterna havellii* Audubon, Birds of Amer., 1844, vol. 7, p. 103, pl. 434.

Marshes and sea coasts; nests in grassy marshes.

MASS.—Rare fall migrant. Aug.—Oct. 2.

## 35. STERNA HIRUNDO Linné.

**Common tern; Mackerel gull; Medrake or Medrick; Red-shank; Sea swallow; Summer gull; Wilson's tern.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 137. "EUROPA." Chapman, Handb. birds east. No. Amer., 1895, p. 81, fig. 61, c; Audubon, Birds of Amer., 1844, vol. 7, p. 97, pl. 433. Egg, Capen, 1886, pl. 23, figs. 4, 5.

Seacoasts; nests on sandy or pebbly shores.

ME.—Abundant summer resident coastwise. May 3—Sept. 2 (late Sept.). Eggs, late May—Aug. 19.

N. H.—Common migrant coastwise, casual inland; formerly summer resident at Isles of Shoals. May; Aug.—Sept.

VT.—Rare visitor: Bennington, May 30, 1907; St. Albans.

## LONGIPENNES.

## LARIDAE.

MASS.—Locally common summer resident coastwise, casual inland; accidental in winter (Woods Hole, Jan. 17, 1891, Feb. 20, 1891). May 1–Oct. 30 (Nov. 14, winter). Eggs, May 20–July 30.

R. I.—Common summer resident coastwise. May 10–Oct. 6. Eggs, June 5–July 11.

CONN.—Common local summer resident coastwise. Eggs, June 20.

## 36. STERNA PARADISAEA Brünnich.

**Arctic tern.**

Brünn., Ornith. borealis, 1764, p. 46. "CHRISTIANSÖE," DENMARK. Chapman, Handb. birds east. No. Amer., 1895, p. 82, fig. 61, *d*.

*Sterna arctica* Temm. Audubon, Birds of Amer., 1844, vol. 7, p. 107, pl. 436. Egg, Capen, 1886, pl. 23, fig. 6–8.

Seacoasts; nests on the ground.

ME.—Common summer resident coastwise. May–Sept. Eggs, June 28.

MASS.—Uncommon local summer resident coastwise; occasional inland. May–Oct.

CONN.—Migrant.

## 37. STERNA DOUGALLI Montagu.

**Roseate tern; Mackerel gull.**

Montagu, Ornith. dict., suppl., 1813, p. [386], 1 pl. "WEST HIGHLANDS OF SCOTLAND." Chapman, Handb. birds east. No. Amer., 1895, p. 83, fig. 61, *e*; Dresser, Birds of Europe, 1871–81, vol. 8, pl. [581]. Egg, Capen, 1886, pl. 23, fig. 9–11.

Seacoasts; nests on sandy or pebbly shores.

## LONGIPEXNES.

## LARIDAE.

ME.—Rare summer visitor coastwise; not certainly known to breed. May–Aug. 31.

N. H.—Formerly rare summer resident at Isles of Shoals.

MASS.—Locally common summer resident coastwise. May 1–Oct. 4. Eggs, May 20–July 30.

R. I.—Common migrant and summer visitor; not known to nest. May–Oct.

CONN.—Common migrant and local summer resident. Eggs, June 20.

## 38. STERNA ANTILLARUM (Lesson) Coues.

**Least tern; Little striker; Oyt.**

*Sternula antillarum* Less., Descr. mamm. et. ois., 1847, p. 256.  
“ANTILLES . . . PRINCIPALEMENT GUADELOUPE.”

*Sterna minuta* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 119, pl. 439. Egg, Capen, 1886, pl. 23, fig. 12–14.

Seacoasts and lakes; nests on the ground.

ME.—Formerly rare summer resident in Casco Bay.

N. H.—Rare migrant: Hampton.

MASS.—Uncommon summer resident, from Cape Cod south (formerly at Ipswich); elsewhere now casual. (April 27) May 4–Oct. Eggs, May 31–June 29.

R. I.—Now rare but formerly a more common summer visitor coastwise. –Aug. 20.

CONN.—Rare migrant and summer visitor, formerly common.

## 39. STERNA FUSCATA Linné.

**Sooty tern; Egg-bird.**

Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 228. “IN INSULA DOMINICENSI.”

## LONGIPENNES.

## LARIDAE.

*Sterna fuliginosa* Lath. Audubon, Birds of Amer., 1844, vol. 7, p. 90, pl. 432. Egg, Seebohm, 1896, p. 104, pl. 30, fig. 4-6.

Seacoasts: nests on sandy shores.

ME.—Accidental visitor: Parkman, Oct. 5, 1878.

N. H.—Accidental visitor: Newmarket, about Sept. 14, 1878; (*nee* Lake Winnepesaukee).

VT.—Accidental visitor: Lake Champlain, early Sept. 1876; Rutland, 1876.

MASS.—Accidental visitor: Chatham, Sept. 1877; Lawrence, Oct. 29, 1876; Williamstown, Sept. 1876.

R. I.—Accidental visitor: (?) Newport, 1877; Point Judith, Sept. 1876; Woonsocket, July 16, 1897.

CONN.—Accidental visitor. Sept.

## 40. HYDROCHELIDON NIGRA SURINAMENSIS (Gmelin) Stejneger.

**Black tern.**

*Sterna surinamensis* Gmel., Syst. nat., 1788, vol. 1, pl. 2, p. 604. "SURINAM."

*Sterna nigra* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 116, pl. 438.

Fresh and salt water; nests in fresh marshes.

ME.—Rare fall migrant. Aug. 14-28.

N. H.—Uncommon migrant, mainly coastwise. June 10; Aug. 24-Sept. 30.

VT.—Rare migrant: Lake Bomoseen, June 6, 1883.

MASS.—Rare spring and uncommon early fall migrant. June 7-10; (July 11) Aug. 8-Sept. 26.

R. I.—Uncommon migrant, mainly in fall, coastwise. Spring; July 30-Sept. 18.

CONN.—Uncommon migrant coastwise, mainly in fall. Aug. 24.

## LONGIPENNES.

## RYNCHOPIDAE.

## 41. RYNCHOPS NIGRA Linné.

**Black skimmer; Cut-water; Scissor-bill; Sea-crow; Sea-dog;  
Razor-bill.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 138. "AMERICA.  
Audubon, Birds of Amer., 1844, vol. 7, p. 67, pl. 428.

Seacoasts; nests on sandy shores.

ME.—Accidental visitor: Matinicus Id.; off Saco, Aug. 31,  
1879; Washington Co.; Wells Bay.

MASS.—Occasional summer visitor coastwise, said to have  
bred about 1830 at Muskeget Id.; casual inland (Springfield).  
July 16–Aug. 19.

CONN.—Accidental visitor: Stonington (Linsley).

## TUBINARES.

## PROCELLARIIDAE.

## 42. FULMARS GLACIALIS (Linné) Stephens.

**Fulmar; Marbleheader; Noddy; Oil-bird.**

*Procellaria glacialis* Linné, Fauna Suec., ed. 2, 1761, p. 51.  
"IN MARI SEPTENTR. INTRA CIRCULUM ARCTICUM." Audubon,  
Birds of Amer., 1844, vol. 7, p. 204, pl. 455. Egg, Seebohm, 1896,  
p. 73, pl. 20, fig. 5.

Mainly pelagic; nests in shallow burrows or on cliffs.

ME.—Rare winter visitor offshore: "coast of Maine," March,  
1879.

MASS.—Rare fall and winter visitor offshore: Georges Bank.  
Oct. 28–winter.



## TUBINARES.

## PROCELLARIIDAE.

## 43. PUFFINUS BOREALIS Cory.

**Cory's shearwater.**

Cory, Bull. Nuttall ornith. club, 1881, vol. 6, p. 84. "NEAR CHATHAM ISLAND, CAPE COD, MASS."

Pelagic; nest unknown.

MASS.—Uncommon summer and fall visitor offshore. Aug. 2–Nov.

R. I.—Uncommon summer and fall visitor offshore. Aug. 15–Oct. 26.

## 44. PUFFINUS GRAVIS (O'Reilly) Salvin.

**Greater shearwater; Hag; Hagdon.**

*Procellaria gravis* O'Reilly, Greenland, adjacent seas, and northwest pass., 1818, p. 140, pl. 12, fig. 1. "LATITUDE OF CAPE FAREWELL AND STATEN HOOK," GREENLAND.

*Puffinus cinereus* Lath. Audubon, Birds of Amer., 1814, vol. 7, p. 212, pl. 456.

Pelagic; nest and eggs unknown.

ME.—Common summer visitor offshore. Aug. 10.

N. H.—Summer visitor offshore: Portsmouth.

MASS.—Common summer and fall visitor offshore; casual inland (Dover). May 26–Nov. (? Dec. 31).

R. I.—Uncommon summer and fall visitor offshore.

CONN.—Uncommon summer and fall visitor offshore.

## 45. PUFFINUS GRISEUS (Gmelin) Finsch.

**Sooty shearwater; Black hag or hagdon.**

Godman, Monogr. pterids, 1908, pt. 2, p. 145, pl. 38.

*Procellaria grisea* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 561. "IN HEMISPHERIO AUSTRALI, INTER 35° & 50°."

Pelagic; nests in burrows on sea islands.

## TUBINARES.

## PROCELLARIIDAE.

ME.—Common summer visitor offshore. May–summer.

MASS.—Common summer and fall visitor offshore. (Mar.)  
July 4–Oct.

R. I.—Uncommon summer visitor offshore. (Mar. 28) May–  
Sept.

46. *AESTRELATA HASITATA* (Kuhl) Coues.

**Black-capped petrel; Diablotin.**

*Procellaria hasitata* Kuhl, Beitr. z. zool., 1820, pt. 2, p. 142.  
No locality.

*Oestrelata haesitata* (Kuhl). Godman, Monogr. petrels, 1908,  
pt. 3, p. 184, pl. 49.

Mainly pelagic; nests in burrows on islands.

N. H.—Accidental visitor: Pittsfield, about Aug. 30, 1893.

Vt.—Accidental visitor: last of August, 1893 (place unknown).

47. *DAPTION CAPENSIS* (Linné) Stephens.

**Pintado petrel; Cape pigeon.**

Salvin, Cat. birds British mus., 1896, vol. 25, p. 428. Gould,  
Birds of Australia, 1848, vol. 7, pl. 53.

*Procellaria capensis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 132.  
“AD CAP[ITEM] B[ONAE] SPEI.”

Pelagic; nests in burrows on islands of southern hemisphere.

ME.—Accidental visitor: Lake Mooseluckmeguntic, Sept. 1872  
(*see* Harpswell).

48. *OCEANODROMA LEUCORHOA* (Vieillot) Stejneger.

**Leach's petrel; Carey chicken; Mother Carey's chicken.**

*Procellaria leucorhoa* Vieill., Nouv. dict. d'hist. nat., 1817, vol.  
25, p. 422. “LES BORDS MARITIMES DE LA PICARDIE,” FRANCE.

## TUBINARES.

## PROCELLARIIDAE.

*Thalassidroma leachii* Temm. Audubon, Birds of Amer., 1844, vol. 7, p. 219, pl. 459. Egg, Capen, 1886, pl. 24, fig. 1.

Pelagic; nests in colonies on islands, in burrows.

ME.—Abundant summer resident on outer islands from Casco Bay north. May–Oct. 21. Eggs, June 10–Aug. 19.

N. H.—Common migrant offshore; occasional inland in fall. Spring–Nov.

VT.—Accidental visitor: Lake Champlain.

MASS.—Common migrant offshore; casual inland in fall. June 21; Sept. 1–Nov. 25.

R. I.—Uncommon migrant offshore. June 9; Aug. 29–Oct. 14.

CONN.—Uncommon migrant offshore, casual inland. Oct. 27.

## 49. OCEANITES OCEANICUS (Kuhl) Lichtenstein.

**Wilson's petrel; Mother Carey's chicken.**

*Procellaria oceanica* Kuhl, Beitr. z. zool., 1820, pt. 2, p. 136, pl. 10, fig. 1. No locality given.

*Thalassidroma wilsonii* Bonap. Audubon, Birds of Amer., 1844, vol. 7, p. 223, pl. 460.

Pelagic; nests within Antarctic circle, in burrows.

ME.—Common summer visitor offshore; casual inland (near Augusta). June–Sept. 17.

N. H.—Common summer visitor offshore.

VT.—Accidental visitor: St. Albans (may have been *O. leucorhoa*).

MASS.—Common summer visitor offshore. June 17–Sept. 23.

R. I.—Common summer visitor offshore. July–Aug.

CONN.—Uncommon summer visitor offshore.

## STEGANOPODES.

## SULIDAE.

50. *SULA LEUCOGASTRA* (Boddaert) Selater & Salvin.

**Booby; Booby gannet.**

*Pelecanus leucogaster* Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 57. Based on Buffon.

*Sula fusca* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 57, pl. 426.

Seacoasts; nests on the ground.

MASS.—Accidental visitor: Cape Cod, about Sept. 17, 1878 (*nee* Essex Co.).

CONN.—Linsley's record (Guilford) said to refer to *S. bassana*.

51. *SULA BASSANA* (Linné) Boie.

**Gannet; Solan goose; White gannet.**

Audubon, Birds of Amer., 1844, vol. 7, p. 44, pl. 425. Egg, Seebohm, 1896, p. 69, pl. 19, fig. 4.

*Pelecanus bassanus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 133. "IN SCOTIA, AMERICA."

Coastwise; nests in colonies on a few rocky islets of North Atlantic.

ME., N. H.—Uncommon migrant. April–May; Sept.–Oct. 9.

MASS.—Common migrant offshore. Mar. 26–June 7; Aug. 3–Dec. 21.

R. I.—Uncommon migrant offshore. April–May 16; Sept. 10–Oct. 26.

CONN.—Rare migrant offshore.

## STEGANOPODES.

## PHALACROCORACIDÆ.

## 52. PHALACROCORAX CARBO (Linné) Leach.

**Cormorant; Common cormorant; Shag; Taunton shag; Taunton turkey.**

Audubon, Birds of Amer., 1843, vol. 6, p. 412, pl. 415. Coues, Key No. Amer. birds, ed. 5, 1903, vol. 2, p. 963. Egg, Capen, 1886, pl. 21, fig. 8.

*Pelecanus carbo* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 133. "IN EUROPA."

Salt water mainly; nests in colonies on sea cliffs.

ME.—Common migrant and uncommon winter resident. Oct.—April 20.

VT.—Rare migrant: Rutland, 1867.

MASS.—Uncommon migrant and rare winter resident coastwise. April–May 14; Sept. 22–Nov. (winter).

R. I.—Uncommon migrant and winter resident. Sept. 15–(June 3).

CONN.—Uncommon migrant and winter resident. Fall–May.

## 53. PHALACROCORAX AURITUS (Lesson) Grant.

**Double-crested cormorant; Shag; Taunton turkey.**

*Carbo auritus* Less., Traité d'ornith., 1831, vol. 1, p. 605. "NOUVELLE-ZÉLANDE." [!]

*Phalacrocorax dilophus* Coues, Key No. Amer. birds, ed. 5, 1903, vol. 2, p. 963, fig. 676; Audubon, Ornith. biogr., 1835, vol. 3, pl. 257. Egg, Capen, 1886, pl. 22, fig. 1.

Coastal; nests in colonies on sea cliffs.

ME.—Common migrant; rare winter and summer (Black Horse Ledge) resident. Eggs, June 23–July 3.

N. H.—Uncommon migrant coastwise; occasional inland. Spring; Oct.

## STEGANOPODES.

## PHALACROCORACIDAE.

Vt.—Rare migrant: Burlington, Nov.; Cambridge, Dec.; Windsor.

Mass.—Common migrant, mainly coastwise. April 4–June 18 (July 7); Aug. 22–Nov. 24.

R. I.—Common migrant, mainly coastwise. April 22–May 22; (Aug. 21) Sept. 16–Nov.

Conn.—Uncommon migrant, mainly coastwise. April–May; mid-July–Nov. 13.

## PELECANIDAE.

## 54. PELECANUS ERYTHORHYNCHOS Gmelin.

**American white pelican.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 571. "IN AMERICA SEPTENTRIONALI."

*Pelecanus americanus* Audubon, Birds of Amer., 1844, vol. 7, p. 20, pl. 422.

Salt and fresh water; nests on the ground.

Me.—Accidental visitor: Calais; Eliot, pair, June 8, 1897; Saponic Lake, May 28, 1892.

N. H.—Formerly a visitor (Belknap).

Mass.—Accidental visitor: Gloucester, 1886; North Scituate, Oct. 5, 1876; Sandwich, May 13, 1905.

## 55. PELECANUS OCCIDENTALIS (Linné) Richmond.

**Brown pelican.**

*Pelecanus onocrotalus occidentalis* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 215. "AMERICA."

*Pelecanus fuscus* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 32, pls. 423, 424.

Coastal; nests on the ground or in low trees.

## STEGANOPODES.

## PELECANIDAE.

ME.— The supposed Bar Harbor specimen was an escape brought from South America.

MASS.— Accidental visitor: (? Ipswich, doubtful); Nantucket, about 1867 (flock of 13).

CONN.— Accidental visitor: near New Haven.

## FREGATIDAE.

## 56. FREGATA AQUILA (Linné) Reichenbach.

**Man-o'-war bird; Frigate-bird; Man-o'-war hawk.**

*Pelecanus aquilus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 133.  
"IN INSULA ADSCENSIONIS ALISQUE PELAGICIS."

*Tachypetes aquilus* Linné. Audubon, Birds of Amer., 1844, vol. 7, p. 40, pl. 421.

Coastal; nests in colonies, in low trees or on ground.

ME.— Doubtfully recorded from Boothbay.

MASS.— Accidental visitor: New Bedford, Oct. 17, 1893 (record not confirmed).

CONN.— Accidental visitor: Faulkner's Id., autumn, 1859.

## ANSERES.

## ANATIDAE.

## 57. MERGUS AMERICANUS Cassin.

**American Merganser; Buff-breasted Merganser; Goosander;  
Pond shelldrake.**

Cassin, Proc. acad. nat. sci. Phila., 1853, vol. 6, p. 187. Based on Wilson, no locality; probably New Jersey or Pennsylvania.

*Mergus merganser* Linné. Audubon, Birds of Amer., 1843, vol. 6, p. 387, pl. 411. Egg, Capen, 1886, pl. 21, fig. 5.

Mainly in fresh water; nests in hollow trees.

## ANSERES.

## ANATIDAE.

ME.—Common migrant and winter resident; less common summer resident inland. Eggs, late May–June 19.

N. H.—Common migrant; less common summer resident from White Mts. north, and winter resident in open streams south of White Mts.

VT.—Uncommon summer resident; rare winter resident in open streams. Eggs, June 18.

MASS.—Uncommon migrant and winter resident, mainly on freshwater lakes and streams; said to have bred formerly. Oct. 12–May 1; summer (Springfield).

R. I.—Uncommon migrant and winter resident, mainly on ponds and rivers. Nov.–May 2.

CONN.—Uncommon migrant and winter resident, mainly on ponds and rivers. Fall–April 17.

58. *Mergus serrator* Linné.**Red-breasted Merganser; Shelldrake.**

Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 129. "IN EUROPA." Audubon, *Birds of Amer.*, 1843, vol. 6, p. 395, pl. 412. Egg Capen, 1886, pl. 21, fig. 6.

Salt and fresh water; nests on the ground.

ME.—Common migrant and winter resident coastwise; uncommon summer resident on coastal islands or inland lakes. Eggs, June 21.

N. H.—Common migrant and winter resident mainly coastwise. Nov.–April.

VT.—Uncommon migrant. April; Oct.

MASS.—Common migrant and winter resident, mainly coastwise; a few non-breeding birds summer offshore. Sept. 23–May 26 (summer).



## ANSERES.

## ANATIDAE.

R. I.—Common migrant and winter resident coastwise; a few non-breeding birds summer offshore. Oct. 29–May 19 (summer).

CONN.—Common migrant and winter resident, mainly coastwise. Fall–April 14.

59. *LOPHODYTES CUCULLATUS* (Linné) Reichenbach.

**Hooded Merganser; Hairy-crown; Hairy-head; Hooded shell-drake; Sawbill diver; Water pheasant.**

*Mergus cucullatus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 129. "AMERICA." Audubon, Birds of Amer., 1843, vol. 6, p. 402, pl. 413. Egg, Capen, 1886, pl. 21, fig. 7.

Fresh water mainly; nests in hollow trees.

ME.—Rare migrant and summer resident. Mar. 31–May; summer; Sept.–late Nov.

N. H.—Rare migrant and summer resident (Lake Umbagog). Mar.–Nov.

VT.—Rare summer resident. Mar. 27–Sept.

MASS.—Uncommon migrant. Mar. 18–April 18; (Aug. 11, Cambridge) Oct. 7–Nov. 30 (Dec. 22).

R. I.—Uncommon migrant and rare winter resident. Oct.–May.

CONN.—Uncommon migrant and rare winter resident. Nov.–Mar. 6.

60. *ANAS PLATYRHYNCHOS* Linné.

**Mallard; Greenhead; Wild mallard.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 125. "IN EUROPAE MARITIMIS."

*Anas boschas* Linné. Audubon, Birds of Amer., 1843, vol. 6, p. 236, pl. 385.

Rivers, ponds, and in migrations, seacoasts; nests in marshes.

## ANSERES.

## ANATIDAE.

ME.—Rare migrant and winter resident. April; Oct. 1–Nov. 15 (winter).

N. H.—Rare migrant. Spring; Oct.–Nov. 9.

VT.—Rare migrant.

MASS.—Uncommon migrant; rare in winter. Mar. 27–May 1; Sept. 22–Dec. 17 (Dec. 26, Jan. 9).

R. I.—Rare migrant and winter resident. Oct.–April.

CONN.—Rare spring and uncommon fall migrant. Spring; Sept. 30–Nov.

61. *ANAS RUBRIPES* (Brewster) A. O. U. comm.

**Black duck; Black mallard; Dusky duck; Red-legged black duck; Spring black duck; Winter black duck.**

*Anas obscura rubripes* Brewst., Auk, 1902, vol. 19, p. 184. "LAKE UMBAGOG (NEW HAMPSHIRE SHORE)."

*Anas obscura* Gmel. Audubon, Birds of Amer., 1843, vol. 6, p. 244, pl. 386. Egg, Capen, 1886, pl. 20, fig. 10.

Salt and fresh water; nests in marshes.

ME., N. H.—Common migrant, less common summer and (coastwise) winter resident. Eggs, April 28 (Me.).

VT.—Common migrant, less common summer resident. Eggs, May.

MASS.—Common migrant and winter resident (mainly coastwise); uncommon local summer resident. Eggs, April 19–May.

R. I.—Common migrant and winter resident, rare summer resident. Sept. 29–May; summer. Eggs, May 5–21.

CONN.—Common migrant and winter resident; rare summer resident. Sept.–May; summer.

62. *CHAULELASMUS STREPERUS* (Linné) Bonaparte.

**Gadwall; Gray duck.**

*Anas strepera* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 125. "IN

## ANSERES.

## ANATIDAE.

EUROPAE AQUIS DULCIBUS." Audubon, Birds of Amer., 1843, vol. 6, p. 254, pl. 388. Egg, Seebohm, 1896, p. 37, pl. 14, fig. 3.

Fresh and salt water; nests in marshes.

ME.—Rare migrant. April 29, 1879 (Scarborough); Oct. 27–29.

N. H.—Rare migrant: Hampton; Portsmouth, spring.

VT.—Rare migrant: Otter Creek, spring 1879; St. Albans.

MASS.—Rare migrant. Oct. 2–Nov. 1.

R. I.—Rare migrant. Feb. 26–May 4; Oct. 8–Dec. 1.

CONN.—Rare migrant. Aug.–Oct. 14.

## 63. MARECA PENELOPE (Linné) Selby.

**Widgeon; European widgeon.**

*Anas penelope* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 126. "IN EUROPAE MARITIMIS & PALUDIBUS." Salvadori, Cat. birds of British mus., 1895, vol. 27, p. 227. Dresser, Birds of Europe, 1871–81, vol. 6, pls. [432, 433]. Egg, Seebohm, 1896, p. 38, pl. 13, fig. 4.

Fresh and salt water; nests on the ground.

VT.—Doubtfully recorded.

MASS.—Accidental visitor: Halifax, Oct. 20, 1899; Marblehead, Dec. 29, 1900; Mass. ? (Samuels).

R. I.—Accidental visitor: Middletown, Sept. 20, 1889.

## 64. MARECA AMERICANA (Gmelin) Stephens.

**Baldpate; American widgeon.**

*Anas americana* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 526. "AMERICA A CAYENNA INSULISQUE VICINI OCEANI AD SINUM HUDSONIS." Audubon, Birds of Amer., 1843, vol. 6, p. 259, pl. 389.

Fresh and salt water; nests in marshes.

## ANSERES.

## ANATIDAE.

ME.— Uncommon migrant, mainly coastwise; accidental in winter. Mar. 18–April; Sept.–Nov. 18 (Feb. 20, 1884, Freeport).

N. H.— Uncommon migrant. April; Sept.–Oct.

VT.— Rare migrant.

MASS.— Uncommon migrant and rare winter resident (Nantucket). Mar. 22–April 17; Sept. 6–Dec. 2 (winter).

R. I.— Uncommon migrant. ? April; Sept. 20–early Dec.

CONN.— Uncommon migrant.

65. *NETTION CRECCA* (Linné) Kaup.**European teal; Teal.**

Salvadori, Cat. birds British mus., 1895, vol. 27, p. 243.

*Anas crecca* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 126. "IN EUROPAE AQUIS DULCIBUS."

*Querquedula crecca* (Linné). Dresser, Birds of Europe, 1871–81, vol. 6, pl. [426]. Egg, Seebohm, 1896, p. 40, pl. 16, fig. 3.

Fresh water mainly; nests in marshes.

ME.— Accidental visitor: Casco Bay, April 6, 1903.

MASS.— Accidental visitor: "Massachusetts" (Samuels); Muskeget Id., Mar. 16, 1890; Sagamore, Feb. 20, 1896.

CONN.— Accidental visitor: East Hartford, Nov. 14, 1889.

66. *NETTION CAROLINENSIS* (Gmelin) Baird.**Green-winged teal.**

*Anas carolinensis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 533. "IN AMERICA, A CAROLINA AD SINUM HUDSONIS, STAGNIS SILVATICIS." Audubon, Birds of Amer., 1843, vol. 6, p. 281, pl. 392. Egg, Capen, 1886, pl. 20, fig. 11.

Fresh and salt water; nests in marshes.

ME.— Common migrant. April; Sept. 15–Nov. 15.

N. H.— Uncommon migrant.

## ANSERES.

## ANATIDAE.

VT.—Common migrant; (? “summer resident”). Mar.; Sept.

MASS.—Uncommon spring and rare fall migrant; accidental in winter. April 4–23; Sept. 6–Dec. 13 (Jan. 1; Feb. 1, 1892, West Barnstable).

R. I.—Uncommon migrant; rare in winter. Mar.; Sept. 2–Nov. 7 (Dec. 30).

CONN.—Uncommon migrant and rare winter resident. Fall–Mar.

67. *QUERQUEDULA DISCORS* (Linné) Stephens.**Blue-winged teal.**

*Anas discors* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 205. “IN AMERICA SEPTENTRIONALI.” Audubon, Birds of Amer., 1843, vol. 6, p. 287, pl. 393. Egg, Capen, 1886, pl. 20, fig. 12.

Fresh and salt water; nests in marshes.

ME.—Common migrant and in Washington Co., a rare summer resident. (Mar. 8) April–May; summer; Sept. 15–Oct. 16.

N. H.—Uncommon migrant. May; Aug. 15–Nov.

VT.—Uncommon migrant; (? “summer resident”). April–May; fall.

MASS.—Rare spring and common fall migrant; probably once bred. April 11–May 31; (June 8 and 20, Cambridge); Aug. 16–Nov. 26.

R. I.—Rare spring and uncommon fall migrant and rare summer resident (at Sakonnet). May (summer); Aug. 27–Nov. 5. Eggs, May.

CONN.—Rare spring and uncommon fall migrant. Spring; Aug.–Oct. 20.

68. *SPATULA CLYPEATA* (Linné) Boie.**Shoveler; Spoonbill.**

*Anas clypeata* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 124. “IN

## ANSERES.

## ANATIDAE.

EUROPAE MARITIMIS." Audubon, Birds of Amer., 1843, vol. 6, p. 293, pl. 394. Egg, Seebohm, 1896, p. 42, pl. 13, fig. 2.

Fresh and salt water; nests in marshes.

ME.—Rare migrant coastwise. April 14–23; Sept.

N. H.—Rare migrant: Hampton; Rye Beach, Aug. 1872.

VT.—Rare migrant.

MASS.—Rare migrant. Spring; Sept. 17–Nov. 7.

R. I.—Rare migrant. April 10; Oct. 2–28.

CONN.—Rare migrant. (Late July) Aug.–Dec. 8.

69. *DAFILA ACUTA* (Linné) Bonaparte.

**Pintail; Gray duck (♀); Pheasant duck; Sprigtail.**

*Anas acuta* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 126. "IN EUROPAE MARITIMIS." Audubon, Birds of Amer., 1843, p. 266, pl. 390. Egg, Seebohm, 1896, p. 38, pl. 13, fig. 3.

Fresh and salt water; nests in marshes.

ME.—Rare migrant. Late Mar.–April; Sept. 20–Nov. 7 (late Nov.).

N. H., VT.—Rare migrant.

MASS.—Rare migrant. (Feb. 21) Mar. 8–April 3; Sept. 11–Dec. 21.

R. I.—Uncommon migrant coastwise. Spring; Sept. 4.

CONN.—Uncommon migrant; (? rare winter resident). Mar. 23; fall.

70. *AIX SPONSA* (Linné) Bonaparte.

**Wood duck; Bride duck; Rainbow duck; Regal duck; Summer duck.**

*Anas sponsa* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 128. "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer., 1843, vol. 6, p. 271, pl. 391. Egg, Capen, 1886, pl. 21, fig. 1.

## ANSERES.

## ANATIDAE.

Freshwater streams and ponds; nests in hollow trees, rarely on the ground.

ME.—Uncommon migrant and summer resident. Mar. 28—late Nov.

N. H.—Uncommon migrant and summer resident. Mar.—Dec.

VT.—Uncommon migrant and summer resident. April 8—Oct. 23.

MASS.—Uncommon migrant and rare summer resident; accidental in winter. (Mar. 10) Mar. 20—April 30; summer; Sept. 15—Nov. 18 (Dec. 31, 1890, So. Weymouth; Jan. 11, 1888, Taunton). Eggs, April 23—May 2.

R. I.—Uncommon migrant and rare summer resident. Mar.—Nov. 11. Eggs, May 10.

CONN.—Uncommon migrant and rare summer resident; accidental in winter. Mar. 18—Oct. (Jan. 9, 1880, Norwich). Eggs, May.

71. *MARILA AMERICANA* (Eytou) A. O. U. comm.**Redhead; American pochard.**

*Fuligula americana* Eyt., Monogr. Anatidae, 1838, p. 155. "NORTH AMERICA."

*Fuligula ferina* Linné. Audubon, Birds of Amer., 1843, vol. 6, p. 311, pl. 396.

Fresh and salt water; nests in marshes.

ME.—Rare migrant; said formerly to have bred in Washington Co. Mar. 27; Sept. 21—Nov. 5.

N. H.—Rare fall migrant.

VT.—Rare fall migrant: Lake Champlain.

MASS.—Rare migrant more frequent in fall, and rare winter resident on south coast. Mar. 6—April 3; Oct. 4—Dec. 21 (winter).

R. I.—Rare migrant and winter resident. Oct. 8—Feb. 3.

CONN.—Rare migrant.

## ANSERES

## ANATIDAE.

72. *MARILA VALLISNERIA* (Wilson) A. O. U. comm.**Canvas-back.**

*Anas valisineria* Wils., Amer. ornith., 1814, vol. 8, p. 103, pl. 70, fig. 5. "HUDSON AND DELAWARE. . . RIVERS. . . AND IN THE NEIGHBOURHOOD OF THE CHESAPEAKE BAY."

*Fuligula valisneriana* (Wils.). Audubon, Birds of Amer., 1843, vol. 6, p. 299, pl. 395.

Fresh and salt water; nests in marshes.

ME.—Very rare fall migrant.

VT.—Very rare migrant: Lake Champlain (Perkins).

MASS.—Rare migrant. Mar. 11–31; Oct. 19–Dec. 18 (Jan. 8, Cambridge).

R. I.—Rare migrant. April; Oct. 2–Nov. 16.

CONN.—Rare migrant. May 7.

73. *MARILA MARILA* (Linné) A. O. U. comm.**Greater scaup; Blackhead; Bluebill; Bluehead; Broadbill; Raft duck; Widgeon.**

*Anas marila* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 196. "IN EUROPAE ALPINIS IN SVECIA, NORVEGIA, AUSTRIA."

*Fuligula marila* (Linné). Audubon, Birds of Amer., 1844, vol. 7, p. 355, pl. 498. Egg, Seebohm, 1896, p. 46, pl. 14, fig. 2.

Fresh and salt water; nests in marshes.

ME.—Common coastwise migrant; rare in winter. Oct.–Nov.; (winter); Mar.–April.

N. H.—Uncommon migrant, mainly coastwise. April; Oct.–Dec. 13.

VT.—Uncommon migrant at Lake Champlain.

MASS.—Rare spring, and common fall migrant; winters locally on coast. Mar. 27–May 1; Sept. 18–Dec. 26; winter.



## ANSERES

## ANATIDAE.

R. I.—Common migrant and less common winter resident coastwise. Oct. 1–May 1 (July 19, 1900, cripple).

CONN.—Common migrant and uncommon winter resident coastwise. Oct.–May 17 (June).

74. *MARILA AFFINIS* (Eytou) Bonaparte.

**Lesser scaup; Blue-billed shoveler; Creek broadbill; Little blackhead; Little bluebill; Little raft duck; River broadbill.**

*Fuligula affinis* Eyt., Monogr. Anatidae, 1838, p. 157. "NORTH AMERICA."

*Fuligula marila* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 316, pl. 397.

*Aythya affinis* (Eyt.). Chapman, Handb. birds east. No. Amer., 1895, p. 112.

Fresh and salt water; nests in marshes.

ME.—Rare migrant. Mar.–April; Oct.–Nov.

N. H.—Rare migrant.

VT.—Rare migrant: Lake Champlain; Otter Creek, May 19, 1883.

MASS.—Rare spring and common fall migrant; rare winter resident on south coast. Mar. 2–April 14 (May 5); Oct. 8–Dec. 6 (winter).

R. I.—Uncommon migrant and winter resident. Oct. 15–April.

CONN.—Uncommon migrant and winter resident. Oct.–May 17 (June).

75. *MARILA COLLARIS* (Donovan) Bonaparte.

**Ring-necked duck; Ring-neck scaup.**

*Anas collaris* Donov., British birds, 1809, vol. 6, pl. 147 and text. "SUPPOSED TO HAVE BEEN . . . LINCOLNSHIRE." ENGLAND.

*Fuligula rafitorques* Bonap. Audubon, Birds of Amer., 1843, vol. 6, p. 320, pl. 398.

## ANSERES.

## ANATIDAE.

*Aythya collaris* (Donov.). Chapman, Handb. birds east. No. Amer., 1895, p. 112.

Fresh and salt water; nests in marshes.

ME.—Rare migrant; once bred in Washington Co. Mar. 31–May 1; (summer); Oct. 11–Nov. 2.

N. H.—Rare migrant: Concord, Nov.; Little Bay near Portsmouth.

MASS.—Rare migrant. Mar.–April; Oct. 4–Nov. 30.

R. I.—Rare migrant. Nov. 11.

CONN.—Rare migrant.

76. *CLANGULA CLANGULA AMERICANA* (Bonaparte) Faxon.

**American goldeneye; Garrot; Whistler; Whistle-wing.**

*Clangula americana* Bonap., Geogr. and comparative list birds Europe and No. Amer., 1838, p. 58. "AMERICA."

*Fuligula clangula* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 362, pl. 406. Egg, Capen, 1886, pl. 21, fig. 2.

Fresh and salt water; nests in hollow trees.

ME.—Common migrant and winter resident coastwise; less common summer resident in northern part. Eggs, May 27–June 8.

N. H.—Common migrant and winter resident coastwise or in open rivers; uncommon summer resident (Lake Umbagog).

VT.—Common migrant on Lake Champlain and Connecticut River.

MASS.—Common migrant and winter resident mainly coastwise. Oct. 8–May 1 (July 26, Boston).

R. I.—Common migrant and winter resident coastwise. Nov. 15–April 20.

CONN.—Common migrant and winter resident mainly coastwise. Late Oct.–April 16.

## ANSERES.

## ANATHIDÆ.

## 77. CLANGULA ISLANDICA (Gmelin) Bonaparte.

**Barrow's goldeneye.**

*Anas islandica* Gmelin, Syst. nat., 1788, vol. 1, pt. 2, p. 541.  
 "IN ISLANDIA."

*Clangula barrovii* Swainson and Richardson, Fauna Boreali-  
 Amer., 1831, vol. 2, p. 456, pl. 70.

Fresh and salt water; nests in hollow trees.

ME.—Rare winter visitor coastwise. Jan.—April.

N. H.—Rare winter visitor: Little Bay, Portsmouth.

VT.—Erroneously recorded.

MASS.—Rare winter visitor coastwise. Dec. 7—Feb. 22.

CONN.—Rare winter visitor coastwise: off Portland, Nov. 14,  
 1867.

## 78. CHARITONETTA ALBEOLA (Linné) Stejneger.

**Bufflehead; Bumble-bee duck; Butterball; Devil diver; Dipper;  
 Hell diver; Spirit duck.**

*Anas albeola* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 124. "AMER-  
 ICA."

*Fuligula albeola* (Linné). Audubon, Birds of Amer., 1843, vol.  
 6, p. 369, pl. 408.

Fresh and salt water; nests in hollow trees.

ME.—Common migrant and on south coast an uncommon winter  
 resident; said to nest in Washington Co. Late Oct.—April (May  
 11) (summer).

N. H.—Uncommon migrant and rare winter resident chiefly  
 coastwise. Mar.—April; Oct. Nov. (winter).

VT.—Uncommon migrant.

MASS.—Uncommon migrant and rare winter resident. Mar. 11—  
 April 8; Oct. 9—Dec. 23 (winter).

## ANSERES.

## ANATIDAE.

R. I.—Uncommon migrant and rare winter resident coastwise. April; Oct. 13–Dec. 7 (winter).

CONN.—Uncommon migrant and winter resident. Nov.—April 15.

79. *HARELDA HYEMALIS* (Linné) L. Brehm.

**Old squaw; Long-tailed duck; Old injun; Old wife; Scolde-  
nore; Scolder; South-southerly.**

*Anas hyemalis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 126. "IN EUROPA & AMERICA ARCTICA."

*Fuligula glacialis* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 379, pl. 410. Egg, Seebohm, 1896, p. 50, pl. 15, fig. 3.

Salt and fresh water; nests on the ground.

ME.—Common migrant and winter resident coastwise; a few non-breeding birds summer. Late Nov.—May 21 (summer).

N. H.—Common migrant and winter resident, chiefly coastwise.

VT.—Rare migrant to Lake Champlain and Connecticut Valley. Oct. 24.

MASS.—Common migrant and winter resident, mainly coastwise. Oct. 10–May 22 (June 17).

R. I.—Common migrant and winter resident. Nov. 7–May 4.

CONN.—Common migrant and winter resident coastwise. Oct.—May.

80. *HISTRIONICUS HISTRIONICUS* (Linné) Boucard.

**Harlequin duck; Lord-and-lady; Sea mouse.**

*Anas histrionica* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 127. "AMERICA."

*Fuligula histrionica* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 374, pl. 409. Egg, Seebohm, 1896, p. 49, pl. 15, fig. 1.

Fresh and salt water; nests on the ground or in hollow trees.

## ANSERES.

## ANATIDAE.

ME.—Uncommon winter resident about rocky islands. Oct. 19–Mar. (April).

N. H.—Rare winter visitor coastwise. Nov.

MASS.—Rare winter visitor coastwise; casual inland. Nov. 1–Jan.

R. I.—Rare winter visitor coastwise. Nov. 28–Jan. 29.

81. *CAMPTORIHYNCHUS LABRADORIUS* (Gmelin) Bonaparte.

**Labrador duck; Pied duck; Sand-shoal duck.**

*Anas labradoria* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 537.  
“IN AMERICA BOREALI.”

*Fuligula labradora* Lath. Audubon, Birds of Amer., 1843, vol. 6, p. 329, pl. 400.

Formerly on seacoast; nests on “tops of low fir trees.” Last specimen taken in 1875.

ME.—Formerly a winter visitor coastwise.

MASS.—Formerly a winter visitor coastwise; last taken in the State at Swampscott, early September, 1862.

CONN.—Formerly a winter resident (Stratford).

82. *SOMATERIA MOLLISSIMA BOREALIS* (C. L. Brehm) A. E. Brehm.

**Northern eider; Greenland eider.**

*Platypus borealis* C. L. Brehm, Lehrb. naturg. europ. vögel, 1824, vol. 2, p. 813. COASTS AND ISLANDS OF BAFFIN BAY AND DAVIS STRAIT (*vide* H. C. Oberholser).

*Somateria mollissima* (Linné). Baird, Brewer, and Ridgway, Water birds No. Amer., 1884, vol. 2, p. 72, figs. on p. 77.

Salt water chiefly; nests on the ground.

ME.—Rare winter visitor coastwise. Oct. 19–April 6.

MASS.—Rare winter visitor coastwise. Late Oct.–April 3.

R. I.—Unsatisfactorily recorded from Newport.

## ANSERES.

## ANATIDAE.

## 83. SOMATERIA DRESSERI Sharpe.

**American eider; Canvasback; Sea duck; Wamp.**

Sharpe, Ann. mag. nat. hist., 1871, ser. 4, vol. 8, p. 51. [NORTH AMERICA].

*Fuligula mollissima* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 349, pl. 405. Egg, Capen, 1886, pl. 21, fig. 3.

Salt water chiefly; nests on the ground.

ME.—Common winter, and from Isle au Haut north, rare summer resident. Nov.—April; summer. Eggs, June 15—July 4.

N. H.—Uncommon migrant and winter resident coastwise.

MASS.—Uncommon migrant and winter resident coastwise. Nov. 5—April 19.

R. I.—Uncommon migrant and winter resident coastwise.

CONN.—Rare winter resident coastwise. Fall—(May 29).

## 84. SOMATERIA SPECTABILIS (Linné) Leach.

**King eider.**

*Anas spectabilis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 123. "IN CANADA, SVECIA."

*Fuligula spectabilis* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 347, pl. 404. Egg, Seebohm, 1896, p. 54, pl. 9, fig. 1.

Salt water chiefly; nests on the ground.

ME.—Uncommon winter resident coastwise. Dec. 11—April 15 (May 29).

N. H.—Rare winter visitor: Hampton.

MASS.—Rare winter visitor coastwise. Oct. 21—April 5.

R. I.—Rare winter visitor coastwise. (Sept. 15) Nov. 27—Feb.

CONN.—Rare winter visitor coastwise. Nov. 4.

## ANSERES.

## ANATIDAE.

85. *OIDEZIA AMERICANA* Swainson.

**American scoter; Black coot; Butterbill; Butterbill coot; Yellow-billed coot.**

Swains., in Swains. and Rich., Fauna Boreali-Amer., 1831, vol. 2, p. 450. "HUDSON'S BAY, LAT. 57°."

*Fuligula americana* (Swains.). Audubon, Birds of Amer., 1843, vol. 6, p. 343, pl. 403.

Salt and fresh water; nests on the ground.

ME., N. II.—Common migrant and winter resident mainly coastwise; a few non-breeding birds summer offshore.

VT.—Uncommon migrant: St. Johnsbury; Windsor.

MASS.—Common migrant and winter resident; a few non-breeding birds summer offshore. Sept. 8–May 24 (summer).

R. I.—Common migrant and winter resident coastwise. Oct.—May 11.

CONN.—Uncommon migrant and winter resident. Oct. 4–May.

86. *OIDEZIA DEGLANDI* Bonaparte.

**White-winged scoter; May white-wing; Velvet duck; White-winged coot.**

Bonap., Rev. crit. de l'orn. Europ. de Dr. Degland, 1850, p. 108. "AMERICA."

*Fuligula fusca* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 332, pl. 401.

Salt and fresh water; nests on the ground.

ME., N. II.—Common migrant and winter resident mainly coastwise; a few non-breeding birds summer offshore.

VT.—Rare migrant: Middlebury.

MASS.—Common migrant and winter resident mainly coastwise; a few non-breeding birds summer offshore. Sept. 6–May 25 (summer).

## ANSERES.

## ANATIDAE.

R. I.—Common migrant and winter resident, mainly coastwise; a few non-breeding birds summer offshore. Sept. 15–May; summer.

CONN.—Abundant migrant and common winter resident, mainly coastwise; a few non-breeding birds summer offshore. Oct. 4–May 15; summer.

87. *OIDEZIA PERSPICILLATA* (Linné) Stephens.

**Surf scoter; Baldheaded coot; Box coot; Gray coot; Hollow-billed coot; Patch-bill coot; Patch-poll coot; Sea coot; Skunk-head.**

Chapman, Handb. birds east. No. Amer., 1895, p. 118.

*Anas perspicillata* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 125.  
“CANADA.”

*Fuligula perspicillata* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 337, pl. 402. Egg, Seebohm, 1896, p. 52, pl. 15, fig. 2.

Salt water mainly; nests on the ground.

ME., N. H.—Common migrant and winter resident mainly coastwise; a few non-breeding birds summer offshore.

VT.—Rare migrant: Bradford.

MASS.—Common migrant and winter resident coastwise; a few non-breeding birds summer offshore. Sept. 4–May 9; summer.

R. I.—Common migrant and winter resident coastwise; a few non-breeding birds summer offshore. Sept. 1–May 21; summer.

CONN.—Common migrant and winter resident; a few non-breeding birds summer offshore. Sept.–April; summer.

88. *ERISMATURA JAMAICENSIS* (Gmelin) Salvadori.

**Ruddy duck; Booby; Broadbill; Dumb-bird; Foolish coot.**

*Anas jamaicensis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 519.  
“JAMAICA.”



## ANSERES.

## ANATIDAE.

*Enligula rubida* Wils. Audubon, Birds of Amer., 1843, vol. 6, p. 324, pl. 399. Egg, Capen, 1886, pl. 21, fig. 4.

Fresh water chiefly; nests in marshes.

ME.—Uncommon migrant, and rare summer resident. April–May; (summer); Oct.–Nov.

N. H.—Uncommon migrant, rare in summer; may breed (Newfound Lake, July 11, 1901; Rye Beach, Aug. 22, 1879).

VT.—Uncommon migrant and rare summer resident (Lake Memphremagog).

MASS.—Common migrant; very rare summer resident (Cape Cod). Mar. 20–(summer); Sept. 30–Dec. 14 (Dec. 25, 1902, Woods Hole; Jan. 9, 1909, Jamaica Plain).

R. I.—Common migrant and occasional summer resident. Mar.; Oct. 1–Dec. 22.

CONN.—Common migrant, mainly in fall; rare in summer (June 5, 1893, flock of 5, Portland). Spring; Oct.–Dec. 25.

89. *NOMONYX DOMINICUS* (Linné) Ridgway.**Masked duck.**

Ridgway, Manual No. Amer. birds, 1887, p. 114.

*Anas dominica* Linné, Syst. nat., ed. 12, 1766, p. 201. "IN AMERICA MERIDIONAL."

*Erismatura dominica* Eyt. Baird, Cassin, and Lawrence, Birds of No. Amer., 1870, vol. 2, pl. 92.

Fresh water; nests among rushes.

VT.—Accidental visitor (or escape): Alburgh Springs on Lake Champlain.

MASS.—Accidental visitor: Malden, Aug. 27, 1889.

90. *CIEN HYPERBOREA* (Pallas) Boie.**Lesser snow goose; Mexican goose; Wavy; White brant.**

Ridgway, Manual No. Amer. birds, 1887, p. 115. Dresser,

## ANSERES.

## ANATIDAE.

Birds of Europe, 1871-81, vol. 6, pl. [417], lower fig. Egg, Seebohm, 1896, p. 30, pl. 11, fig. 3.

*Anser hyperboreus* Pall., Spicilegia zool., 1769, fasc. 6, p. 25.  
 "IN TERRIS BOREALIBUS, AD ORIENTEM 130°. LONGITUDINIS,  
 SIVE CIRCA LENAM ET IANAM FLUVIOS."

Salt and fresh water; nests on the ground.

ME.—Rare fall migrant. Oct. 2-Dec.

N. H.—Rare fall migrant: Lake Umbagog, Oct. 2, 1896; Seabrook.

VT.—Rare fall migrant: Lake Champlain; ? Lunenburg; St. Johnsbury.

MASS.—Rare migrant. April 13, 1908, Townsend; (? July) Oct. 15-Nov. 21.

R. I.—Rare fall migrant. Sept. 3-Oct. 16.

CONN.—Rare fall migrant: Saybrook, fall, 1875; Stratford, Oct. 5, 1842.

## 91. CUEN HYPERBOREA NIVALIS (Forster) Ridgway.

**Greater snow goose; Wavy.**

Ridgway, Manual No. Amer. birds, 1887, p. 115.

*Anas nivalis* Forster, Phil. trans. roy. soc. London, 1772, vol. 62, p. 413. "SEVERN RIVER," HUDSON BAY.

*Anser hyperboreus* Gmel. Audubon, Birds of Amer., 1843, vol. 6, p. 212, pl. 381, fig. 1.

Salt and fresh water; nests on the ground.

ME.—Now an accidental visitor: Georgetown, April 25, 1903; mouth of Kennebec River, April 7, 1890; Lubec, April 30, 1906.

MASS.—Now an accidental visitor: near Springfield (Morris); various doubtful records.

CONN.—Now an accidental visitor: Stratford, three wintered, 1867-68 (*vide* Averill); also listed by Linsley (1843), but may have been *C. hyperborea*.

## ANSERES.

## ANATIDAE.

92. *CHEN CAERULESCENS* (Linné) Gundlach.**Blue goose; Blue wavy.**

Ridgway, Manual No. Amer. birds, 1887, p. 111. Elliot, New and heretofore unfigured birds No. Amer., 1869, vol. 2, pl. 43.

*Anas caeruleus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 124. "CANADA."

Salt and fresh water; nests on the ground.

N. H.— Rare fall migrant: Lake Umbagog, Oct. 2, 1896.

MASS.— Rare fall migrant: West Gloucester, Oct. 20, 1876.

R. I.— Rare migrant: Westerly, Mar. 16, 1894. Other records in error for *C. hyperborea*.

93. *ANSER ALBIFRONS GAMBELI* (Hartlaub) Coues.**American white-fronted goose; Laughing goose.**

*Anser gambelli* (*sic*) Hartl., Rev. and mag. zool., ser. 2, 1852, vol. 4, p. 7. "TEXAS."

*Anser albifrons* Bechst. Audubon, Birds of Amer., 1843, vol. 6, p. 209, pl. 380.

Fresh and salt water; nests on the ground.

VT.— Doubtfully recorded.

MASS.— Accidental migrant (? formerly regular): Ipswich, Aug. 1907; Plymouth, Nov. 26, 1897; Quincy.

94. *BRANTA CANADENSIS* (Linné) Bannister.**Canada goose; Wild goose.**

*Anas canadensis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 123. "CANADA."

*Anser canadensis* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 178, pl. 376. Egg, Seebohm, 1896, p. 35, pl. 12, fig. 2.

Fresh and salt water; nests on the ground, rarely in trees.

## ANSERES.

## ANATIDAE.

ME.— Common migrant. (Feb. 28) Mar. 5–May; Oct. 13–Dec. 8.

N. H., VT.— Common migrant. Mar. 11–April; Oct.–Dec. 7.

MASS.— Common migrant, formerly summer resident (said to have bred at Lexington, late April, 1888); may winter rarely on south coast. Mar. 8–June 5; (Aug. 3) Sept. 21–Dec. 30 (Jan. 10; Jan. 31).

R. I.— Common migrant. Mar. 13–May 7; Sept. 17–Dec. 29.

CONN.— Common migrant and less common winter resident. Mar.–May 22; Oct.–winter.

95. *BRANTA CANADENSIS HUTCHINSII* (Richardson) Coues.**Hutchins's goose; ? Southern goose.**

Coues, Key to No. Amer. birds, ed. 5, 1903, vol. 2, p. 905.

*Anser hutchinsii* Rich., in Swainson and Rich., Fauna Boreali-Amer., 1831, vol. 2, p. 470. "MELVILLE PENINSULA," ARCTIC AMERICA. Audubon, Ornith. biogr., 1835, vol. 3, p. 226, pl. 277.

Fresh and salt water; nests on the ground.

ME.— Accidental migrant: Cape Elizabeth, Nov. 13, 1894; Portland, spring 1899; Rangeley.

MASS.— Supposed to occur as a migrant.

CONN.— ? Uncommon fall migrant; said to arrive later than *B. canadensis* and to pass at once south (Merriam, 1877).

96. *BRANTA BERNICLA GLAUCOGASTRA* (C. L. Brehm) Coues.**White-bellied brant; Blue brant; Brant; May brant.**

*Bernicla glaucogaster* C. L. Brehm, Handb. d. naturgesch. aller vögel Deutschl., 1831, p. 849. "AUS DEM HOHEN NORDEN IM HERBSTE AN DIE KÜSTEN DER DEUTSCHEN OSTSEE."

*Anser bernicla* Linné. Audubon, Birds of Amer., 1843, vol. 6, p. 203, pl. 379.

Salt water; nests on the ground.

## ANSERES.

## ANATIDAE.

ME.—Common coastwise migrant. April 27–May; Oct.–Nov.

N. H.—Uncommon coastwise migrant, casual inland.

VT.—Rare migrant.

MASS.—Common coastwise migrant; uncommon winter resident on south coast. Mar. 12–May 17 (June 2); Sept. 12–Dec. 12 (winter).

R. I.—Common migrant and rare winter resident offshore. Mar. 24–April 28; Nov.–Dec. 16 (winter).

CONN.—Common migrant and occasional winter resident offshore. April 14–May; Oct. 15–(winter).

97. *BRANTA NIGRICANS* (Lawrence) Bannister.**Black brant.**

*Anser nigricans* Lawr., Ann. lyceum nat. hist. New York, 1846, vol. 4, p. 171, pl. 12. "EGG HARBOR, N. J."

Seacoasts; nests on the ground.

MASS.—Accidental visitor: Chatham, spring, 1883.

98. *BRANTA LEUCOPSIS* (Bechstein) Bannister.**Barnacle goose.**

*Anas leucopsis* Bechst., Ornith. taschb. Deutschl., 1803, p. 424.  
NORTHERN REGIONS OF THE EARTH; GERMANY IN MIGRATION.

*Anser leucopsis* (Bechst.). Audubon, Birds of Amer., 1843, vol. 6, p. 200, pl. 378. Egg, Seebohm, 1896, p. 34, pl. 10, fig. 2.

Seacoasts and grassy flats; nests on the ground among rocks.

ME.—Doubtfully reported from near Portland about 1878.

VT.—Accidental visitor: Marshfield, 1878.

MASS.—Accidental visitor: North Chatham, Nov. 1, 1895.

CONN.—Recorded from Stonington (Linsley).

## ANSERES.

## ANATIDAE.

## 99. OLOR CYGNUS (Linné) Bonaparte.

**Whooping swan; Whooper; Wild swan.**

*Anas cygnus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 122. "IN EUROPA, AMERICA SEPTENTRIONALI."

*Cygnus musicus* Bechst. Dresser, Birds of Europe, 1871-81, vol. 6, p. [433], pl. [419], fig. 4. Egg, Seebohm, 1896, p. 29, pl. 7, fig. 1.

Seacoasts, lakes, swamps; nests on the ground.

ME.—Accidental visitor: Poke-a-moon-shine Lake, Washington Co., Sept. 10, 1903.

## 100. OLOR COLUMBIANUS (Ord) Stejneger.

**Whistling swan; American swan.**

*Anas columbianus* Ord, in Guthrie's Geogr., hist., and commercial grammar, 2d Amer. ed., 1815, p. 319. Based on Lewis and Clarke: "BELOW THE GREAT NARROWS OF THE COLUMBIA."

*Cygnus americanus* Sharpless. Audubon, Birds of Amer., 1843, vol. 6, p. 226, pl. 384.

Salt and fresh water; nests on the ground.

ME.—Unsatisfactorily recorded from mouth of Kennebec River Nov., 1881; and near Crawford Lake, Sept. 10, 1903.

N. H.—Rare migrant: Great Bay (*bis*), Dec. 16, 1902, and —; Seabrook, Oct. 18, 1878.

VT.—Rare migrant: Alburg, Lake Champlain.

MASS.—Rare migrant and winter resident (at Nantucket) formerly commoner. Oct. 16-Dec. 27 (winter-Mar. 4).

R. I.—Rare migrant: Quonocontaug Pond, Nov. 9, 1879; West-erly, Nov. 13, 1879.

CONN.—Rare migrant. Middle Mar.; fall-Nov. 2.

## ANSERES.

## ANATIDAE.

## 101. OLOR BUCCINATOR (Richardson) Wagler.

**Trumpeter swan.**

*Cygnus buccinator* Rich., in Swainson and Rich., Fauna Boreali-Amer., 1831, vol. 2, p. 464. "HUDSON'S BAY." Audubon, Birds of Amer., 1843, vol. 6, p. 219, pls. 382, 383.

Fresh and salt water; nests on the ground.

ME.— Once erroneously recorded for *O. cygnus*.

N. H.— Formerly a migrant (Belknap).

MASS.— Believed to have occurred formerly as a migrant.

R. I., CONN.— Doubtfully reported, and believed to have occurred formerly as a migrant.

## HERODIONES.

## IBIDES.

## IBIDIDAE.

## 102. GUARA ALBA (Linné) Stejneger.

**White Ibis.**

*Scolopax alba* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 145. "AMERICA."

*Ibis alba* (Linné). Audubon, Birds of Amer., 1843, vol. 6, p. 54, pl. 360.

Swamps and lagoons; nests in trees and reedy marshes.

VT.— Accidental visitor: South Woodstock, summer, 1878.

CONN.— Accidental visitor: near Milford, May 23, 1875.

## 103. PLEGADIS AUTUMNALIS (Hasselquist) Stejneger.

**Glossy Ibis.**

*Tringa autumnalis* Hasselq., Reise nach Paläst., deutsche ausg., 1762, p. 306. [EGYPT.]

HERODIONES.

IBIDES.

IBIDIDAE.

*Ibis falcinellus* Linné. Bonaparte, Amer. ornith., 1833, vol. 4, p. 23, pl. 23, fig. 1.

Marshes and borders of lagoons; nests among low bushes or reeds.

N. H.—Accidental visitor: Alton, Oct. 1858.

MASS.—Accidental visitor: noteworthy flights in May of 1850 and 1878.

CONN.—Accidental visitor: Middletown, May 16, 1850; Stratford (5 specimens) about 1836.

CICONIAE.

CICONIIDAE.

104. MYCTERIA AMERICANA Linné.

**Wood Ibis.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 140. "IN AMERICA CALIDIORE."

*Tantalus loculator* Linné. Audubon, Birds of Amer., 1843, vol. 6, p. 64, pl. 361.

Swamps and marshes; nests in trees.

ME.—Accidental visitor: Berwick, July 16, 1896.

N. H.—Erroneously recorded for Glossy ibis.

VT.—Accidental visitor: pair in summer, about 1897, near Burlington.

MASS.—Accidental visitor: (? Georgetown, June 19, 1880); Seekonk, July 17, 1896.

R. I.—Accidental visitor: Barrington, Aug. 8, 1896.



HERODIONES.

HERODIIL.

## ARDEIDAE.

105. *BOTaurus LENTIGINOSUS* (Montagu) Stephens.

**American bittern; Bog hen; Indian hen; Meadow hen; Stake driver; Thunder pump.**

*Ardea lentiginosa* Montagu, Ornith. dict., suppl., 1813, [p. 186] "Her." "THE WEST OF ENGLAND." Audubon, Birds of Amer., 1843, vol. 6, p. 94, pl. 365. Egg, Capen, 1886, pl. 19, fig. 14.

Fresh and salt marshes and borders of streams; nests on ground.

ME.—Uncommon migrant and summer resident. April 27–Oct. 28. Eggs, late May–June 6.

N. H.—Uncommon migrant and summer resident. Mar.–Nov. (Dec. 11). Eggs, May 27.

VT.—Uncommon migrant and summer resident. April 30–fall.

MASS.—Common migrant and less common summer resident; casual in winter. Mar. 31–Nov. 26 (Dec. 30; Jan. 5). Eggs, May 5–June 10.

R. I.—Common migrant and uncommon summer resident. April 6–Nov. 10. Eggs, May 23–June 13.

CONN.—Common migrant and uncommon summer resident. April–Nov.

106. *IXOBRYCHUS EXILIS* (Gmelin) A. O. U. comm.

**Least bittern.**

*Ardea exilis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 645. "JAMAICA." Audubon, Birds of Amer., 1843, vol. 6, p. 100, pl. 366. Egg, Capen, 1886, pl. 19, fig. 15.

Fresh marshes and swamps; nests among rushes or in bushes.

ME.—Rare summer resident in southern counties. (Mar. 4, 1891, Georgetown) May 12–Aug. (Nov. 13).

N. H.—Rare in summer; may breed in southeastern part: Hampton; Hampton Beach, 1869; Seabrook.

VT.—Said to have occurred "formerly."

HERODIONES.

HERODII.

## ARDEIDAE.

MASS.—Rare local summer resident, mainly near the coast. April 15–Sept. 1. Eggs, June 1–29.

R. I.—Common summer resident; casual in winter. (Mar. 1)–Sept. 14 (Feb. 28, 1881, Providence). Eggs, May 23–June 16.

CONN.—Uncommon local summer resident. Spring–Sept. Eggs, June 10–27.

## 107. ARDEA HERODIAS Linné.

**Great blue heron; Blue crane; Crane; Frog stabber; Poke.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 143. "AMERICA." Audubon, Birds of Amer., 1843, vol. 6, p. 122, pl. 369. Egg, Capen, 1886, pl. 19, fig. 11.

Swamps and borders of ponds; nests in trees.

ME.—Common migrant and local summer resident. Mar. 29–Nov. 12 (Dec. 9). Eggs, late April–May 16.

N. H.—Common migrant and, northward from L. Winnepesaukee, a local summer resident. April 5–Nov. (Dec. 21; Dec. 31, 1901, Randolph).

Vt.—Common migrant and, in northern part, a local summer resident. Summer–(Dec. 22, 1884, Milton).

MASS.—Common migrant and rare winter resident; occasional in summer but not now known to breed (formerly at Naushon). Mar. 15–June 4; (summer); July 30–Dec. 20 (winter).

R. I.—Common migrant. Mar.–May 4; July 15–Dec. 11.

CONN.—Common migrant, occasional in summer; rare winter resident. Mar. 15–May (June 2); Aug.–Dec. (winter).

## 108. HERODIAS EGRETTE (Gmelin) Cabanis.

**American egret.**

*Ardea egretta* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 629. "IN INSULA S. DOMINICI, INSULIS FALKLAND & AMERICA AUSTRALI AD

HERODIONES.

HERODII.

## ARDEIDAE.

LOUISIANAM." Audubon, Birds of Amer., 1843, vol. 6, p. 132, pl. 370.

Swamps and lowlands; nests in small trees.

ME.— Occasional summer visitor. April 7–Aug. 22.

N. H.— Rare summer visitor: near Newmarket, 1897.

VT.— Rare summer visitor: New Haven, two in Aug. 1882; Winooski River, summer, 1869.

MASS.— Occasional summer and fall visitor. April–Nov. 22.

R. I.— Occasional summer visitor; may have bred at West-quage Pond in 1894. June 1–Oct. 12.

CONN.— Occasional summer visitor. July–Oct.

## 109. EGRETTA CANDIDISSIMA (Gmelin) Gosse.

**Snowy heron.**

*Ardea candidissima* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 633.

"IN INUNDATIS PROPE CARTHAGENAM AMERICAE." Audubon, Birds of Amer., 1843, vol. 6, p. 163, pl. 374.

Swamps and lowlands; nests in small trees.

ME.— Erroneously accredited.

VT.— Accidental visitor: St. Albans Bay, two in Oct., 1890.

MASS.— Accidental visitor: near Boston, 1862; Nantucket, March, 1881; Northampton, 1887 (? *nec* Essex Co.).

CONN.— Accidental visitor: Hartford; Stratford ("seen").

## 110. FLORIDA CAERULEA (Linné) Baird.

**Little blue heron.**

*Ardea caerulea* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 143. "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer., 1843, vol. 6, p. 148, pl. 372.

Swamps and lowlands; nests in bushes.

## HERODIONES.

## HERODII.

## ARDEIDAE.

ME.— Accidental visitor. April 1–Sept.

N. H.— Accidental visitor: Amherst, April 28, 1897.

VT.— Doubtfully recorded.

MASS.— Occasional visitor. April 14–Aug

R. I.— Rare visitor. June 7–Aug.

CONN.— Rare visitor. May–Sept. 7.

## 111. BUTORIDES VIRESCENS (Linné) Bonaparte.

**Little green heron; Fly-up-the-creek; Green heron; Poke.**

*Ardea virescens* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 144.  
“AMERICA.” Audubon, Birds of Amer., 1843, vol. 6, p. 105, pl. 367.  
Egg, Capen, 1886, pl. 19, fig. 12.

Mainly freshwater streams and ponds; nests in small trees.

ME.— Uncommon migrant and summer resident in southern counties. May 2–Sept. 19.

N. H.— Uncommon summer resident in southern half, and in Connecticut valley. Late April–Oct. Eggs, June 16.

VT.— Uncommon summer resident in southern part. May–Oct.

MASS.— Common summer resident. (April 9 and 16) April 26–Oct. 16. Eggs, May 10–June 2.

R. I.— Common summer resident. April 20–Sept. 24. Eggs, May 21, 23.

CONN.— Common summer resident. Late April–Oct. Eggs, May 30–June 17.

## 112. NYCTICORAX NYCTICORAX NAEVIUS (Boddaert) Zeledon.

**Black-crowned night heron; Buttermunk; Qua-bird; Quok; Shitepoke; Squawk.**

*Ardea naevia* Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 56. Based on Buffon: “CAYENNE.”

HERODIONES.

HERODII.

## ARDEIDAE.

*Ardea nycticorax* Linné. Audubon, Birds of Amer., 1843, vol. 6, p. 82, pl. 363. Egg, Capen, 1886, pl. 19, fig. 13.

Swamps and marshes; nests in colonies in trees, rarely on the ground.

ME.— Common summer resident, chiefly near the coast. Spring–Sept. 16. Eggs, May 26.

N. H.— Uncommon summer resident north to southern White Mt. valleys. April–Oct.

VT.— Uncommon summer resident near lakes and streams.

MASS.— Common migrant and summer resident; rare winter resident near coast. Mar. 27–Nov. 1 (winter). Eggs, May 2–June 13.

R. I.— Common summer and rare winter resident. April 1–Sept. 23 (winter). Eggs, May 7–June 7.

CONN.— Common migrant and less common summer resident. April–Oct. Eggs, April 17.

113. *NYCTANASSA VIOLACEA* (Linné) Sharpe.

**Yellow-crowned night heron.**

*Ardea violacea* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 143. "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer., 1843, vol. 6, p. 89, pl. 364.

Swamps and lowlands; nests in low trees.

ME.— Accidental visitor: Deering, April 13, 1901; Portland, April 11, 1906.

MASS.— Accidental visitor: six instances. July–Oct.

R. I.— Accidental visitor: Newport, Aug. 1892; Tiverton, April 23, 1886.

## PALUDICOLAE.

## GRUES.

## GRUIDAE.

## 114. GRUS AMERICANA (Linné) Vieillot.

**Whooping crane.**

Audubon, Birds of Amer., 1842, vol. 5, p. 188, pl. 313.

*Ardea americana* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 142.

“IN AMERICA SEPTENTRIONALI.”

Freshwater marshes and open lowlands; nests in marshes.

VT., MASS.— Supposed to have been a migrant in colonial times.

## 115. GRUS CANADENSIS (Linné) Temminck.

**Little brown crane.**

Ridgway, Manual of No. Amer. birds, 1887, p. 135.

*Ardea canadensis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 141.

“IN AMERICA SEPTENTRIONALI.”

*Grus fraterculus* Cassin. Baird, Cassin, and Lawrence, Expl. and surveys for railroad to Pacific, 1858, birds, pl. 37.

Grassy flats and marshes; nests on the ground.

R. I.— Accidental visitor: Natick Hill, Oct. 9, 1889.

## 116. GRUS MEXICANA (Müller) Vieillot.

**Sandhill crane.**

Coues, Key to No. Amer. birds, ed. 5, 1903, vol. 2, p. 848.

*Ardea (grus) mexicana* Müll., Linné's Vollstand. natursyst., suppl., 1776, p. 110. “MEXICO.”

*Grus americana* Forst. Audubon, Birds of Amer., 1842, vol. 5, pl. 314.

Freshwater marshes and open lowlands; nests in marshes.

N. H.— Believed to have occurred formerly; now accidental visitor: Wakefield, 1896 or 1897.

VT.— Believed to have occurred formerly as a migrant; now accidental visitor: Lunenburg.

MASS., CONN.— Believed to have occurred formerly as a migrant.

## PALUDICOLAE.

## RALLI.

## RALLIDAE.

117. *RALLUS ELEGANS* Audubon.

**King rail; Fresh-marsh hen; Royal rail.**

Audubon, Ornith. biogr., 1835, vol. 3, p. 27, pl. 203. "SOUTHERN STATES." Coues, Key No. Amer. birds, ed. 5, 1903, vol. 2, p. 854.

Freshwater marshes; nests on the ground.

ME.— Occasional fall visitor. Sept. 19–Dec. 17.

MASS.— Occasional visitor at all seasons; may breed.

R. I.— Rare visitor (four records). Jan., Feb., Mar., fall.

CONN.— Rare summer resident (Saybrook, Stratford) and occasional at other seasons.

118. *RALLUS CREPITANS* Gmelin.

**Clapper rail; Salt-marsh hen.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 713. "IN NOVEBORACO." Audubon, Birds of Amer., 1842, vol. 5, p. 165, pl. 310. Egg, Capen, 1886, pl. 20, figs. 1, 2.

Saltwater marshes; nests on the ground.

ME.— Accidental visitor in southern counties: Popham Beach, Oct. 12 and (25), 1900; Sabattus Pond, 1874; ? York and Cumberland Cos.

N. H.— Accidental visitor: Portsmouth.

VT.— Accidental visitor: Burlington (Conger).

MASS.— Occasional visitor, may breed.

R. I.— Doubtfully recorded.

CONN.— Rare visitor and summer resident (New Haven Stratford).

119. *RALLUS VIRGINIANUS* Linné.

**Virginia rail; Water-hen.**

Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 263. "IN AMERICA

## PALUDICOLAE.

## RALLI.

## RALLIDAE.

SEPTENTRIONALI." Audubon, Birds of Amer., 1842, vol. 5, p. 174, pl. 311. Egg, Capen, 1886, pl. 20, figs. 3, 4.

Fresh and brackish marshes; nests on the ground.

ME.—Rare summer resident. Late April–Oct.

N. H.—Rare summer resident south of White Mts. April–Oct. 1. Eggs, May 21–28.

VT.—Rare summer resident locally: Burlington; Rutland Co.

MASS.—Summer resident, more common near coast; rare winter resident (Cape Cod). April 11–Nov. 27 (Jan. 1, 1891, Worcester) (winter). Eggs, May 12–June 1.

R. I.—Common migrant and summer resident. April–Nov. 2. Eggs, May 29.

CONN.—Common migrant and summer resident. Eggs, June 6.

## 120. PORZANA CAROLINA (Linné) Baird.

**Sora; Carolina rail; Meadow-hen; Rail-bird.**

*Rallus carolinus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 153. "IN AMERICA SEPTENTRIONALI."

*Ortygometra carolinus* (Linné). Audubon, Birds of Amer., 1842, vol. 5, p. 145, pl. 306. Egg, Capen, 1886, pl. 20, fig. 5.

Fresh and brackish marshes; nests on the ground.

ME.—Uncommon summer resident. Late April–Oct. 26. Eggs, –July 27.

N. H.—Uncommon summer resident northward to White Mt. valleys. May–Oct.

VT.—Uncommon summer resident.

MASS.—Common migrant and local summer resident. (Mar. 20) April 14–Nov. 1 (Dec. 20). Eggs, June 9–11.

R. I.—Common migrant and summer resident. (Mar. 2) April–Nov. 10. Eggs, May 24.

CONN.—Common migrant and summer resident. April–Oct. (Dec. 29, 1881, Hartford).



PALUDICOLAE.

RALLI.

RALLIDAE.

121. *COTURNICOPS NOVEBORACENSIS* (Gmelin) Bonaparte.**Yellow rail.**

*Fulica noveboracensis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 701.  
"IN NOVEBORACO."

*Rallus noveboracensis* (Lath.). Bonaparte, Amer. ornith., 1833,  
vol. 4, p. 136, pl. 27, fig. 2. Egg, Capen, 1886, pl. 20, fig. 6.

Freshwater marshes; nests on the ground.

ME.— Rare migrant and said to have bred near Calais. May 31;  
Oct. 4–Nov. 15.

N. H.— Rare migrant: Hampton.

VT.— Rare migrant: Newport; northern Vt.; Rutland Co.

MASS.— Rare migrant and possible summer resident. April–  
May 26; Sept. 2–Oct. 22.

R. I.— Rare migrant (five records). Sept. 28.

CONN.— Rare spring and uncommon fall migrant; said to have  
bred at Middletown, 1874, 1875. (Mar. 24, Gaylordsville); Sept.–  
Oct. 14 (Nov. 10).

122. *CRECISCUS JAMAICENSIS* (Gmelin) Cabanis.**Black rail; Little black rail.**

*Rallus jamaicensis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 718.  
"JAMAICA."

*Porzana jamaicensis* (Gmel.). Chapman, Handb. birds east.  
No. Amer., 1895, p. 144; Allen, Auk, 1900, vol. 17, pl. 1.

Freshwater marshes; nests on the ground.

ME.— Doubtfully recorded from Scarborough, Oct. 4, 1881.

MASS.— Very rare visitor and summer resident (at Chatham,  
Hazardville). May 16–Sept. 20. Eggs, May.

CONN.— Very rare summer resident (at Lyme, Saybrook);  
also taken at Hazenville. Eggs, June 6–July 10.

PALUDICOLAE.

RALLI.

RALLIDAE.

123. *CREX CREX* (Linné) Sharpe.**Corn crane; Daker-hen; European land rail.**

Coues, Key to No. Amer. birds, ed. 5, 1903, vol. 2, p. 859.

*Rallus crex* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 153. "IN EUROPAE AGRIS, CARECTIS."*Crex pratensis* Bechst. Dresser, Birds of Europe, 1871-81, vol. 7, pl. [499]. Egg, Seebohm, 1896, p. 83, pl. 22, fig. 9.

Grassy meadows and grain fields; nests on the ground.

ME.—Accidental from Europe: near Falmouth, Oct. 14, 1889.

R. I.—Accidental visitor: Cranston, 1857.

CONN.—Accidental visitor: Saybrook, Oct. 20, 1887.

124. *IONORNIS MARTINICA* (Linné) Reichenbach.**Purple gallinule.***Fulica martinica* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 259. "IN MARTINICAE INUNDATIS."*Gallinula martinica* (Linné). Audubon, Birds of Amer., 1842, vol. 5, p. 128, pl. 303.

Freshwater marshes; nests among rushes.

ME.—Accidental visitor. April 11-Dec. 17.

N. H.—Accidental visitor: Dover; Rye.

VT.—Doubtfully recorded.

MASS.—Occasional visitor, mainly in April and Oct.-Nov., once in June.

R. I.—Occasional visitor. Jan.; May; June 8, 1902, Seacommet Point; Aug.; fall.

CONN.—Rare visitor: Middletown, 1855.

125. *GALLINULA GALEATA* (Lichtenstein) Bonaparte.**Florida gallinule; Pond hen; Red-billed mud-hen.***Crex galeata* Licht., Verz. d. doubletten zool. mus. Berlin, 1823, p. 80. "SAN PAULO," BRAZIL.

## PALUDICOLAE.

## RALLI.

## RALLIDAE.

*Gallinula chloropus* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 132, pl. 304. Egg, Capen, 1886, pl. 20, figs. 7, 8.

Freshwater marshes; nests among rushes near the ground.

ME.—Rare migrant and (?) summer resident. May 5; Sept. 20–Oct. 16.

N. H.—Accidental visitor: Rollinsford.

VT.—Rare local summer resident at Lake Bomoseen, Lake Champlain, St. Albans. April 28–fall. Eggs, May 28–June 5.

MASS.—Rare local summer resident, mainly near coast. April 29–Oct. 25 (Nov. 9). Eggs, June 5–July 17.

R. I.—Uncommon local summer resident. May–Nov. 29.

CONN.—Uncommon local summer resident. Eggs, June 25 (Stratford).

## 126. FULICA AMERICANA Gmelin.

**American coot; Blue Peter; Coot; Marsh hen; Mud-hen;  
Pond crow; Pond hen.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 704. "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer., 1842, vol. 5, p. 138, pl. 305. Egg, Capen, 1886, pl. 20, fig. 9.

Freshwater ponds and marshes mainly; nests among reeds.

ME.—Uncommon migrant, mainly in fall. (Mar. 23) April 14–May 13; Oct. 26–Nov.

N. H.—Uncommon migrant, mainly in southern part. Spring; late Aug.–Oct. 15.

VT.—Uncommon migrant and rare summer resident (Lake Bomoseen).

MASS.—Rare spring and uncommon fall migrant; rare winter resident near coast; may breed rarely (pair, Cheshire reservoir, June 21, 1892). Mar. 29–April 25; (June 21); (Aug. 16) Sept. 3–Nov. (winter).

## PALUDICOLAE.

## RALLI.

## RALLIDAE.

R. I.—Common migrant, mainly in fall. April–May 28; Sept. 26–Dec. 20.

CONN.—Common migrant, mainly in fall. Spring; Sept. 29–Nov. 14.

## LIMICOLAE.

## PHALAROPODIDAE.

127. PHALAROPUS FULICARIUS (Linné) Bonaparte.

**Red phalarope; Bank-bird; Brown bank-bird; Gray phalarope; Gulf-bird; Sea goose; Whale-bird.**

Audubon, Birds of Amer., 1842, vol. 5, p. 291, pl. 339. Egg, Capen, 1886, pl. 18, fig. 11; pl. 19, fig. 1.

*Tringa fulicaria* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 148. "AMERICA."

Chiefly maritime; nests on the ground in marshes.

ME.—Uncommon migrant, mainly offshore, and rare summer resident (near Calais). May 8–17; (summer); July 25–Oct. 17.

N. H.—Migrant, mainly offshore; sometimes common. May; Oct.

VT.—Rare migrant at Lake Champlain.

MASS.—Offshore migrant, irregularly common. May 1–26; Aug. 31–Nov. 24.

R. I., CONN.—Offshore migrant, sometimes common. Late May; Sept. 26–Dec. 20.

128. LOBIPES LOBATUS (Linné) Baird, Brewer, & Ridgway.

**Northern phalarope; Bank-bird; Jersey goose; Mackerel goose; Red-necked phalarope; Sea goose; Sea snipe; Whale-bird; White bank-bird.**

*Tringa lobata* Linné, Syst. nat., ed. 10, 1758, vol. 1, pp. 148, 824. "IN AMERICA SEPTENTRIONALI, LAPPONIA."

*Lobipes hyperboreus* Lath. Audubon, Birds of Amer., 1842, vol. 5, p. 295, pl. 340. Egg, Seebohm, 1896, p. 131, pl. 38, fig. 1.

Chiefly maritime; nests on the ground in marshes.

## LIMICOLAE.

## PHALAROPODIDAE.

ME.—Common migrant, mainly offshore. (Feb.) May 3–June 15; July 24–Nov.

N. H.—Common migrant, mainly offshore. May; Aug. 9–Sept. 22.

MASS.—Common migrant, mainly offshore. May 1–26; Aug. 10–Oct. 13.

R. I.—Common migrant, mainly offshore. May; Aug. 16–Sept. 15.

CONN.—Offshore migrant. Spring?; Aug. 4–Sept. 27 (Dec. 25).

## 129. STEGANOPUS TRICOLOR Vieillot.

**Wilson's phalarope.**

Vieill., Nouv. dict. d'hist. nat., 1819, vol. 32, p. 136. "PARAGUAY."

*Phalaropus wilsonii* Sabine. Swainson and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 405, pl. 69.

Chiefly freshwater marshes; nests on the ground.

ME.—Accidental visitor: Sabbattus Pond, Sept. or Oct. 1906; Scarborough, three, June 9, 1891.

N. H.—Accidental visitor: Rye Beach, Aug. 15, 1872.

MASS.—Accidental visitor (five records). May 20; Aug. 18–31.

R. I.—Accidental (fall) visitor (five records). Aug. 2–Sept. 13.

CONN.—Accidental visitor: Bridgeport.

## RECURVIROSTRIDAE.

## 130. RECURVIROSTRA AMERICANA Gmelin.

**American avocet.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 693. "IN AMERICA SEPTENTRIONALI & NOVA HOLLANDIA." Audubon, Birds of Amer., 1843, vol. 6, p. 24, pl. 353.

## LIMICOLAE

## RECURVIROSTRIDAE.

Open lowlands and marshes; nests on the ground.

ME.—Accidental visitor: Cape Elizabeth, Nov. 5, 1878.

VT.—Accidental visitor: St. Albans, fall, 1875, and 1890; (*nec* Rutland, 1882).

MASS.—Accidental visitor: Ipswich, three, Sept. 13, 1896; Lynn; Natick, Oct. 19, 1880.

CONN.—Accidental visitor: between Saybrook and East Lyme, 1871.

## 131. HIMANTOPUS MEXICANUS (Müller) Ord.

**Black-necked stilt.**

*Charadrius mexicanus* Müll., Linné's Vollstand. natursyst., suppl., 1776, p. 117. "MEXICO."

*Himantopus nigricollis* Vieill. Audubon, Birds of Amer., 1843, vol. 6, p. 31, pl. 354.

Salt marshes and lagoons; nests on the ground.

ME.—Accidental visitor: Rockland, May, 1889; (*nec* Boardman, New Brunswick).

N. H.—Accidental visitor: Rye Beach.

MASS.—Accidental visitor: Lynn; Mass. coast.

## SCOLOPACIDAE.

## 132. PHILOHELA MINOR (Gmelin) Gray.

**American woodcock; Bog-sucker; Woodcock.**

*Scolopax minor* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 661. "IN AMERICA, A SEPTEMBRI INDE IN CAROLINAE, AB APRILIS FINE IN NOVEBORACI SILVIS HUMIDIS."

*Microptera americana* Audubon, Birds of Amer., 1843, vol. 6, p. 15, pl. 352. Egg, Capen, 1886, pl. 19, fig. 2.

Freshwater swamps, damp woods and thickets; nests on the ground.

## LIMICOLAE.

## SCOLOPACIDAE.

ME.— Common migrant and summer resident; possibly winters. (Mar. 10) Mar. 23–Nov. 30 (Dec. 18) (Feb., two instances). Eggs, April 26–May 19.

N. H.— Uncommon migrant and summer resident. Late Mar.–Oct. Eggs, May 10.

VT.— Uncommon migrant and summer resident. Mar. 27–Oct. Eggs, May 4.

MASS.— Common migrant and uncommon summer resident; rare in winter. Mar. 4–Nov. 25 (winter). Eggs, April 11–May 6.

R. I.— Common migrant and now a rare summer resident. Mar.–Dec. 4. Eggs, April 5–May 17.

CONN.— Common migrant, uncommon summer and winter resident. Feb. 26–Nov.; winter. Eggs, (Mar. 30) April 3–June 3; July 20 (2d brood).

133. *GALLINAGO DELICATA* (Ord) A. O. U. check-list.

**Wilson's snipe; Alewife-bird; Common snipe; English snipe; Jack snipe.**

*Scolopax delicata* Ord, Wilson's Ornith., 1825, vol. 9, p. ccviii. "KENTUCKY, . . . ILLINOIS, . . . PENNSYLVANIA."

*Scolopax wilsonii* Temm. Audubon, Birds of Amer., 1842, vol. 5, p. 339, pl. 350. Egg, Capen, 1886, pl. 19, fig. 3.

Fresh and brackish meadows and marshes; nests on the ground.

ME.— Common migrant and rare summer resident. April 19–May 15; summer; Sept.–Nov. 15.

N. H.— Common migrant, mainly coastwise; very rare winter resident (Nashua). Mar.–May 7; July 22–mid-Dec. (winter).

VT.— Uncommon migrant and rare winter resident (Rutland Co.); ? "summer resident."

MASS.— Common migrant, rare winter resident; rare and irregular summer resident (at Carlisle, Concord). (Mar. 8) Mar. 20–May 18; (summer); Aug. 1–Nov. 30 (winter).

## LIMICOLAE.

## SCOLOPACIDAE.

R. I.—Common migrant and rare winter resident. Mar. 9–May 6; (July 29) Aug. 12–Nov. 14 (Dec., Jan.).

CONN.—Common migrant, rare summer and winter resident. Mar. 18–Dec.; winter. Eggs, May 13 (Portland).

134. *MACRORHAMPHUS GRISEUS* (Gmelin) Leach.

**Dowitcher; Brownback; Deutscher; Fool plover; German snipe; Grayback; Gray snipe; Red-breasted snipe; Robin snipe.**

*Scolopax grisea* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 658. "IN NOVEBORACI MARITIMIS."

*Scolopax noveboracensis* Gmel. Audubon, Birds of Amer., 1843, vol. 6, p. 10, pl. 351. Egg, Seebohm, 1896, p. 143, pl. 41, fig. 8.

Chiefly maritime marshes and mudflats; nests on the ground.

ME.—Common migrant coastwise. May 21–29; late July–Sept. (Oct. 29).

N. H.—Migrant, common in fall. May 20–25; July 20–Sept.

MASS.—Rare spring and uncommon fall migrant, mainly coastwise. May 1–June 7 (June 18); July 5–Sept. 30.

R. I.—Uncommon migrant coastwise. May; July 5–Oct. 20.

CONN.—Uncommon migrant, mainly in fall.

135. *MACRORHAMPHUS SCOLOPACEUS* (Say) Lawrence.

**Long-billed dowitcher.**

Elliot, New and unfigured birds No. Amer., 1869, vol. 2, pl. 40.

*Limosa scolopacea* Say, Long's exped. to Rocky Mts., 1823, vol. 1, p. 170. "NEAR THE BOWYER CREEK," IOWA.

Chiefly maritime marshes and mudflats; nests on the ground.

MASS.—Rare fall migrant. Aug. 29–Nov. 3.

R. I.—Rare fall migrant; Middletown, Oct. 8, 1890.



## LIMICOLAE.

## SCOLOPACIDAE.

## 136. MICROPALAMA HIMANTOPUS (Bonaparte) Baird.

**Stilt sandpiper; Bastard yellow-leg; Long-legged sandpiper;  
Mongrel.**

*Tringa himantopus* Bonap., Ann. lyceum nat. hist. New York, 1826, vol. 2, p. 157. "LONG BRANCH, NEW JERSEY." Bonap., Amer. ornith., 1833, vol. 4, p. 89, pl. 25, fig. 3.

Marshes and mudflats; nests on the ground.

ME.—Rare fall migrant coastwise. July 19–Sept. 26 (Oct. 2, 1908, Pine Point).

N. H.—Rare fall migrant coastwise. July 31–Aug. 28.

MASS.—Uncommon fall migrant coastwise. July 19–Sept. 29.

R. I.—Very rare spring and uncommon fall migrant. May 9, 1895, Sakonnet Point; July 6–Sept. 19.

CONN.—Rare migrant: New Haven, Sept. 16, 1886.

## 137. TRINGA CANUTUS Linné.

**Knot; Ash-colored sandpiper; Blue plover; Grayback; Gray snipe; Red-breasted plover or sandpiper; Robin snipe; Silver plover.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 149. "IN EUROPA." Egg, Seebohm, 1896, p. 144, pl. 43, fig. 10.

*Tringa islandica* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 254, pl. 328.

Sandy beaches; nests on the ground.

ME., N. H.—Uncommon migrant coastwise. May 24–June 11; Aug.–Sept.

MASS.—Common migrant coastwise, casual inland (Springfield); rare winter resident. May 11–June 5 (June 25); July 13–Nov. 8 (winter).

R. I.—Common migrant coastwise. May; July 11–Sept. 14.

CONN.—Common migrant coastwise.

## LIMICOLAE.

## SCOLOPACIDAE.

138. *ARQUATELLA MARITIMA* (Brünnich) Coues.

**Purple sandpiper; Rock snipe; Winter peep; Winter snipe.**

*Tringa maritima* Brünn., Ornith. borealis, 1764, p. 54. "CHRISTIANSÖE" I., DENMARK, AND "NORVEGIA." Audubon, Birds of Amer., 1842, vol. 5, p. 261, pl. 330. Egg, Seebohm, 1896, p. 146, pl. 43, figs. 2, 5.

Rocky islands and coasts of the sea; nests on the ground.

ME.—Common migrant and winter resident coastwise; accidental in summer. Sept. 15–April (Aug. 6, 1907, Metinic Green Id.; Aug. 11, 1902, Metinic reef).

N. H.—Rare winter visitor coastwise.

VT.—Doubtfully recorded.

MASS.—Common fall and winter visitor coastwise. (July 30) Sept. 6–April 19 (May 11).

R. I.—Common winter resident on rocky coasts. Sept. 13–April 15.

CONN.—Common winter resident on rocky islands coastwise.

139. *PISOBIA MACULATA* (Vieillot) A. O. U. comm.

**Pectoral sandpiper; Brown-back; Creaker or Krieker; Grass-bird; Grass snipe; Jack snipe; Pert; Squatter.**

*Tringa maculata* Vieill., Nouv. dict. d'hist. nat., 1819, vol. 34, p. 465. "AUX ÎLES ANTILLES ET DANS LES PARTIES MÉRIDIONALES DES ÉTATS-UNIS."

*Tringa pectoralis* Bonap. Audubon, Birds of Amer., 1842, vol. 5, p. 259, pl. 329. Egg, Seebohm, 1896, p. 148, pl. 43, fig. 12.

Marshes and mudflats; nests on the ground.

ME.—Rare spring and common fall migrant, mainly coastwise. (Spring); Aug. 13–Oct. 5.

N. H.—Migrant, common in fall, mainly coastwise. July–Oct. 10.

## LIMICOLAE.

## SCOLOPACIDAE.

VT.— Uncommon fall migrant.

MASS.— Rare spring and common fall migrant. (April 4) April 9–May 16 (May 23); July 15–Nov. 11.

R. I.— Rare spring and common fall migrant. April 25; July 16–Oct. 20.

CONN.— Rare spring and common fall migrant.

140. *PISOBIA FUSCICOLLIS* (Vieillot) A. O. U. comm.

**White-rumped sandpiper; Bonaparte's sandpiper; Bull-peep; Schinz's sandpiper.**

*Tringa fuscicollis* Vieill., Nouv. dict. d'hist. nat., 1819, vol. 34, p. 461. "PARAGUAY." Egg, Seebohm, 1896, p. 145, pl. 43, fig. 8.

*Actodromas fuscicollis* (Vieill.). Baird, Brewer, and Ridgway, Water birds of No. Amer., 1884, vol. 1, p. 227, fig.

*Tringa schiuzii* Brehm. Audubon, Ornith. biogr., 1835, vol. 3, pl. 278.

Beaches and marshy shores; nests on the ground.

ME.— Uncommon migrant, mainly coastwise. May 29, 30; late July–Oct. 23 (Nov. 18).

N. H.— Uncommon migrant, mainly coastwise. ? Spring; July–Oct. 14.

VT.— Uncommon fall migrant. Aug.–Sept.

MASS.— Rare spring and common fall migrant mainly coastwise. May 15–June 5; July 10–Nov. 10.

R. I.— Uncommon fall migrant. July 11–Oct.

CONN.— Uncommon migrant. Aug. 31.

141. *PISOBIA BAIRDII* (Coues) A. O. U. comm.

**Baird's sandpiper.**

*Actodromas bairdii* Coues, Proc. acad. nat. sci. Phila., 1861, p. 194. "NORTH AMERICA." Baird, Brewer, and Ridgway, Water birds No. Amer., 1884, vol. 1, p. 230, colored fig. of head.

## LIMICOLAE.

## SCOLOPACIDAE.

Beaches and marshes; nests on the ground.

ME.—Rare fall migrant. Aug. 7–Sept. 20 (Nov. 1, 1881, Bangor).

N. H.—Rare fall migrant. Aug. 26–Sept. 5.

MASS.—Rare fall migrant. July 27–Oct. 1.

R. I.—Rare fall migrant (seven records). Aug. 26–Sept. 17.

CONN.—Rare fall migrant: New Haven, Oct. 19, 1889, Oct. 28, 1887; Stratford, Nov. 3, 1888.

142. *PISOBIA MINUTILLA* (Vieillot) A. O. U. comm.

**Least sandpiper; Bumble bee; Meadow ox-eye; Mud peep; Peep; Wilson's sandpiper.**

*Tringa minutilla* Vieill., Nouv. dict. d'hist. nat., 1819, vol. 34, p. 466. "EN AMÉRIQUE JUSQU'AU DELÀ DU CANADA." Egg, Seebohm, 1896, p. 149, pl. 44, fig. 8.

*Tringa pusilla* Wils. Audubon, Birds of Amer., 1842, vol. 5, p. 280, pl. 337.

Marshes and mudflats; nests on the ground.

ME.—Common migrant. May 13–24 (June); July 5–Oct. 1 (Oct. 15).

N. H.—Common migrant. Late May–June 3; July 10–Sept. VT.—Uncommon migrant.

MASS.—Common migrant. (April 18) May 5–June 7; (summer) July 6–Sept. 21 (Oct. 13).

R. I.—Common migrant. (April 25)–May 26; July 15–Oct.

CONN.—Common migrant. May–June; Aug.–Sept.

143. *PELIDNA ALPINA* (Linné) C. L. Brehm.

**European dunlin.**

Sharpe, Cat. birds British mus., 1896, vol. 24, p. 602. Dresser, Birds of Europe, 1871–81, vol. 8, p. [21], pl. [548].

## LIMICOLAE.

## SCOLOPACIDAE.

*Tringa alpina* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 119. "IN LAPONIA." Egg, Seebohm, 1896, p. 144, pl. 43, figs. 7, 9.

Beaches and marshes; nests on the ground.

MASS.—Accidental from Old World: Chatham, Aug. 11, 1900.

144. *PELIDNA ALPINA SAKHALINA* (Vieillot) Buturlin.

**American dunlin; Black-bellied sandpiper; Black-breast; Brant-bird; Crooked-billed snipe; Fall snipe; Lead-back; Ox-bird; Ox-eye; Red-back; Red-backed sandpiper.**

*Scolopax sakhalina* Vieill., Nouv. dict. d'hist. nat., 1816, vol. 3, p. 359. "EN RUSSIE."

*Tringa alpina* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 266, pl. 332.

Beaches and marshes; nests on the ground.

ME.—Rare spring and commoner fall migrant coastwise. Late May; Oct.—Nov. 17.

N. H.—Uncommon coastwise migrant.

MASS.—Rare spring and common fall migrant coastwise. (? Mar. 15) late April—May 20 (June 18); Sept. 1—Dec. 13 (Dec. 24).

R. I.—Rare spring and common fall migrant coastwise. May; Aug. 29—Oct. 29.

CONN.—Rare spring and common fall migrant; "may winter."

145. *EROLIA FERRUGINEA* (Brünnich) A. O. U. comm.

**Curlew sandpiper.**

*Tringa ferruginea* Brünn., Ornith. bor., 1764, p. 53. "ISLANDIA & CHRISTIANSÖE" ISLAND.

*Tringa subarquata* Temm. Audubon, Birds of Amer., 1842, vol. 5, p. 269, pl. 333.

Seacoasts and tundras; nests on the ground.

## LIMICOLAE.

## SCOLOPACIDAE.

ME.— Accidental visitor: Pine Point, Cumberland Co., Sept. 15, 1881.

MASS.— Accidental visitor: five records. May; Aug.

R. I.— Doubtfully recorded.

CONN.— Accidental visitor: four records. (June 1874, New Haven); Aug. 30–Oct. 3.

146. *EREUNETES PUSILLUS* (Linné) Cassin.

**Semipalmated sandpiper; Black-legged peep; Peep; Sand ox-eye; Sand peep.**

*Tringa pusilla* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 252.  
“IN DOMINGO.”

*Tringa semipalmata* Wils. Audubon, Birds of Amer., 1842, vol. 5, p. 277, pl. 336.

Sand beaches and marshes; nests on the ground.

ME.— Common migrant. May–June 5; July 29–Sept. 28 (Oct.).

N. H.— Common migrant. May; July–early Sept.

VT.— Uncommon migrant to larger lakes.

MASS.— Common migrant; non-breeding birds occur in summer coastwise. (May 1) May 13–June 14; (summer); July 3–Oct. 8 (Oct. 30).

R. I.— Common migrant. May 13–June 2; July 14–Oct. 10.

CONN.— Common migrant. May–early June; July–Oct.

147. *EREUNETES MAURI* Cabanis.

**Western semipalmated sandpiper**

Cab., Journ. f. ornith., 1856, p. 419. “CUBA.”

*Ereunetes pusillus occidentalis* (Lawrence). Baird, Brewer, and Ridgway, Water birds of No. Amer., 1884, vol. 1, p. 206, fig. of head.

Sand beaches and marshes; nests on the ground.

## LIMICOLAE.

## SCOLOPACIDAE.

N. H.— Rare fall migrant: Hampton, Oct. 10, 1899.

MASS.— Uncommon fall migrant. July 19–Sept. 20.

R. I.— Rare fall migrant: Middletown, Aug. 25, and Aug. 29, 1899.

CONN.— Rare fall migrant: Lyme, Sept. 4, 1889 (six).

148. *CALIDRIS LEUCOPHAEA* (Pallas) A. O. U. comm.

**Sanderling; Beach-bird; Beach plover; Bull-peep; Gray-back; Ruddy plover; Surf snipe; Whitey; Whiting.**

*Tringa leucophaea* Pallas, in Vroeg's Catalogue, 1764, p. 32.

## NORTH SEA COAST.

*Tringa arenaria* Bonap. Audubon, Birds of Amer., 1842, vol. 5, p. 287, pl. 338. Egg, Seebohm, 1896, p. 150, pl. 41, fig. 4.

Sand beaches; nests on the ground.

ME., N. H.— Uncommon spring and common fall migrant coastwise. May 5–30; late July–Oct. 13 (Nov. 5, Portland).

VT.— Rare migrant. Sept.

MASS.— Common coastwise migrant and, south of Cape Cod, a winter resident. (Late April) May 19–June 4 (June 24); July 6–Dec. 6 (Dec. 13); winter.

R. I.— Common migrant, casual inland; probably a rare winter resident. (Mar. 20) May–June; Aug. 1–Nov. 5.

CONN.— Common migrant coastwise. May; July–Oct.

149. *LIMOSA FEDOA* (Linné) Sabine.

**Marbled godwit; Brown marlin; Red curlew.**

Audubon, Birds of Amer., 1842, vol. 5, p. 331, pl. 348.

*Scolopax fedoa* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 146. "IN AMERICA SEPTENTRIONALIS."

Beaches and marshes; nests on the ground.

ME.— Rare migrant. May, 1884 (Scarborough); Aug. 8–Sept. 13.

## LIMICOLAE.

## SCOLOPACIDAE.

N. H.— Rare migrant: Rye Beach, Aug. 27, 1868.

Vt.— Formerly a migrant.

MASS.— Rare migrant. May 20–23; July 17–Sept. 5.

R. I.— Rare migrant. Aug. 6–Oct. 2.

CONN.— Rare migrant. Aug.

## 150. LIMOSA HAEMASTICA (Linné) Coues.

**Hudsonian godwit; Black-tail; Goose-bird; Ring-tailed marlin;  
Spot-rump.**

*Scolopax haemastica* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 147.  
“IN AMERICA SEPTENTRIONALI.”

*Limosa hudsonica* Lath. Audubon, Birds of Amer., 1842, vol. 5,  
p. 335, pl. 349.

Beaches and marshes; nests on the ground.

ME.— Rare fall migrant coastwise.

N. H.— Rare fall migrant coastwise. Aug. 2–Oct.

Vt.— Rare migrant: Burlington.

MASS.— Rare (? spring and) fall migrant. (? April 29–May  
25); July 15–Nov. 7.

R. I.— Rare (? spring and) fall migrant coastwise. July 22–  
Oct. 13.

CONN.— Rare migrant.

## 151. TOTANUS MELANOLEUCUS (Gmelin) Vieillot.

**Greater yellow-legs; Cucu; Greater tattler or telltale; Long-  
legged tattler; Stone snipe; Winter yellow-legs.**

*Scolopax melanoleuca* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 659.  
“AUCTUMNO IN ARENIS LITTORIS LABRADOR.”

*Totanus vociferus* Wils. Audubon, Birds of Amer., 1842, vol. 5,  
p. 316, pl. 345.

Salt and fresh marshes; nests on the ground.



## LIMICOLAE.

## SCOLOPACIDAE.

ME.—Common migrant and possibly a rare summer resident in extreme north (Woolastook valley). Late April–May 19 (July); late July–Oct. 22.

N. H.—Common migrant, mainly coastwise. May; July 17–Oct. 27.

VT.—Uncommon migrant. May; fall.

MASS.—Common migrant mainly coastwise; a few non-breeding birds rarely summer. (Mar. 27) April 4–June 16 (June 28); July 6–Nov. 12.

R. I.—Common migrant. (? Mar. 10) April 10–May 26; July 20–Nov. 4.

CONN.—Common migrant. May 14–June 1 (June 13, 1898 New London); Aug–Nov. 15.

152. *TOTANUS FLAVIPES* (Gmelin) Vieillot.

**Yellow-legs; Lesser yellow-legs; Summer yellow-legs; Wandering tattler.**

Audubon, *Birds of Amer.*, 1842, vol. 5, p. 313, pl. 344. Egg, Seebohm, 1896, p. 139, pl. 44, fig. 11.

*Scolopax flavipes* Gmel., *Syst. nat.*, 1788, vol. 1, pt. 2, p. 659. "IN NOVEBORACO."

Salt and fresh marshes; nests on the ground.

ME.—Rare spring and common fall migrant. May 16; late July–(Oct. 17).

N. H.—Rare spring and common fall migrant. May; Aug.–Sept. 4.

VT.—Uncommon migrant.

MASS.—Rare spring and common early fall migrant. April 29–June 2 (June 15); July 10–Sept. 29 (Oct. 11 and 30).

R. I.—Rare spring and common fall migrant. April 28; July 3–Oct. 1.

CONN.—Rare spring and common fall migrant. May 7; Aug. 17–Oct.

## LIMICOLAE.

## SCOLOPACIDAE.

## 153. HELODROMAS SOLITARIUS (Wilson) Sharpe.

**Solitary sandpiper or tattler.**

*Tringa solitaria* Wils., Amer. ornith., 1813, vol. 7, p. 53, pl. 58, fig. 3. "KENTUCKY TO NEW YORK."

*Totanus solitarius* (Wils.). Audubon, Birds of Amer., 1842, vol. 5, p. 309, pl. 343. Egg, Capen, 1886, pl. 19, fig. 6.

Borders of woodland streams and ponds, rarely near salt water; nests on the ground or in other birds' deserted nests.

ME.—Common migrant and in northern part a rare summer resident (Woolastock valley). May 7–30; summer–Oct. 2 (Oct. 21).

N. H.—Common migrant, and possibly a rare summer resident. May 9–June; July 17–(Nov. 1).

VT.—Common migrant; reported breeding at Lake Bomoseen, St. Albans Bay, Wells River. May–Sept. 3. Eggs, middle of May.

MASS.—Common migrant, mainly inland. May 2–26 (June); (July 8) July 20–Oct. 14 (Nov. 28).

R. I.—Uncommon migrant. May 3–20; July 18–Sept. 25.

CONN.—Common migrant. May 2–(June 10); Aug.–Sept.

## 154. CATOPTROPHORUS SEMIPALMATUS (Gmelin) Bonaparte.

**Willet; Goose-bird; Humility.**

*Scolopax semipalmata* Gmelin, Syst. nat., 1788, vol. 1, pt. 2, p. 659. "IN NOVEBORACO."

*Totanus semipalmatus* Lath. Audubon, Birds of Amer., 1842, vol. 5, p. 324, pl. 347. Egg, Capen, 1886, pl. 19, figs. 4, 5.

Salt and fresh marshes, and beaches; nests on the ground.

ME., N. H.—Rare migrant coastwise. Oct. 25.

VT.—Rare migrant: Middlebury.

MASS.—Rare migrant coastwise; formerly rare summer resident

## LIMICOLAE.

## SCOLOPACIDAE.

(Muskeget, about 1869, New Bedford). May 2-31 (June 17); July 8-Sept. 11.

R. I.—Rare migrant coastwise, mainly in fall. May; July 2-Sept. 18.

CONN.—Formerly a rare summer resident (Madison, Stratford), now only a rare migrant. May; (summer); Aug. 9-Oct. Eggs, June 5, 1873.

155. CATOPTROPHORUS SEMIPALMATUS INORNATUS (Brewster)  
A. O. U. comm.

**Western willet.**

*Symphemia semipalmata inornata* Brewst., Auk, 1887, vol. 4, p. 145. "LARIMER COUNTY, COLORADO."

*Totanus semipalmatus* Temm. Swainson and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 388, pl. 67.

Salt and fresh marshes, and beaches; nests on the ground.

MASS.—Rare fall migrant: Newburyport, Aug. 5, 1908 (and probably most of those now found).

CONN.—Rare fall migrant: Stony Creek, Aug. 15, 1897, and other fall birds.

156. PAVONCELLA PUGNAX (Linné) Leach.

**Ruff (♂); Reeve (♀).**

*Tringa pugnax* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 148. "IN EUROPA MINUS BOREALI."

*Machetes pugnax* Linné. Dresser, Birds of Europe, 1871-81, vol. 8, p. [87], pls. [557, 558]. Egg, Seebohm, 1896, p. 135, pl. 42, figs. 4, 6.

Shores and marshes; nests on the ground.

ME.—Accidental from Old World: Camden, Sept. 14, 1900; Scarborough, April 10, 1870; Upton, Sept. 8, 1874.

## LIMICOLAE.

## SCOLOPACIDAE.

N. H.— Accidental visitor: Seabrook, Sept. 23, 1907.

MASS.— Accidental visitor: Chatham, Sept. 12, 1880; Nantucket, July, 1901; Newburyport marshes, May 20, 1871.

R. I.— Accidental visitor: Point Judith, Aug. 31, 1903; Sakonnet, July 30, 1900.

157. *BARTRAMIA LONGICAUDA* (Bechstein) Bonaparte.

**Bartramian sandpiper; Field plover; Grass plover; Pasture plover; Upland plover.**

*Tringa longicauda* Bechst., Kurze uebers. vögel Lathams, 1812, vol. 2, p. 453. NORTH AMERICA.

*Tringa bartramia* Wils. Audubon, Birds of Amer., 1842, vol. 5, p. 248, pl. 327. Egg, Capen, 1886, pl. 19, figs. 9, 10.

Grassy prairies and fields; nests on the ground.

ME.— Common migrant, and uncommon local summer resident. April 19—summer; Aug. 4—Sept. 16. Eggs, June 6.

N. H.— Now uncommon migrant and summer resident in southern part. May 4—Sept.

VT.— Uncommon migrant and local summer resident. April 24—Sept. Eggs, May 26.

MASS.— Rare spring and uncommon fall migrant, rare local summer resident; formerly commoner. April 3—May 10; summer; July 16—Sept. 14. Eggs, May 25—June 3 (June 22).

R. I.— Uncommon migrant; formerly more common and probably bred. April; Aug. 2—Sept. 18.

CONN.— Formerly common summer resident; now an uncommon migrant. May 2; fall.

158. *TRYNGITES SUBRUFICOLLIS* (Vieillot) Ridgway.

**Buff-breasted sandpiper; Hill grass-bird.**

*Tringa subruficollis* Vieill., Nouv. dict. d'hist. nat., 1819, vol. 34, p. 465. "PARAGUAY."

## LIMICOLAE

## SCOLOPACIDAE.

*Tringa rufescens* Vieill. Audubon, Birds of Amer., 1842, vol. 5, p. 264, pl. 331. Egg, Seebohm, 1896, p. 151, pl. 43, fig. 11.

Grassy uplands and marshes; nests on the ground.

ME.—Rare fall migrant coastwise. Aug.—Sept. 5.

N. H.—Rare fall migrant coastwise: Portsmouth; Rye Beach, Aug. 25 and 28, 1871.

VT.—Doubtfully recorded.

MASS.—Rare fall migrant coastwise. July 28—Sept. 24.

R. I.—Rare fall migrant coastwise. Sept. 6–15.

CONN.—Rare fall migrant coastwise. Aug.—Sept. 30.

159. *ACTITIS MACULARIA* (Linné) Naumann.

**Spotted sandpiper; Ox-eye; Peet-weet; Sand lark; Teeter peep; Tip-tail; Tip-up.**

*Tringa macularia* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 249. "IN EUROPA & AMERICA SEPTENTRIONALI."

*Totanus macularius* Wils. Audubon, Birds of Amer., 1842, vol. 5, p. 303, pl. 342. Egg, Capen, 1886, pl. 19, figs. 7, 8.

Seacoast and freshwater streams and lakes; nests on the ground.

ME.—Common migrant and summer resident. April 26–Oct. 3 (Nov. 19). Eggs, May 20–July 1.

N. H.—Common migrant and summer resident at lower elevations. May 1–Oct. Eggs, May 31.

VT.—Common migrant and summer resident at lower elevations. Late April–Sept.

MASS.—Common migrant and summer resident. (April 19) April 26–Nov. 6 (Nov. 14). Eggs, May 25–July 4.

R. I.—Common migrant and summer resident. (April 16)–Oct. 3. Eggs, May 27–June 6.

CONN.—Common migrant and summer resident. Late April–Oct. 6. Eggs, May 31–June 1.

## LIMICOLAE.

## SCOLOPACIDAE.

160. *NUMENIUS AMERICANUS* Bechstein.

**Long-billed curlew; Great curlew; Hen curlew; Sickle-bill.**

Bechst., Kurze uebers. vögel Lathams, 1812, vol. 4, pt. 2, p. 432.  
 "NEW YORK & HUDSON BAY."

*Numenius longirostris* Wils. Audubon, Birds of Amer., 1843,  
 vol. 6, p. 35, pl. 355.

Marshes and grassy flats; nests on the ground.

ME.— Very rare migrant. May 2; Aug.

N. H.— Formerly rare, now accidental migrant. Aug. 12–25.

MASS.— Now an accidental migrant. July 20–Oct. 18.

R. I.— Now an accidental migrant; last record Jamestown,  
 Sept. 9, 1897. July 15–Sept. 9.

CONN.— Formerly uncommon, now accidental migrant. Aug. 3.

161. *NUMENIUS HUDSONICUS* Latham.

**Hudsonian curlew; Blue-legs (young); Foolish curlew (young);  
 Jack curlew; Short-billed curlew.**

Lath., Index ornith., 1790, vol. 2, p. 712. "IN SINU HUDSONIS."  
 Audubon, Birds of Amer., 1843, vol. 6, p. 42, pl. 356.

Beaches and marshes; nests on the ground.

ME.— Rare migrant. May 20; July 19–late Sept. (Oct. 12,  
 1898, Dover).

N. H.— Rare migrant. Aug. 6–Sept. 2.

MASS.— Rare spring and uncommon fall migrant. April 10–  
 May 30 (June 8); (June 23) July 6–Oct. 1.

R. I.— Rare spring and uncommon fall migrant. April 27–May;  
 July 2–Sept. 25.

CONN.— Rare migrant, mainly in fall. July 20–Aug. 27.

162. *NUMENIUS BOREALIS* (Forster) Latham.

**Eskimo curlew; Dough-bird; Fute.**

Swainson and Richardson, Fauna Boreali-Amer., 1831, vol. 2,  
 p. 378, pl. 65. Egg, Seebohm, 1896, p. 133, pl. 45, fig. 3.

## LIMICOLAE.

## SCOLOPACIDAE.

*Scolopax borealis* Forst., Phil. trans. roy. soc. London, 1772, vol. 62, pp. 411, 431. "ALBANY FORT," HUDSON BAY.

Marshes, beaches, and grassy uplands; nests on the ground.

ME., N. H.—Formerly common fall migrant, coastwise. Late Aug.—Sept.

MASS.—Formerly a common fall migrant, accidental in spring. (Late May 1873, Cape Cod); July 15—Oct. 2.

R. I.—Formerly a common fall migrant coastwise. Aug. 27—Sept.

CONN.—Formerly a common fall migrant coastwise, now very rare. Aug.—Oct. 13.

## CHARADRIIDAE.

163. *SQUATAROLA SQUATAROLA* (Linné) Cuvier.

**Black-bellied plover; Beetlehead; Blackbreast; Blackheart; Frost-bird; Ox-eye; Palebelly (young); Whistling field plover.**

*Tringa squatarola* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 149. "IN EUROPA."

*Charadrius helveticus* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 199, pl. 315.

Beaches and marshes; nests on the ground.

ME., N. H.—Common migrant chiefly coastwise. May 12—(June 24); Aug.—Oct. 24.

VT.—Rare migrant: Lunenburg; Middlebury.

MASS.—Common migrant coastwise. April 15—June 13 (June 25); July 8—Nov. 16 (Dec.).

R. I.—Common migrant. May 15—22; Aug. 9—Oct. 15.

CONN.—Common migrant coastwise.

## LIMICOLAE.

## CHARADRIIDAE.

## 164. CHARADRIUS DOMINICUS Müller.

**American golden plover; Bullhead; Frost-bird; Greenback;  
Greenhead; Green plover; Muddy breast; Pale belly;  
Pale breast; Three toes; Toad-head.**

Müll., Linné's Vollstand. natursyst., suppl., 1776, p. 116. "ST. DOMINGO."

*Charadrius marmoratus* Wagl. Audubon, Birds of Amer., 1842, vol. 5, p. 203, pl. 316. Egg, Seebohm, 1896, p. 125, pl. 39, fig. 5.

Beaches, marshes, and open hilltops; nests on the ground.

ME.—Formerly common, now rare fall migrant coastwise. Aug. 12–Nov. 15.

N. H.—Rare migrant coastwise. (May); Aug. 26–mid-Oct.

VT.—Rare migrant: Lunenburg.

MASS.—Very rare spring, and formerly common but now rare fall migrant coastwise. (May); Aug. 12–Nov. 2 (Nov. 19; Nov. 25, 1908, Ipswich).

R. I.—Migrant, formerly rare in spring and common in fall coastwise; now very rare. (May); Aug. 14–Oct. 25.

CONN.—Formerly common migrant, now rare; mainly coastwise. Oct. 24.

## 165. OXYECHUS VOCIFERUS (Linné) Reichenbach.

**Killdeer.**

*Charadrius vociferus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 150. "IN AMERICA SEPTENTRIONALI."

*Aegialitis vocifera* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 173; Bird-life, 1898, pl. 11. Egg, Capen, 1886, pl. 18, figs. 8, 9.

Meadows, fields, and marshes; less often on beaches; nests on the ground.

ME.—Rare migrant and accidental visitor. Aug. 17–Oct. 24 (Nov. 28–Dec. 4, 1888).



## LIMICOLAE.

## CHARADRIIDAE.

N. H.— Rare migrant and accidental winter resident (1888-89); formerly rare summer resident.

Vt.— Rare migrant and summer resident (Lake Champlain).

MASS.— Rare migrant and summer resident; accidental winter resident (1888-89). Mar. 1-Dec. 15 (winter). Eggs, May 30.

R. I.— Uncommon migrant and rare summer resident; accidental winter resident in 1888-89. Mar.-Dec. (winter). Eggs, May 10-31.

CONN.— Rare migrant and summer resident, formerly more plentiful. (Feb. 21) late Mar.-May; summer. Eggs, June 6.

166. *AEGIALITIS SEMIPALMATA* (Bonaparte) Bonaparte.

**Semipalmated plover; Little ring-neck; Ring-neck.**

*Charadrius semipalmatus* Bonap., Journ. acad. nat. sci. Phila., 1825, vol. 5, p. 98. Based on Wilson: "NEW JERSEY." Audubon, Birds of Amer., 1842, vol. 5, p. 218, pl. 320.

Sandy beaches and salt-marsh flats; nests on the ground.

ME.— Common migrant; mainly coastwise. May-(June 22); July 29-Sept.

N. H.— Common migrant, mainly coastwise.

Vt.— Rare migrant.

MASS.— Common migrant, mainly coastwise. Late April-June 14 (June 26); July 12-Oct. 28.

R. I.— Common migrant, mainly coastwise. April 19-May 15; July 6-Sept. 25.

CONN.— Common migrant, mainly coastwise. May 17-June; July-Sept.

167. *AEGIALITIS MELODA* (Ord) Bonaparte.

**Piping plover; Clam-bird; Mourning-bird; Pale ring-neck;  
Ring-neck.**

*Charadrius melodus* Ord, Wilson's Amer. ornith., 1824, vol. 7,

## LIMICOLAE.

## CHARADRIIDAE.

p. 71. "GREAT EGG-HARBOR," N. J. Audubon, *Birds of Amer.*, 1842, vol. 5, p. 223, pl. 321. Egg, Capen, 1886, pl. 18, fig. 10.

Sandy beaches; nests on the ground.

ME.— Formerly a rare summer resident coastwise (Wells Beach).  
May 2–Aug.

N. H.— Rare migrant coastwise.

VT.— Doubtfully recorded.

MASS.— Uncommon migrant and summer resident coastwise.  
(Mar. 26) April 5–Sept. 22. Eggs, May 10–June 14 (July 20).

R. I.— Uncommon migrant and summer resident coastwise.  
(Mar. 24) May 15–Sept. Eggs, June 5–14.

CONN.— Uncommon migrant and summer resident coastwise.

168. *OCHTHODROMUS WILSONIUS* (Ord) Reichenbach.

**Wilson's plover.**

*Charadrius wilsonia* Ord, *Wilson's Amer. ornith.*, 1814, vol. 9, p. 77, pl. 73, fig. 5. "CAPE ISLAND, NEWJERSEY." Audubon, *Birds of Amer.*, 1842, vol. 5, p. 214, pl. 319.

Sandy beaches and mudflats; nests on the ground.

VT., N. H.— Doubtfully recorded.

MASS.— Accidental visitor: Gernet Pt., Plymouth, Aug. 22, 1877; Ipswich, May 8, 1904 (*nec* Nahant, Aug. 1839).

R. I.— Doubtfully included.

CONN.— Accidental visitor: Bridgeport, July 28, 1888 ("seen," Averill); Stratford (Linsley).

## APHRIZIDAE.

169. *ARENARIA INTERPRES* (Linné) Vieillot.

**European turnstone; Tangle-picker.**

*Tringa interpres* Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 148.  
"IN EUROPA & AMERICA SEPTENTRIONALI."

## LIMICOLAE.

## APIRIZIDAE.

*Streptilas interpres* (Linné). Dresser, Birds of Europe, 1871-81, vol. 7, p. [555], pl. [532].

Seacoasts; nests on the ground.

MASS.—Accidental visitor: Monomoy Island, Sept. 8, 1892, (Bishop).

170. *ARENARIA INTERPRES MORINELLA* (Linné) A. O. U. comm.

**Ruddy turnstone; Bead-bird; Brant-bird; Calico-back; Calico-bird; Chicken plover; Horsefoot snipe; Rock-bird; Rock plover; Stone plover; Turnstone.**

*Tringa morinella* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 249. "AD MARIS LITTORA AMERICAE SEPTENTRIONALIS; & EUROPÆ."

*Streptilas interpres* (Linné). Audubon, Birds of Amer., 1842, vol. 5, p. 231, pl. 323.

Seacoasts; nests on the ground.

ME.—Common migrant. May 22-27; July 28-Sept.

N. H.—Uncommon migrant.

VT.—Rare migrant: Lake Champlain.

MASS.—Uncommon migrant. May 1-31 (June 24); July 25-Oct. 16.

R. I.—Common migrant. May 14-20; Aug. 4-Oct. 8.

CONN.—Common migrant. May 23; Aug.-Sept.

## HAEMATOPODIDAE.

171. *HAEMATOPUS PALLIATUS* Temminck.

**American oyster-catcher.**

Temm., Man. d'ornith., ed. 2, 1820, vol. 2, p. 532. "L'AMÉRIQUE MÉRIDIONALE." Audubon, Birds of Amer., 1842, vol. 5, p. 236, pl. 324. Egg, Gentry, Nests and eggs birds U. S., 1882, p. 277, pl. 47.

Seacoasts; nests on the ground.

## LIMICOLAE.

## HAEMATOPODIDAE.

ME.— Formerly at Portland (Audubon).

MASS.— Now an accidental visitor. Late April; Aug.

CONN.— Now very rare migrant; formerly more plentiful.

## GALLINAE.

## PHASIANI.

## TETRAONIDAE.

172. *COLINUS VIRGINIANUS* (Linné) Stejneger.

**Bob-white; Quail; Virginia partridge.**

Chapman, Handb. birds east. No. Amer., 1895, p. 178, pl. 1.  
Egg, Bendire, 1892, vol. 1, pl. 1, fig. 1.

*Tetrao virginianus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 161.  
"AMERICA."

Weedy and brushgrown fields and edges of woods; nests on the ground.

ME., N. H.— Formerly common, now rare resident in southern counties; also introduced from the South.

VT.— Rare resident at lower elevations; largely introduced.

MASS.— Uncommon resident; also introduced. Eggs, May–June 30 (Sept.).

R. I.— Common resident; also frequently introduced. Eggs, May 25–June 20 (! Oct. 10).

CONN.— Common resident, also introduced. Eggs, June 14–Aug. (! Sept. 2).

173. *CANACHITES CANADENSIS CANACE* (Linné) Norton.

**Canadian spruce grouse; Black grouse; Fool-hen; Spotted grouse; Spruce partridge; Wood partridge.**

*Tetrao canace* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 275.  
"CANADA."

*Tetrao canadensis* Linné, Audubon, Birds of Amer., 1842, vol. 5, p. 83, pl. 294. Egg, Bendire, 1892, vol. 1, pl. 1, fig. 23.

GALLINAE.

PHASIANI.

## TETRAONIDAE.

Dense fir and spruce woods; nests on the ground.

ME.—Uncommon resident in northern forests. Eggs, May.

N. H.—Uncommon resident in northern part, and above 3000 ft. in White Mts. south to Mt. Passaconaway. Eggs, late May.

VT.—Uncommon resident in northernmost part (Victoria).

MASS.—Accidental visitor: Gloucester, Sept. 1851; Roxbury Nov. about 1865.

174. *BOXASA UMBELLUS* (Linné) Stephens.

**Ruffed grouse; Birch partridge; Partridge; Pheasant.**

*Tetrao umbellus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 275. "IN PENNSYLVANIA." Audubon, Birds of Amer., 1842, vol. 5, p. 73, pl. 293. Egg, Bendire, 1892, vol. 1, pl. 2, fig. 1.

Woods and thickets; nests on the ground.

VT.—Nearly typical birds rarely found in south part; resident.

MASS.—Uncommon resident. Eggs, April 27–June 2.

R. I.—Common in northern and western parts. Eggs, May 8–19.

CONN.—Common resident. Eggs, April 21–May 20.

175. *BOXASA UMBELLUS TOGATA* (Linné) Ridgway.

**Canadian ruffed grouse; Partridge; Pheasant.**

Ridgway, Manual No. Amer. birds, 1887, p. 198. Egg, Bendire, 1892, vol. 1, pl. 2, fig. 2.

*Tetrao togatus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 275. "CANADA."

Woods and thickets; nests on the ground.

ME.—Common resident. Eggs, April 27–May 25.

N. H.—Common resident up to tree limit on mountains.

VT.—Common resident.

## GALLINAE.

## PHASIANI.

## TETRAONIDAE.

MASS.—Resident; nearly typical birds occur in Berkshire Co. (North Adams).

## 176. LAGOPUS LAGOPUS (Linné) Stejneger.

**Willow ptarmigan.**

*Tetrao lagopus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 159. "IN EUROPAE ALPIS."

*Lagopus albus* Gmel. Audubon, Birds of Amer., 1842, vol. 5, p. 114, pl. 299 (summer). Egg, Bendire, 1892, vol. 1, pl. 2, fig. 5-10.

Dense thickets; nests on the ground.

ME.—Accidental visitor: near Dennisville (Audubon); Kenduskeag, April 23, 1892.

MASS.—Accidental visitor (or escape): Manchester, May 10, 1859.

R. I.—Doubtfully recorded.

## 177. TYMPANUCHUS CUPIDO (Linné) Ridgway.

**Heath-hen; Eastern pinnated grouse; Hethen.**

*Tetrao cupido* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 160. "IN VIRGINIA." De Kay, Zool. New York, 1844, vol. 2, p. 205, pl. 77, fig. 175.

*Cupidonia cupido* Brewster, Auk, 1885, vol. 2, p. 82. Egg, Bendire, 1892, vol. 1, pl. 3, fig. 2.

Scrub-oak thickets; nests on the ground.

ME.—Ascribed by Audubon to Maine, but perhaps erroneously.

MASS.—Resident, now confined to Martha's Vineyard, but (till about 1815) occurring coastwise nearly to Boston, and in lower Connecticut valley. Eggs, June 2-July 24.

R. I.—Formerly common, but practically extirpated by 1800.

CONN.—Formerly common; extirpated before 1840 (Westford, about 1832, Nuttall).

## GALLINAE.

## PHASIANI.

## TETRAONIDAE.

## 178. MELEAGRIS GALLOPAVO SILVESTRIS (Vieillot) Allen.

**Wild turkey.**

*Meleagris silvestris* Vieill., Nouv. dict. d'hist. nat., 1817, vol. 9, p. 447. "ILLINOIS JUSQU'À L'ISTHME DE PANAMA."

*Meleagris gallopavo* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 42, pls. 287, 288. Egg, Bendire, 1892, vol. 1, pl. 3, fig. 14. Woods and thickets; nests on the ground.

ME.— Formerly a resident in southern counties (to Mt. Desert?); now extirpated.

N. H.— Formerly a resident in southern part; last record in 1841 or '42.

VT.— Formerly resident in southern part; now extirpated.

MASS.— Formerly common resident; last recorded capture at Mt. Tom, winter of 1850-51.

R. I.— Formerly common resident.

CONN.— Formerly common resident; last killed about 1813 on Totoket Mt., Northford.

## COLUMBAE.

## COLUMBIDAE.

## 179. ECTOPISTES MIGRATORIUS (Linné) Swainson.

**Passenger pigeon; Wild pigeon.**

*Columba migratoria* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 285. "IN AMERICA SEPTENTRIONALI."

*Ectopistes migratoria* (Linné). Audubon, Birds of Amer., 1842, vol. 5, p. 25, pl. 285. Egg, Bendire, 1892, vol. 1, pl. 4, fig. 6.

Woods and fields; nests in trees in colonies.

ME.— Formerly an abundant migrant and local summer resident; last specimens taken near Dexter, Aug. 16, 1896, and at Bar Harbor, early summer 1904. April 7-Sept.

## COLUMBAE.

## COLUMBIDAE.

N. II.— Formerly abundant migrant and local summer resident; last specimen taken at Concord, 1885. April 2–18; summer–Oct. 10.

VT.— Formerly abundant migrant and summer resident. Mar. 20–Oct. 10.

MASS.— Formerly abundant migrant and common local summer resident, rare in winter; last recorded in 1889. Mar. 10–Oct. 21 (winter). Eggs, May 22.

R. I.— Formerly abundant migrant; last killed in 1886.

CONN.— Formerly an abundant migrant, and common summer resident. April 2–fall. Eggs, late May.

180. *ZENAIDURA MACROURA CAROLINENSIS* (Linné) A. O. U. comm.

**Mourning dove; Carolina dove; Long-tailed dove; Turtle dove.**

*Columba carolinensis* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 286. "IN AMERICA."

*Zenaidura macroura* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 188; Bird-life, 1898, pl. 13. Egg, Bendire, 1892, vol. 1, pl. 4, figs. 8, 9.

Open woods and fields; nests in trees, or rarely on the ground.

ME.— Rare migrant and summer resident in southern part; accidental in winter. Mar. 21–Oct. 24 (Nov. 21 and winter).

N. II.— Rare migrant and summer resident in southern part. Late Mar.–Oct. (Dec. 15).

VT.— Rare summer resident: Rutland Co.

MASS.— Uncommon migrant and summer resident; rare winter resident. Mar. 5–Nov. 15 (winter). Eggs, April 19–May 20.

R. I.— Uncommon summer resident in northern and western parts. April 3–Oct. 15. Eggs, May 15–25.

CONN.— Uncommon summer and rare winter resident. Mar.–Nov. (winter). Eggs, May 24–June 20 (July 27).



## COLUMBAE.

## COLUMBIDAE.

## 181. STARNOENAS CYANOCEPHALA (Linné) Bonaparte.

**Blue-headed quail-dove.**

*Columba cyanocephala* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 163. "AMERICA." Audubon, Birds of Amer., 1842, vol. 5, p. 23, pl. 284.

Thickets and woods; nests among vines.

MASS.—One taken at Brookline, Oct. 10, 1870, probably an escape.

## RAPTORES.

## SARCORHAMPHI.

## CATHARTIDAE.

## 182. CATHARTES AURA SEPTENTRIONALIS (Wied) Nelson.

**Turkey vulture; Turkey buzzard.**

*Cathartes septentrionalis* Wied, Reise Nord-America, 1839, vol. 1, p. 162. Locality elsewhere given as NEAR NEW HARMONY, INDIANA.

*Cathartes aura* Linné. Audubon, Birds of Amer., 1840, vol. 1, p. 15, pl. 2. Egg, Bendire, 1892, vol. 1, pl. 4, figs. 1, 3.

Open country; nests on ground in shallow caves or in hollow logs.

ME.—Accidental visitor at all seasons (eight records).

N. H.—Accidental visitor: Hampton Falls, April 1882; North Weare, spring 1887.

VT.—Accidental visitor: Eutaw.

MASS.—Occasional visitor at all seasons.

R. I.—Occasional visitor; four records. May 10–Nov.

CONN.—Occasional (mainly summer) visitor. April 20–Oct. 18.

## 183. CATHARISTA URUBU (Vieillot) Vieillot.

**Black vulture; Black-headed buzzard; Carrion crow; Jim crow.**

*Vultur urubu* Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 1,

## RAPTORES.

## SARCORHAMPHI.

## CATHARTIDAE.

p. 23, pl. 2. "L'AMÉRIQUE, DEPUIS LA NOUVELLE-ÉCOSSE JUSQU'À LA TERRE DE FEU."

*Cathartes atratus* Wils. Audubon, Birds of Amer., 1840, vol. 1, p. 17, pl. 3. Egg, Bendire, 1892, vol. 1, pl. 4, figs. 7, 10.

Open country and vicinity of dwellings; nests on the ground.

ME.—Accidental visitor. Aug. 20–Nov. 3.

VT.—Accidental visitor: Lunenburg; Woodbury, July 10, 1884.

MASS.—Accidental visitor. July–Nov.

CONN.—Accidental visitor: East Lyme, July 6, 1901.

## FALCONES.

## FALCONIDAE.

184. ELANOIDES FORFICATUS (Linné) Coues.

**Swallow-tailed kite.**

Chapman, Handb. birds east. No. Amer., 1895, p. 196; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 1. Egg, Bendire, 1892, vol. 1, pl. 5, figs. 1, 2.

*Falco forficatus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 89. "AMERICA."

Open country; nests in tall trees.

ME.—Erroneously recorded.

N. H.—Doubtfully recorded (Franklin, 1875).

VT.—Doubtfully recorded (Williams).

MASS.—Accidental visitor: Amesbury (= West Newbury) about Sept. 25, 1882; Whately; (?) near Northampton, 1880.

CONN.—Accidental visitor: Lyme, July 2, 1877; Portland, summer 1861; Saybrook, June 16, 1889.

185. CIRCUS HUDSONIUS (Linné) Vieillot.

**Marsh hawk; Blue hawk; Bog-trotter; Frog hawk; Harrier; Mole hawk; Mouse hawk; Mouser; Snake hawk.**

A. K. Fisher, Hawks and owls of U. S., 1893, p. 26, pl. 3. Egg, Bendire, 1892, vol. 1, pl. 5, fig. 8–10.

## RAPTORES.

## FALCONES.

## FALCONIDAE.

*Falco hudsonius* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 128.  
 "AD ERETUM HUDSONIS."

Open country, fields, marshes; nests on the ground.

ME.—Common migrant, less common summer resident. Mar. 19–Oct. 7 (Nov. 12). Eggs, June 4.

N. H.—Common migrant, uncommon summer resident. Mar. 16–Nov. 12.

VT.—Uncommon summer resident. April 14–Oct.

MASS.—Common migrant, uncommon summer resident; rare winter resident on the coast. Mar. 1–Nov. 29 (winter). Eggs, May 12–June 8.

R. I.—Common migrant, rare summer and winter resident. Mar. 15–Oct. 29 (winter). Eggs, May 23, 25.

CONN.—Common migrant, uncommon summer resident; winters rarely. Mar.–Nov. 6 (winter). Eggs, May 9–June 9.

186. ACCIPITER VELOX (Wilson) Vigors.

**Sharp-shinned hawk; Pigeon hawk.**

Chapman, Handb. birds east. No. Amer., 1895, p. 198; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 4. Egg, Bendire, 1892, vol. 1, pl. 5, fig. 11–17.

*Falco velox* Wils., Amer. ornith., 1812, vol. 5, p. 116, pl. 45, fig. 1.  
 "BANKS OF THE SCHUYLKILL."

Open or wooded country; nests in trees.

ME.—Common migrant, less common summer resident; rare winter resident (Portland, 1905; Westbrook, 1907). Mar. 27–Nov. 23 (winter). Eggs, June 1.

N. H.—Common migrant, less common summer resident; rare winter resident in southern part. Eggs, May 23–June 1.

VT.—Uncommon summer and rare winter resident.

MASS.—Common migrant, uncommon summer and winter resident. Eggs, May 3–June 10.

RAPTORES.

FALCONES.

## FALCONIDAE.

R. I.— Uncommon migrant, rare summer and winter resident. Eggs, May 23, 25.

CONN.— Common migrant, less common summer and winter resident. Eggs, May 15–June 26.

## 187. ACCIPITER COOPERII (Bonaparte) Gray.

**Cooper's hawk; Chicken hawk.**

Chapman, Handb. birds east. No. Amer., 1895, p. 199; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 5. Egg, Bendire, 1892, vol. 1, pl. 5, fig. 18–20.

*Falco cooperii* Bonap., Amer. ornith., 1828, vol. 2, p. 1, pl. 10, fig. 1. "NEAR BORDENTOWN, NEW JERSEY."

Open country and woods; nests in trees.

ME.— Uncommon summer resident. April 4–Oct. 24 (? Jan. 16).

N. H.— Uncommon migrant and summer resident at lower elevations. Mar. 26–Oct. Eggs, May 12–30.

VT.— Uncommon migrant and summer resident; (? winter). Eggs, April 22.

MASS.— Uncommon migrant and summer resident; rare in winter. Eggs, April 25–June 11.

R. I.— Common migrant, less common summer resident; probably winters rarely. Eggs, May 12–14.

CONN.— Common migrant, less common summer and winter resident. Eggs, April 25–June 18.

## 188. ASTUR ATRICAPILLUS (Wilson) Bonaparte.

**American goshawk; Blue hawk; Partridge hawk.**

*Falco atricapillus* Wils., Amer. ornith., 1812, vol. 6, p. 80, pl. 52, fig. 3. NEAR "PHILADELPHIA."

*Accipiter atricapillus* (Wils.). Chapman, Handb. birds east.

## RAPTORES.

## FALCONES.

## FALCONIDAE.

No. Amer., 1895, p. 199; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 6. Egg, Bendire, 1892, vol. 1, pl. 6, fig. 1.

Open country and heavy woods; nests in trees.

ME.—Common winter resident, less common summer resident. Oct. 26–May; summer. Eggs, April 25–May.

N. H.—Irregularly common winter resident; rare summer resident south to Alstead.

VT.—Irregular winter visitor; rare summer resident south to Rutland, Mendon. Eggs, April 23–May 1.

MASS.—Winter visitor, irregularly common; casual summer visitor (Hoosac Mt., Aug. 15, 1900). Oct. 4–Feb. 26.

R. I.—Irregular winter visitor, sometimes common. Oct. 27–Feb. 14.

CONN.—Irregular winter visitor, sometimes common; accidental in late summer. (Aug. 24, 1893, West Haven) Oct. 2–winter.

189. *BUTEO BOREALIS* (Gmelin) Vieillot.

**Red-tailed hawk or buzzard; Hen hawk.**

Chapman, Handb. birds east. No. Amer., 1895, p. 200; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 7. Egg, Bendire, 1892, vol. 1, pl. 6, figs. 5, 6.

*Falco borealis* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 266. "IN AMERICA BOREALI, POTISSIMUM CAROLINA."

Open country and forests; nests in large trees.

ME.—Uncommon summer resident, and in southern counties rare winter resident. Eggs, May 6–26.

N. H.—Uncommon resident, wintering from White Mts. south.

VT.—Uncommon resident. Eggs, April 22.

MASS.—Uncommon migrant and winter resident; rare summer resident. Eggs, April 6–27.

R. I.—Uncommon winter, and rare summer resident. Nov. 30–Mar. 25; summer. Eggs, April 5–30.

RAPTORES.

FALCONES.

## FALCONIDAE.

CONN.— Uncommon migrant and winter resident; rare summer resident. Sept.—Mar.; summer. Eggs, Mar. 28—April 24 (May 25).

## 190. BUTEO LINEATUS (Gmelin) Vieillot.

**Red-shouldered hawk; Chicken hawk; Hen hawk; Winter hawk.**

Chapman, Handb. birds east. No. Amer., 1895, p. 201; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 8. Egg, Bendire, 1892, vol. 1, pl. 7, fig. 1-5.

*Falco lineatus* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 268. "IN INSULA LONGA," NEW YORK.

Open country and woods; nests in large trees.

ME.— Uncommon summer resident. (Mar. 4) Mar. 20—Oct. 21. Eggs, late April—May 24.

N. H.— Uncommon summer resident north to southern valleys of White Mts. Mar. 27—Nov.

Vt.— Uncommon summer resident (? winters). Mar. 17—fall. Eggs, April 22.

MASS.— Common migrant, less common summer and winter resident. Eggs, April 3—June 6.

R. I.— Uncommon resident. Eggs, April 5-13.

CONN.— Common migrant and winter resident, less common summer resident. Eggs, April 5—June 1.

## 191. BUTEO SWAINSONI Bonaparte.

**Swainson's hawk.**

Bonap., Geogr. and comparative list birds Europe and No. Amer., 1838, p. 3. Based on Audubon: "NEAR THE COLUMBIA RIVER." Chapman, Handb. birds east. No. Amer., 1895, p. 203; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 9. Egg, Bendire, 1892, vol. 1, pl. 8, fig. 1-6.

## RAPTORES.

## FALCONES.

## FALCONIDAE.

Open prairie country or edges of woods near streams; nests in trees.

ME.— Occasional visitor from West. May 19; Sept. 15, Oct. 8.

VT.— Rare visitor: Lunenburg.

MASS.— Rare visitor (four records). April 20–Nov. 7.

192. *BUTEO PLATYPTERUS* (Vieillot) Faxon.**Broad-winged hawk.**

*Sparrius platypterus* Vieill., *Encycl. méthod. ornith.*, 1823, vol. 3, p. 1273. Based on Wilson: "NEAR THE SCHUYLKILL."

*Buteo latissimus* (Wils.). Chapman, *Handb. birds east. No. Amer.*, 1895, p. 204; A. K. Fisher, *Hawks and owls of U. S.*, 1893, pl. 10. Egg, Bendire, 1892, vol. 1, pl. 7, fig. 10–13.

Woodlands, and along streams; nests in trees.

ME.— Uncommon summer resident. (Mar. 31) April 11–Sept. 17. Eggs, May 2–June 12.

N. H.— Uncommon migrant and summer resident. April 8–Sept. 15. Eggs, May 16.

VT.— Uncommon summer resident. April 14–fall.

MASS.— Uncommon migrant, and in remoter districts, a rare summer resident. April 12–Sept. 30 (Oct. 16). Eggs, May 7–31.

R. I.— Uncommon migrant and rare summer resident. April–Oct. Eggs, May 15–19.

CONN.— Uncommon migrant, and rare summer (and ? winter) resident. Eggs, May 3–18.

193. *ARCHIBUTEO LAGOPUS SANCTI-JOHANNIS* (Gmelin) Coues.**American rough-legged hawk; Black hawk.**

Chapman, *Handb. birds east. No. Amer.*, 1895, p. 205; A. K. Fisher, *Hawks and owls of U. S.* 1893, pl. 11. Egg, Bendire, 1892, vol. 1, pl. 8, fig. 10–12.

## RAPTORES.

## FALCONES.

## FALCONIDAE.

*Falco S. johannis* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 273.  
 "IN FRETO HUDSONIS ET NOVA TERRA."

Open country; nests on cliffs or in trees.

ME.— Uncommon fall and winter visitor. Oct. 28–Mar.

N. H.— Uncommon irregular fall and winter visitor. Oct.–  
 (May 6).

VT.— Rare fall and winter visitor. Oct.

MASS.— Rare fall and winter visitor. Oct.–April 13 (May 8).

R. I.— Uncommon fall and winter visitor. Oct.–April 12.

CONN.— Fall and winter visitor, irregular, but sometimes com-  
 mon. Nov. 20–April.

194. *AQUILA CHRYSÆTOS* (Linné) Sprüngli.

**Golden eagle; Ring-tailed eagle.**

Chapman, Handb. birds east. No. Amer., 1895, p. 206; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 13. Egg, Bendire, 1892, vol. 1, pl. 9, figs. 3, 5.

*Falco chrysaetos* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 88. "IN EUROPA."

Mountainous country; nests on cliffs or in large trees.

ME.— Rare visitor; not now known to breed (young (?), Sandy Bay Mt., 1895).

N. H.— Rare visitor; formerly rare resident in White Mts.

VT.— Rare visitor; formerly a resident.

MASS.— Rare visitor at all seasons, not now known to breed.

R. I.— Rare visitor in fall and winter.

CONN.— Rare visitor, at all seasons.

195. *HALIAEETUS LEUCOCEPHALUS* (Linné) Boie.

**Bald eagle; Gray eagle; Washington eagle (young); White-headed eagle.**

Chapman, Handb. birds east. No. Amer., 1895, p. 207; A. K.



## RAPTORES.

## FALCONES.

## FALCONIDAE.

Fisher, Hawks and owls of U. S., 1893, pl. 14. Egg, Bendire, 1892, vol. 1, pl. 9, fig. 7.

*Falco leucocephalus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 124. "IN AMERICA, EUROPA."

Open wild country, especially near lakes and seacoasts; nests in tall trees or on the ground.

ME.—Uncommon migrant and summer resident; less common winter resident coastwise. Eggs, April 4–21.

N. H.—Uncommon migrant and summer resident; rare in winter in southern part.

VT.—Uncommon migrant and summer resident.

MASS.—Uncommon migrant and, near coast, occasional at all seasons. No recent breeding records (formerly on Mt. Tom).

R. I.—Occasional at all seasons, though not known to breed.

CONN.—Occasional at all seasons, but most frequent as a migrant; formerly bred (Mt. Carmel, Saltonstall Ridge, West Peak).

196. *FALCO ISLANDUS* Brünnich.**White gyrfalcon; White or Speckled hawk.**

Brünn., Ornith. borealis, 1764, p. 2. "ISLANDIA."

*Falco islandicus* Lath. Audubon, Birds of Amer., 1840, vol. 1, p. 81, pl. 19. Egg, Seebohm, 1896, p. 4, pl. 1, fig. 2.

Open country and barrens; nests on sea cliffs.

ME.—Accidental visitor: South Winn, Oct. 8, 1893; (*nec* Cape Elizabeth).

N. H.—Doubtfully recorded (Errol, Nov. 4, 1868, Maynard).

MASS.—Doubtfully recorded (Cambridge, Jan.–Feb., 1833, Nuttall).

197. *FALCO RUSTICOLUS* Linné.**Gray gyrfalcon.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 88. "IN SVEDIA."

## RAPTORES.

## FALCONES.

## FALCONIDAE.

Coues, Key to No. Amer. birds, ed. 5, 1903, vol. 2, p. 665; Brisson, Ornith., 1760, vol. 1, pl. 31.

Open country; nests on cliffs.

ME.—Accidental visitor: Cape Elizabeth, Oct. 13, 1877.

198. *FALCO RUSTICOLUS GYRFALCO* (Linné) Stejneger.

**Gyrfalcon; MacFarlane's gyrfalcon.**

Ridgway, Manual No. Amer. birds, 1887, p. 245-246.

*Falco gyrfalco* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 91. "IN EUROPA." Dresser, Birds of Europe, 1871-81, vol. 6, pl. [367]. Egg, Bendire, 1892, vol. 1, pl. 9, figs. 6, 8, 9.

Open or mountainous country; nests on cliffs.

ME.—Very rare winter visitor: Jacques Id., Mar. 7, 1907; Ktaadn Iron Works, December, 1876; "Maine;" North Deering, Dec. 11, 1906.

VT.—Doubtfully recorded.

MASS.—Very rare winter visitor: Melrose, Jan. 1, 1891; Northampton, Feb., 1880; Stowe, 1881.

R. I.—Very rare winter visitor: Point Judith, Oct. 11, 1883; near Providence, winter, 1864-65.

199. *FALCO RUSTICOLUS OBSOLETUS* (Gmelin) Stejneger.

**Black gyrfalcon; Labrador gyrfalcon.**

Ridgway, Manual No. Amer. birds, 1887, p. 245-246.

*Falco obsoletus* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 268. "IN FRETO HUDSONIS."

*Falco labradora* Audubon, Ornith. biogr., vol. 2, pl. 196. Egg, Bendire, 1892, vol. 1, pl. 10, fig. 1.

Open and mountainous country; nests on cliffs or rarely in trees.

ME.—Rare fall and winter visitor. Sept.—Mar. 22.

N. H.—Rare winter visitor: Milford, January, 1891.

RAPTORES.

FALCONES.

## FALCONIDAE.

MASS.—Rare fall and winter visitor (four records). Oct.—Mar. 11.

R. I.—Rare fall and winter visitor: Conanicut Id., Nov. 22, 1891; Newport, Oct. 28, 1896; Tiverton, Dec. 26, 1896.

200. *FALCO PEREGRINUS ANATUM* (Bonaparte) Blasius.

**Duck hawk; American peregrine falcon; Black hawk (Vt.);  
Bullet hawk; Great-footed hawk; Ledge hawk.**

Chapman, Handb. birds east. No. Amer., 1895, p. 209; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 15. Egg, Bendire, 1892, vol. 1, pl. 10, fig. 5-7.

*Falco anatum* Bonap., Geogr. and comparative list birds Europe and No. Amer., 1838, p. 4. "NORTHERN PARTS" of No. America.

Open and mountainous country; nests on ledges or in cavities of large trees.

ME.—Rare summer resident. Mar. 14–Oct. Eggs, April 14.

N. H., VT.—Rare migrant and local summer resident. Mar. 14–Oct. VT., Eggs, May 2–June 30.

MASS.—Rare migrant, and in western mountains, rare local summer resident; rare in winter. Eggs, April 15–May 23.

R. I.—Rare migrant. May; Oct.

CONN.—Rare migrant, summer and winter resident. Eggs, May 9 (large young, May 25, Talcott Mt.).

201. *FALCO COLUMBARIUS* Linné.

**Pigeon hawk.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 90. "AMERICA." Chapman, Handb. birds east. No. Amer., 1895, p. 210; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 16. Egg, Bendire, 1892, vol. 1, pl. 10, figs. 4, 8.

Woods and open country; nests in low trees.

## RAPTORES.

## FALCONES.

## FALCONIDAE.

ME.— Uncommon migrant, rare in winter and summer; probably breeds in northern counties (fledglings noted, Knight). Mar. 28–May 13; (summer ; Sept.–Oct. 10 (Dec. 1, 1906, Westbrook; Feb. 22, 1907, Scarborough).

N. H.— Uncommon migrant. May; Aug.–Sept.

VT.— Uncommon migrant; doubtfully recorded in summer.

MASS.— Uncommon migrant and rare winter resident. April 10–May 16; Sept. 7–Nov. 24; winter.

R. I.— Uncommon migrant, mainly in fall, and rare winter resident. April 24–May; Sept. 3–Nov. 21 (Jan.).

CONN.— Uncommon migrant and rare winter resident; doubtfully recorded in summer. April; (? Aug.); Sept. 25–Nov.; winter.

202. *FALCO TINNUNCULUS* Linné.**Kestrel.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 90. “IN EUROPAE TURRIBUS.” Dresser, Birds of Europe, 1871–81, vol. 6, p. [113], pl. [384]. Egg, Seebohm, 1896, p. 7, pl. 4, fig. 5.

Open and wooded country; nests in dense woods, on cliffs, or in towers of buildings.

MASS.— Accidental visitor from Old World: Nantasket Beach, Sept. 29, 1889.

203. *FALCO SPARVERIUS* Linné.**American sparrow hawk; Killyhawk; Pigeon hawk.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 90. “AMERICA.” Chapman, Handb. birds east. No. Amer., 1895, p. 211; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 17. Egg, Bendire, 1892, vol. 1, pl. 10, fig. 11–16.

Open country; nests in cavities of trees or rarely of buildings.

ME.— Uncommon migrant and summer resident; rare winter

## RAPTORES.

## FALCONES.

## FALCONIDAE.

resident in southern part (Portland, 1907-08; Yarmouth, Jan. 10, 1904). Mar. 19-Oct. 15 (winter). Eggs, May 30.

N. H.—Uncommon migrant and rare summer resident north to White Mt. valleys. Mar. 15-Oct. Eggs, May 16.

VT.—Uncommon migrant and summer resident. Mar. 21-fall.

MASS.—Uncommon migrant and summer resident; uncommon winter resident near coast. Eggs, April 27-May 26.

R. I.—Uncommon migrant, rare winter and summer resident. Eggs, May 10.

CONN.—Uncommon resident. Eggs, April 26-May 7.

204. *PANDION HALIAËTUS CAROLINENSIS* (Gmelin) Ridgway.**American osprey; Fish hawk.**

Chapman, Handb. birds east. No. Amer., 1895, p. 212; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 18. Egg, Bendire, 1892, vol. 1, pl. 10, fig. 17; pl. 11, fig. 5-9.

*Falco haliaëtus*  $\gamma$ . *carolinensis* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 263. "AMERICA."

Seacoasts and vicinity of large streams and lakes; nests in high or low trees, on rocks or artificial sites.

ME.—Common summer resident. (Mar. 28) April 5-Oct. 15. Eggs, May 8-June 19.

N. H.—Uncommon migrant; formerly bred at Lake Umbagog. April; Aug.-Nov. 1.

VT.—Uncommon migrant.

MASS.—Common migrant, and in Bristol Co. a summer resident. (Mar. 16) April 2-May 29; summer; Aug. 26-Nov. 11. Eggs, May 8-July.

R. I.—Common migrant and summer resident coastwise. Mar. 24-Oct. 21. Eggs, May 2-24.

CONN.—Common migrant and uncommon summer resident coastwise. Mar. 20-Nov. 13 (Dec. 1). Eggs, May 5-July 4.

RAPTORES.

STRIGES.

## ALUCONIDAE.

205. ALUCO PRATINCOLA (Bonaparte) A. O. U. comm.

**American barn owl; Monkey-faced owl.**

*Strix pratincola* Bonap., Geogr. and comparative list birds Europe and No. Amer., 1838, p. 7. "NORTHERN PARTS" of No. America. Chapman, Handb. birds east. No. Amer., 1895, p. 213; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 19. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 1.

Open country, hiding in hollow trees and cavities during the day; nests in hollow trees, burrows, cavities in cliffs or buildings.

ME.—Erroneously recorded from Falmouth.

VT.—Accidental visitor: Danby, 1902; Lyndon, June 4, 1894.

MASS.—Rare visitor. May–Dec.

R. I.—Rare visitor; five records.

CONN.—Rare visitor. June 11–Oct. 28.

## STRIGIDAE.

206. ASIO WILSONIANUS (Lesson) Coues.

**American long-eared owl; Cat owl.**

Chapman, Handb. birds east. No. Amer., 1895, p. 214; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 20. Egg, Bendire, 1892, vol. 1, pl. 1, fig. 2.

*Otus wilsonianus* Less., Traité d'ornith., vol. 1, 1831, p. 110. "LES ÉTATS-UNIS. . . . PENSYLVANIE. . . . NEW-YORK."

Dense woods and swamps; nests in trees, usually in old crow or hawk nests.

ME., N. H., VT.—Uncommon resident. Eggs, May 1.

MASS.—Uncommon fall and winter visitor, rare summer resident. Eggs, April 1–May 14.

R. I.—Rare resident. Eggs, April 10.

CONN.—Uncommon resident. Eggs, early April–April 25.

RAPTORES.

STRIGES.

## STRIGIDAE.

207. ASIO FLAMMEUS (Pontoppidan) A. O. U. comm.

**Short-eared owl; Bog owl; Gray marsh owl; Marsh owl.**

*Strix flammea* Pontopp., Danske atlas, 1763, vol. 1, p. 617, pl. 25. Based on Linné: [Sweden].

*Asio accipitrinus* (Pallas). Chapman, Handb. birds east. No. Amer., 1895, p. 215; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 21. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 3.

Open country and marshes; nests on the ground.

ME.—Uncommon resident, more common migrant near coast.

N. H.—Uncommon migrant. April; Nov.

VT.—Uncommon migrant (? winters).

MASS.—Uncommon migrant mainly near coast, rare in winter; rare summer resident. Mar. 15–April 15; summer; Sept. 24–Nov. 30 (winter).

R. I.—Uncommon migrant and rare summer resident; may winter rarely. April; Oct. 26–Dec. 23. Eggs, April 28.

CONN.—Uncommon migrant; rare summer and winter resident.

208. STRIX VARIA Barton.

**Barred owl; Barn owl (Me.); Black-eyed owl; Hoot owl; Round-headed owl.**

*Strix varius* Barton, Fragm. nat. hist. Penna., 1799, p. 11. "PENNSYLVANIA."

*Syrnium nebulosum* (Forst.). Chapman, Handb. birds east. No. Amer., 1895, p. 215; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 22. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 4.

Heavy woods, preferably near water; nests in hollows of trees or in abandoned hawk or crow nests.

ME.—Rather common resident. Eggs, Mar. 27.

N. H.—Rather common resident up to 3000 feet.

VT.—Rather common resident.

RAPTORES.

STRIGES.

## STRIGIDAE.

MASS.— Uncommon resident, more frequent in fall migration. Eggs, Mar. 27–May 13.

R. I.— Uncommon resident, occasionally more common in fall and winter. Eggs, Mar. 15–April 12.

CONN.— Uncommon resident. Eggs (Feb.), Mar. 1–April 30 (May 10).

209. *SCOTIAPTEX NEBULOSA* (Forster) Preble.

**Great gray owl; Cinereous owl; Spectral owl.**

*Strix nebulosa* Forst., Phil. trans. roy. soc. London, 1772, vol. 62, p. 424. "CIRCA SINUM HUDSONIS."

*Strix cinerea* Gmel. Swainson and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 77, pl. 31. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 5.

Wooded country; nests in trees.

ME.— Rare and irregular winter visitor. Nov. 8–Mar. 2.

N. H.— Rare winter visitor.

VT.— Rare and irregular winter visitor. Nov. 27–Feb. 11.

MASS.— Rare winter visitor. Nov. 10–Mar. 4.

R. I.— Very rare winter visitor: Fox Island, Mar. 25, 1883; near Providence, 1870, Nov. 19, 1906.

CONN.— Rare winter visitor: North Haven, Jan. 18, 1893; Stratford, Jan. 6, 1843.

210. *GLAUX FUNEREA RICHARDSONI* (Bonaparte) A. O. U. comm.

**Richardson's owl; Sparrow owl.**

*Nyctale richardsoni* Bonap., Geogr. and comparative list birds Europe and No. Amer., 1838, p. 7. Based on Audubon: "AT BANGOR, IN MAINE, ON THE PENOBSCOT RIVER."

*Strix tengmalmi* Linné. Swainson and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 94, pl. 32. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 6.



## RAPTORES.

## STRIGES.

## STRIGIDAE.

Wooded country; nests in holes in trees.

ME.— Winter visitor of irregular abundance. (Sept.) Nov. 30–Mar. 16.

N. H.— Rare winter visitor. Dec. 15–Mar. 1.

VT.— Rare winter visitor.

MASS.— Rare winter visitor. Nov. 26–Mar.

R. I.— Rare winter visitor (four records). Dec. 18.

CONN.— Rare winter visitor: East Windsor, 1859.

211. *GLAUX ACADICUS* (Gmelin) A. O. U. comm.

**Saw-whet owl; Acadian owl; Little owl.**

*Strix acadica* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 296. “IN AMERICA SEPTENTRIONALI.”

*Nyctala acadica* (Gmel.). Chapman, Handb. birds east. No. Amer., 1895, p. 217.

*Ulula acadica* (Gmel.). Audubon, Birds of Amer., 1840, vol. 1, p. 123, pl. 33. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 7.

Dense woods; nests in old woodpecker holes or cavities of trees.

ME., N. H.— Uncommon resident. Eggs, April 6.

VT.— Uncommon resident, mainly in northern part.

MASS.— Uncommon resident, breeding locally. Eggs, April 4–May 1 (July 3).

R. I.— Uncommon winter visitor; may breed rarely. Nov.–Feb.

CONN.— Rare resident.

212. *OTUS ASIO* (Linné) Stone.

**Screech owl; Cat owl; Little horned owl; Mottled owl; Red owl.**

*Strix asio* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 92. “AMERICA.”

## RAPTORES.

## STRIGES.

## STRIGIDAE.

*Megascops asio* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 218; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 23. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 8.

Open woods, orchards, and shade trees; nests in cavities of trees.

ME.— Rare resident.

N. H.— Uncommon resident of southern part, becoming rare in southern valleys of White Mts.

VT.— Uncommon resident at lower elevations.

MASS.— Common resident. Eggs, April 15–May 5.

R. I.— Common resident. Eggs, April 7–May 2.

CONN.— Common resident. Eggs, April 7–26.

213. *BUBO VIRGINIANUS* (Gmelin) Bonaparte.

**Great horned owl; Cat owl; Eagle owl; Hoot owl.**

Chapman, Handb. birds east. No. Amer., 1895, p. 219; A. K. Fisher, Hawks and owls of U. S., 1893, pl. 24. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 12.

*Strix virginiana* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 287. "IN OMNI AMERICA, IN ASTRACANIA AD CAMSCHATCAM USQUE, ET POLO ARCTICO PROPIUS."

Heavy woods; nests in cavities of trees, in open nests on the branches, or rarely on the ground.

ME., N. H.— Uncommon resident. Eggs, Mar. 11–23.

VT.— Uncommon resident.

MASS.— Uncommon resident. Eggs, Feb. 17–April 20.

R. I.— Rare resident. Eggs, Feb. 28–Mar. 11.

CONN.— Uncommon resident. Eggs, Feb. 19–Mar. 8 (April 12, second laying).

214. *BUBO VIRGINIANUS SUBARCTICUS* (Hoy) Ridgway.

**Arctic horned owl.**

*Bubo subarcticus* Hay, Proc. acad. nat. sci. Phila., 1852, vol. 6, p. 211. "WISCONSIN."

## RAPTORES.

## STRIGES.

## STRIGIDAE.

*Strix arctica* Swains. Swains. and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 87, pl. 32.

Heavy woods or open country; nests probably in cavities of trees, or on their branches.

ME.—Accidental visitor: "Maine"; Brownfield, 1886. (Knight supposes these to be "pale or faded individuals of the typical Horned Owl.")

MASS.—Accidental visitor: Waltham, Nov. 30, 1867.

215. *BUBO VIRGINIANUS HETEROCNEMIS* (Oberholser) A. O. U. comm.

**Labrador horned owl.**

*Asio magellanicus heterocnemis* Oberh., Proc. U. S. nat. mus., 1904, vol. 27, p. 187 "LANCE AU LOUP, LABRADOR."

Dense forests; nests in trees.

ME.—Accidental visitor: near Portland, Mar. 1870 (recorded as *B. v. saturatus*).

216. *NYCTEA NYCTEA* (Linné) Lichtenstein.

**Snowy owl; Arctic owl; White owl.**

*Strix nyctea* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 93. "IN EUROPA & AMERICA SEPTENTRIONALI."

*Surnia nyctea* (Linné). Audubon, Birds of Amer., 1840, vol. 1, p. 113, pl. 28. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 19.

Open country and barren grounds; nests on the ground.

ME.—Uncommon fall and winter visitor. (Sept.) Oct. 15–Feb.

N. H.—Late fall and winter visitor, sometimes common; accidental in summer (Concord, July). Oct. 10–April (July).

VT.—Irregular fall and winter visitor. Oct. 20.

MASS.—Irregular fall and winter visitor, sometimes common. (Oct. 12) Oct. 28–April 15 (May 20).

## RAPTORES.

## STRIGES.

## STRIGIDAE.

R. I.—Uncommon and irregular winter visitor. Nov. 3–Feb. 15.  
CONN.—Uncommon and irregular winter visitor. Oct. 17.

217. *SURNIA ULULA CAPAROCH* (Müller) Stejneger.**American hawk owl; Day owl.**

*Strix caparoch* Müll., Linné' Vollstand. natursyst., suppl., 1776, p. 69. "EUROPA."

*Surnia funerea* Gmel. Audubon, Birds of Amer., 1840, vol. 1, p. 112, pl. 27. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 18.

Open country or woods; nests in trees or rarely in cliffs.

ME.—Rare fall and winter visitor. Oct. 24–Mar. 10 (April 16, 1906, Van Buren).

N. H.—Rare fall and winter visitor. Oct. 25–(April 20, Connecticut Lakes).

VT.—Rare fall and winter visitor; (? summer in north). Nov. 2–winter.

MASS.—Rare fall and winter visitor. Nov. 16–Mar. 25.

R. I.—Rare winter visitor: West Greenwich, Nov. 16, 1906; (also one doubtful record).

CONN.—Rare winter visitor: New Haven, Nov. 1869.

218. *SPEOTYTO CUNICULARIA HYPOGAEA* (Bonaparte) Coues.**Burrowing owl.**

A. K. Fisher, Hawks and owls of U. S., 1893, p. 195, pl. 25. Egg, Bendire, 1892, vol. 1, pl. 12, fig. 14.

*Strix hypogaea* Bonap., Amer. ornith., 1825, vol. 1, p. 72, pl. 7, fig. 2. "WESTERN AMERICAN TERRITORY."

Open prairie country; nests in old burrows of prairie-dogs and spermophiles.

MASS.—Accidental visitor: Newburyport, May 5, 1875.

## COCCYGES.

## CUCULI.

## CUCULIDAE.

219. *COCCYZUS AMERICANUS* (Linné) Bonaparte.

**Yellow-billed cuckoo; Chow-chow; Egg sucker; Milk sourer;  
Rain crow; Rain dove.**

Audubon, *Birds of Amer.*, 1842, vol. 4, p. 293, pl. 275. Egg, Bendire, 1895, vol. 2, pl. 5, fig. 1.

*Cuculus americanus* Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 111. "CAROLINA."

Thickets, shade trees, and open woods; nests in low trees or bushes.

ME.—Rare summer resident in southern counties. May 18–Aug.

N. H.—Rare summer resident of coast and southern valleys. May–Sept.

VT.—Rare summer resident at lower elevations. May–summer. Eggs, June 4–July 10.

MASS.—Uncommon summer resident, mainly of south and east parts. (May 4) May 9–Sept. 26 (Oct. 6; Nov.). Eggs, May 20–June 30.

R. I.—Summer resident, sometimes common. May 9–Sept. 25 (Oct. 23). Eggs, May 24–30.

CONN.—Common summer resident. May 7–Sept. (Oct. 11). Eggs, June 1–July 24 (Aug. 4, 18, 26, Sept. 23).

220. *COCCYZUS ERYTHROPHthalmus* (Wilson) Bonaparte.

**Black-billed cuckoo; Chow-chow; Egg sucker; Milk sourer;  
Rain crow; Rain dove.**

Audubon, *Birds of Amer.*, 1842, vol. 4, p. 300, pl. 276. Egg, Bendire, 1895, vol. 2, pl. 5, figs. 3, 4.

*Cuculus erythrophthalmus* Wils., *Amer. ornith.*, 1811, vol. 4, p. 16, pl. 28. "GEORGIA."

Thickets, shade trees, and open woods; nests in low trees or bushes.

## COCCYGES.

## CUCULI.

## CUCULIDAE.

ME.—Common summer resident. May 11–Sept. 14. Eggs, June 17.

N. H.—Common summer resident at lower elevations. May 12–Sept. 22 (Oct. 5).

VT.—Common summer resident. Eggs, (Aug. 13).

MASS.—Common summer resident. May 4–Sept. 27 (Oct. 21). Eggs, May 20–June 17.

R. I.—Common summer resident. May 11–Sept. 16. Eggs, May 27–June 1.

CONN.—Common summer resident. May 15–Sept. 28 (Nov. 28, 1888, Stratford). Eggs, June 7–July 24 (Aug. 26).

## ALCYONES.

## ALCEDINIDAE.

221. CERYLE ALCYON (Linné) Bonaparte.

**Belted kingfisher; Kingfisher; Lazy-bird.**

*Alcedo alcyon* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 115. "AMERICA." Audubon, Birds of Amer., 1842, vol. 4, p. 205, pl. 255. Egg, Bendire, 1895, vol. 2, pl. 1, fig. 3.

Vicinity of bodies of water; nests in burrows excavated in banks.

ME.—Common summer resident. April 12–Sept. (Dec. 27). Eggs, June 1.

N. H.—Common summer resident; rare winter resident near open streams. Mar.–Nov. 24 (winter). Eggs, May 20.

VT.—Common summer resident; rare winter resident. Mar. 27–Oct. (winter, Brattleboro).

MASS.—Common summer resident, and rare winter resident. Mar. 15–Dec. 5 (winter). Eggs, May 15–June 6.

R. I.—Common summer and uncommon winter resident. Mar.–Oct. 20; winter. Eggs, May 15–20.

CONN.—Common summer and uncommon winter resident. Mar. 29–Nov. 13; winter. Eggs, May 14.

## PICI.

## PICIDAE.

222. *DRYOBATES VILLOSUS* (Linné) Cabanis.

**Hairy woodpecker; Big guinea woodpecker; Big sapsucker.**

*Picus villosus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 175.  
 "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer.,  
 1842, vol. 4, p. 244, pl. 262. Egg, Capen, 1886, pl. 4, fig. 9.

Forests; nests in holes dug in dead or live trees.

ME.—Common resident. Eggs, May 28.

N. H.—Common resident up to 4500 ft.

VT.—Uncommon resident. Eggs, June 8.

MASS.—Uncommon winter visitor and less common summer resident, especially in western part. Eggs, April 22–May 30.

R. I.—Uncommon fall and winter visitor and rare summer resident (Chepachet, Cranston). Sept. 1.–Mar.; summer. Eggs, May 12–28.

CONN.—Uncommon fall and winter visitor, rare summer resident. Eggs, May 2–9.

223. *DRYOBATES PUBESCENS MEDIANUS* (Swainson) Brewster.

**Downy woodpecker; Little guinea woodpecker; Little sapsucker.**

*Picus (Deudrocopus) medianus* Swains., in Swains. and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 308. "MIDDLE PARTS OF NORTH AMERICA. . . . NEW JERSEY."

*Dryobates pubescens* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 230; Bird-life, 1898, pl. 24. Egg, Capen, 1886, pl. 14, fig. 10.

Open woods, shade and orchard trees; nests in holes excavated in dead trees.

ME.—Common resident. Eggs, May 27.

N. H.—Common resident, mainly below 3000 ft.

VT.—Common resident.

PICI.

## PICIDAE.

MASS.— Common resident, most conspicuous in fall and winter. Eggs, May 20–June 21.

R. I.— Uncommon summer and common winter resident. Eggs, May 12–25.

CONN.— Common resident.

224. *PICOIDES ARCTICUS* (Swainson) Gray.

**Arctic three-toed woodpecker; Black-backed three-toed woodpecker.**

*Picus (Apternus) arcticus* Swains., in Swains. and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 313, pl. 57. “NEAR THE SOURCES OF THE ATHABASCA RIVER, LAT. 57°.” Egg, Capen, 1886, pl. 14, fig. 11.

Evergreen forests; nests in holes excavated in dead trees.

ME.— Uncommon winter visitor, and in northern and western counties, rare resident. Eggs, May 23.

N. H.— Rare resident above 3000 ft. in White Mts., south to Mt. Passaconaway and possibly on Mt. Monadnock (Keene, breeding), elsewhere rare visitor.

VT.— Rare resident in northern part and on higher peaks (nests at Burlington, Lunenburg, Sherburne, Pico Peak, etc.); occasional visitor to lower country in fall and winter. Eggs, May 29, June 1.

MASS.— Occasional fall and winter visitor. (Late Sept.) Oct. 16–April 27 (July, 1888, Boxford; Aug. 11, 1883, Dorchester).

CONN.— Rare winter visitor: East Windsor Hill; Simsbury, 1860.

225. *PICOIDES AMERICANUS* Brehm.

**American three-toed woodpecker; Banded-backed woodpecker; Ladder-backed woodpecker.**

*Picoides americana* Brehm, Handb. d. naturgeseh. aller vögel Deutschl., 1831, p. 195. “AMERIKA.”



## PICI.

## PICIDAE.

*Picus hirsutus* Vieill. Audubon, Birds of Amer., 1842, vol. 4, p. 268, pl. 269. Egg, Capen, 1886, pl. 14, fig. 12.

Evergreen forests; nests in holes excavated in dead trees.

ME.— Rare winter visitor, and in north and west, a rare resident. Eggs, June 2.

N. H.— Rare resident above 3000 ft., south to Sandwich range; elsewhere rare winter visitor. Eggs, June 4.

VT.— Rare winter visitor and local resident (Pico Peak above 2500 ft.).

MASS.— Rare winter visitor. April 28.

226. *SPHYRAPICUS VARIUS* (Linné) Baird.

**Yellow-bellied sapsucker; Red-throated sapsucker; Squealing or Whining woodpecker.**

*Picus varius* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 176. "IN AMERICA SEPTENTRIONALI." DeKay, Zool. New York, 1844, vol. 2, p. 188, pl. 18, fig. 38; Bonaparte, Amer. ornith., 1825, vol. 1, pl. 8, figs. 1, 2. Egg, Capen, 1886, pl. 14, fig. 13.

Woods, orchards, and shade trees; nests in holes excavated in dead trees.

ME.— Common migrant and in northern counties a common summer resident. (Mar. 8 and 31) April 10 — Oct. 26 (Nov. 15). Eggs, May 15–June 8.

N. H.— Common migrant and in northern half a common summer resident up to 3000 ft.; accidental winter resident (Charlestown, 1898–99). April 14–Oct. 20 (winter).

VT.— Common migrant, less common summer resident.

MASS.— Uncommon migrant and accidental winter resident; in Berkshire Co. a rare summer resident. April 4–May 10; summer; Sept. 10–Nov. 1 (winter).

R. I.— Uncommon migrant. (Mar. 23)–April; Sept. 26–Oct. 11.

CONN.— Rare spring and uncommon fall migrant. Mar. 30–May 3; Sept. 28–Oct. (Dec. 21, 1908, New Haven).

PICI.

## PICIDAE.

227. *PHILOEOTOMUS PILEATUS ABIETICOLA* (Bangs) A. O. U. comm.

Northern pileated woodpecker; Black woodpecker; Cock of the woods; Laughing woodpecker; Logcock; Stump-breaker; Woodchuck; Woodcock; Woodhen.

*Ceophloeus pileatus abieticola* Bangs, Auk, 1898, vol. 15, p. 176. "GREENVILLE, MAINE."

*Picus pileatus* Linné. DeKay, Zool. New York, 1844, vol. 2, pl. 18, fig. 35. Egg, Capen, 1886, pl. 14, fig. 6.

Heavy timber; nests in holes excavated in dead trees.

ME.— Uncommon resident in wilder portions. Eggs, May 11–May 28.

N. H.— Rare resident mainly in northern and western parts.

VT.— Rare resident, breeds south to Taftsville, Townshend, Wells River. Eggs, May 22.

MASS.— Rare resident in western counties; elsewhere casual. Eggs, May 17.

CONN.— Rare or now an accidental visitor (Granby, Nov. 1, 1890, 1894–95; Litchfield, June 20, 1908).

228. *MELANERPES ERYTHROCEPHALUS* (Linné) Swainson.**Red-headed woodpecker.**

*Picus erythrocephalus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 113. "AMERICA." Audubon, Birds of Amer., 1842, vol. 4, p. 274, pl. 271. Egg, Capen, 1886, pl. 14, fig. 14.

Open woodlands and shade trees; nests in holes excavated in dead trees.

ME.— Occasional visitor, mainly in spring and fall; said to have bred in York Co.

N. H.— Rare and irregular visitor in southern half. May 25–Sept.

VT.— Irregular visitor at all seasons, sometimes common; resident and bred in western part about 1880 (Orwell, Rutland).

PICI.

## PICIDAE.

MASS.— Rare irregular visitor at all seasons, sometimes breeding (Agawam, Brookline, Newton). Eggs, May–June 17.

R. I.— Irregular visitor, sometimes common in fall; rare summer resident (Providence Co.; Warwick, young in nest, July 28, 1882). Eggs, May 28.

CONN.— Rare summer resident and irregular visitor at all seasons; formerly more common.

229. *CENTURUS CAROLINUS* (Linné) Bonaparte.**Red-bellied woodpecker.**

*Picus carolinus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 113. "IN AMERICA SEPTENTRIONALI." DeKay, Zool. New York, 1844, vol. 2, p. 189, pl. 17, fig. 37.

Wooded bottomlands; nests in holes excavated in dead trees.

MASS.— Accidental visitor (five records). May–Nov.

R. I.— Accidental visitor: near Providence, two specimens.

CONN.— Accidental visitor: Hartford; Stratford, Oct. 16, 1842. Suffield, July 30, 1874.

230. *COLAPTES AURATUS LUTEUS* Bangs.

**Northern flicker; Anteater; Clape; Golden-winged woodpecker; Grasshopper woodpecker (Vt.); Harry wicket; High-hole; Pigeon woodpecker; Wake-up; Yaffle; Yellow jay; Yellow-hammer; Yellow-wing (Mass.); Yucker.**

Bangs, Auk, 1898, vol. 15, p. 177. "WATERTOWN, MASS."

*Colaptes auratus* (Linné). Chapman, Bird-life, 1898, p. 116, pl. 26. Egg, Capen, 1886, pl. 14, fig. 15.

Open woods, orchards, fields, clearings; nests in holes excavated in dead trees, or in cavities.

ME.— Common summer, and near coast, a rare winter resident. Mar. 11–Nov. 13 (winter). Eggs, May 27–June 14.

## PICI.

## PICIDAE.

N. H.— Common migrant and summer resident below 2000 ft.; rare winter resident near coast. Eggs, May 24.

Vt.— Common migrant and summer resident. April 18–fall.

MASS.— Common migrant and summer resident; less common winter resident, mainly near coast. Eggs, May 6–June 20.

R. I.— Common migrant, less common summer and winter resident. Mar. 18–Nov.; winter. Eggs, April 29–May 19.

CONN.— Common migrant and summer resident; less common winter resident. Mar. 28–Nov. 28; winter. Eggs, May 18–June 9.

## MACROCHIRES.

## CAPRIMULGI.

## CAPRIMULGIDAE.

## 231. ANTROSTOMUS CAROLINENSIS (Gmelin) Gould.

**Chuck-will's-widow.**

*Caprimulgus carolinensis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 1028. "VIRGINIA & CAROLINA." Audubon, Birds of Amer., 1840, vol. 1, p. 151, pl. 41. Egg, Bendire, 1895, vol. 2, pl. 1, figs. 8, 9.

Woods and thickets, active at night; nests on the ground.

MASS.— Accidental visitor: Revere (Oct. or Dec.), 1884.

CONN.— Accidental visitor: New Haven, May 19, 1889 (after southerly storm).

## 232. ANTROSTOMUS VOCIFERUS (Wilson) Bonaparte.

**Whip-poor-will.**

*Caprimulgus vociferus* Wils., Amer. ornith., 1812, vol. 5, p. 71, pl. 41, fig. 1–3. "KENTUCKY." Audubon, Birds of Amer., 1840, vol. 1, p. 155, pl. 42. Egg, Bendire, 1895, vol. 2, pl. 1, figs. 10, 11.

Woods and thickets, active at night; nests on the ground.

ME.— Common summer resident. April 28–Sept. 21.

MACROCHIRES.

CAPRIMULGI.

## CAPRIMULGIDAE.

N. II.— Common summer resident north to White Mt. valleys. May 6–Sept. 26.

Vt.— Common summer resident at lower elevations. May 5–Sept. 16. Eggs, May 24–31.

MASS.— Common summer resident. (April 22) April 30–Sept. 25. Eggs, May 18–June 9.

R. I.— Common summer resident. April 25–fall. Eggs, May 25–June 8.

CONN.— Common summer resident. Late April–Sept. (Nov. 1, 1880, Noank). Eggs, May 20–June 5 (July 4).

233. *CHORDEILES VIRGINIANUS* (Gmelin) Swainson.

**Nighthawk; Bull-bat; Mosquito hawk; Pork-and-beans.**

Audubon, Birds of Amer., 1840, vol. 1, p. 159, pl. 43. Egg, Bendire, 1895, vol. 2, pl. 2, fig. 1–3.

*Caprimulgus virginianus* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 1028. "IN AMERICA SEPTENTRIONALI, PRAESERTIM VIRGINIA."

Woods, thickets, or in cities and open country; nests on the ground or on flat gravelled roofs; largely nocturnal.

ME.— Common migrant and summer resident. (April 11, East Hebron; April 22) May 3–Sept. 14. Eggs, June 26–July 4.

N. II.— Common migrant and at lower altitudes a less common summer resident. (April 17) May 9–Sept. 15. Eggs, June 10.

Vt.— Common migrant and summer resident. May–Sept.

MASS.— Common migrant, less common summer resident. (April 23) May 9–Oct. 6. Eggs, June 5–15.

R. I.— Common migrant (mainly inland), and summer resident. (April 7) May–Sept. (Oct.) Eggs, June 5–8.

CONN.— Common migrant and less common summer resident. May 11–Oct. Eggs, June 3–23.

MACROCHIRES.

CYPSELI.

## MICROPODIDAE.

234. *CHAETURA PELAGICA* (Linné) Stephens.**Chimney swift; Chimney-bird; Chimney swallow.**

Chapman, Handb. birds east. No. Amer., 1895, p. 239; Bird-life, 1898, pl. 28. Egg, Bendire, 1895, vol. 2, pl. 1, fig. 25.

*Hirundo pelagica* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 192. "AMERICA."

Lives largely on the wing; rests and nests in hollow trees, chimneys, or barns.

ME.—Common summer resident. April 26–Sept. 11. Eggs, June 15–July 10.

N. H.—Common summer resident. April 27–Sept. 7 (Oct. 1).

VT.—Common summer resident. May 1–Sept.

MASS.—Common summer resident. (April 18) April 25–Sept. 25 (Oct. 12, 1907, Sharon). Eggs, June 10–30.

R. I.—Common summer resident. April 26–Sept. 17. Eggs, June 10–July 1.

CONN.—Common summer resident. (April 14) April 21–Sept. 27. Eggs, June 24.

TROCHILI.

## TROCHILIDAE.

235. *TROCHILUS COLUBRIS* Linné.**Ruby-throated hummingbird.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 120. "IN AMERICA, IMPRIMIS SEPTENTRIONALI." DeKay, Zool. New York, 1844, vol. 21, p. 46, pl. 40, fig. 87. Egg, Bendire, 1895, vol. 2, pl. 1, fig. 27.

Open country, haunting flower growths; nests in trees.

ME.—Common summer resident. May 6–Sept. 26 (Oct. 1).

N. H.—Common migrant, less common summer resident at lower elevations. May 13–Sept. 25. Eggs, June 29–July 15.

VT.—Uncommon summer resident. May 15–Sept.

## MACROCHIRES.

## TROCHILI.

## TROCHILIDAE.

MASS.— Common migrant, less common summer resident. (May 1) May 6–Sept. 23 (Oct. 15). Eggs, May 24–June 15.

R. I.— Common summer resident. (May 3) May 6–Sept. 23. Eggs, May 20–July 4.

CONN.— Common summer resident. May 5–Oct. 5 (Oct. 16). Eggs, May 27–June 3 (Aug. 4).

## PASSERES.

## CLAMATORES.

## TYRANNIDAE.

236. *MUSCIVORA FORFICATA* (Gmelin) Oberholser.

**Scissor-tailed flycatcher.**

*Muscivora forficata* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 931. "IN NOVA HISPANIA." Bonaparte, Amer. ornith., 1825, vol. 1, p. 15, pl. 2, fig. 1. Egg, Bendire, 1895, vol. 2, pl. 1, figs. 12, 13.

Open tree-grown country; nests in trees.

VT.— Accidental visitor: St. Johnsbury, ? 1884.

CONN.— Accidental visitor: Wauregan, about April 27, 1876.

237. *TYRANNUS TYRANNUS* (Linné) Jordan.

**Kingbird.**

Chapman, Handb. birds east. No. Amer., 1895, p. 243; Bird-life, 1898, pl. 30. Egg, Bendire, 1895, vol. 2, pl. 1, figs. 14, 15.

*Lanius tyrannus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 94. "IN AMERICA SEPTENTRIONALIS."

Orchards and tree growth along streams and lakes; nests in trees.

ME.— Common summer resident. May 3–Sept. 12 (Sept. 25). Eggs, June 15.

N. H.— Common summer resident north to the White Mt. valleys. May 3–Sept. 11.

VT.— Common summer resident. May 5–Sept.

MASS.— Common summer resident. (April 27) May 3–Sept. 19 (Sept. 26). Eggs, May 30–July 4 (July 17).

PASSERES.

CLAMATORES.

## TYRANNIDAE.

R. I.— Common summer resident. May 1–Sept. 12 (Sept. 28).  
Eggs, May 31–June 17.

CONN.— Common summer resident. (April 13, 26) May 1–  
Sept. 15.

238. *TYRANNUS DOMINICENSIS* (Gmelin) Richardson.

**Gray kingbird.**

*Lanius tyrannus*  $\beta$ . *dominicensis* Gmel., Syst. nat., 1788, vol. 1,  
pt. 1, p. 302. "INSULA S. DOMINICI ET JAMAICA."

*Muscicapa dominicensis* Brisson. Audubon, Birds of Amer.,  
1840, vol. 1, p. 201, pl. 55. Egg, Bendire, 1895, vol. 2, pl. 2, figs.  
3, 4.

Tree growth along seacoasts and large streams; nests in low trees.

MASS.— Accidental visitor: Lynn, Oct. 23, 1869.

239. *TYRANNUS VERTICALIS* Say.

**Arkansas kingbird.**

Say, Long's Exped. to Rocky Mts., 1823, vol. 2, p. 60. On the  
"ARKANSA" River.

*Muscicapa verticalis* (Say). Bonaparte, Amer. ornith., 1825,  
vol. 1, p. 18, pl. 2, fig. 2. Egg, Bendire, 1895, vol. 2, pl. 1, figs. 16, 17.

Open country, usually near water; nests in trees.

ME.— Accidental visitor: Eliot, Oct., ? 1864.

240. *MYIARCHUS CRINITUS* (Linné) Lichtenstein.

**Crested flycatcher; Great crested flycatcher.**

*Muscicapa crinita* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 325.  
"IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer., 1840,  
vol. 1, p. 209, pl. 57. Egg, Bendire, 1895, vol. 2, pl. 2, figs. 8, 9.

Open woods; nests in hollow limbs.



## PASSERES.

## CLAMATORES.

## TYRANNIDAE.

ME.— Uncommon summer resident, mainly in southern part. May 8–Sept. 20. Eggs, June 11–25.

N. H.— Uncommon summer resident north to White Mt. valleys. May 13–Sept. Eggs, early June–July 7.

VT.— Common summer resident. May 8–Sept.

MASS.— Uncommon summer resident. (May 1) May 9–Sept. 26 (Oct. 15). Eggs, June 13–27.

R. I.— Uncommon summer resident. May 2–Aug. 23. Eggs, June 5, 6.

CONN.— Common summer resident. May 4–Sept. 17. Eggs, May 27–June 13.

241. SAYORNIS PHOEBE (Latham) Stejneger.

**Phoebe; Barn, Bridge, or House pewee; Bridge-bird; Pewit.**

Ridgway, Bull. 50, U. S. nat. mus., 1907, pt. 4, p. 594. Egg, Bendire, 1895, vol. 2, pl. 1, fig. 28.

*Muscicapa phoebe* Lath., Index ornith., 1790, vol. 2, p. 489. "IN AMERICA SEPTENTRIONALI, NOVEBORACO."

*Muscicapa fusca* Bonap. Audubon, Ornith. biogr., 1834, vol. 2, p. 122, pl. 120.

Borders of streams and ponds, or about buildings; nests under bridges or culverts, or on projections of buildings or ledges.

ME.— Common summer resident. Mar. 22–Oct. 19 (Nov. 24, 1884, Brewer). Eggs, May 7–June 24.

N. H.— Common summer resident at lower elevations. Mar. 22–Oct. 17. Eggs, June 7–July.

VT.— Common summer resident. Mar. 22–fall.

MASS.— Common summer resident. (Feb. 20) Mar. 12–Oct. 26 (Nov. 15). Eggs, April 27–May 10; June 13–July 3 (2d brood); Aug. 15 (? 3d brood).

R. I.— Common summer resident. Mar. 8–Oct. 27. Eggs, May 7–12.

PASSERES.

CLAMATORES.

## TYRANNIDAE.

CONN.— Common summer and rare or accidental winter resident (New Haven, Dec. 21, 1906–Mar. 10, 1907; Dec. 25, 1907). (Feb. 25) Mar. 11–Oct. 30 (winter). Eggs, May 12–June 11; July 10.

242. *SAYORNIS SAYA* (Bonaparte) Baird.

**Say's phoebe.**

*Muscicapa saya* Bonap., Amer. ornith., 1825, vol. 1, p. 20, pl. 2, fig. 3. "NEAR THE ARKANSAW RIVER, ABOUT TWENTY MILES FROM THE ROCKY MOUNTAINS." Egg, Bendire, 1895, vol. 2, pl. 1, fig. 29.

Open country; nests under bridges, projections of buildings and ledges.

MASS.— Accidental visitor: North Truro, Sept. 30, 1889.

243. *NUTTALLORNIS BOREALIS* (Swainson) Oberholser.

**Olive-sided flycatcher; Three-deer.**

*Tyrannus borealis* Swains., in Swains. and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 141, pl. 35. "CUMBERLAND HOUSE, LAT. 54.°" Egg, Bendire, 1895, vol. 2, pl. 2, figs. 15, 16.

Mixed and coniferous woods; nests in trees.

ME.— Uncommon summer resident, south to Cape Elizabeth. May 7–Sept. 15. Eggs, June 17–23.

N. H.— Uncommon summer resident. May 18–Sept. 14.

VT.— Uncommon summer resident, mainly of mountainous regions. May 10–fall. Eggs, June 10–16.

MASS.— Uncommon summer resident in western part, rare and local in the east. May 10–Sept. 10. Eggs, June 16–July.

R. I.— Rare migrant. May 14–24; ? fall.

CONN.— Rare migrant; may breed in northwestern part. Spring; Aug. 5–Oct. 18.

PASSERES.

CLAMATORES.

TYRANNIDAE.

244. *MYIOCHANES VIRENS* (Linné) Ridgway.**Wood pewee; Dead-limb bird.**

*Muscicapa virens* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 327. "IN CAROLINA AD RIPAS." Audubon, Birds of Amer., 1840, vol. 1, pl. 64.

*Contopus virens* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 247. Egg, Bendire, 1895, vol. 2, pl. 2, figs. 18, 19. Woods and shade trees; nests in trees.

ME.—Common summer resident. May 12–Sept. 27. Eggs, June 20.

N. H.—Common summer resident at lower elevations. May 17–Sept. 17. Eggs, June 9–16.

Vt.—Common summer resident. May 19–Sept.

MASS.—Common summer resident. (May 10) May 12–Sept. 26 (Oct. 1). Eggs, June 10–July 10.

R. I.—Common summer resident. May 12–Sept. 24. Eggs, June 15–30.

CONN.—Common summer resident. May 10–Sept.

245. *EMPIDONAX FLAVIVENTRIS* (Baird) Baird.**Yellow-bellied flycatcher.**

*Tyrannula flaviventris* Baird, Proc. acad. nat. sci. Phila., 1843, vol. 1, p. 283. "NEAR CARLISLE, PENNSYLVANIA."

*Muscicapa flaviventris* (Baird). Audubon, Birds of Amer., 1844, vol. 7, p. 341, pl. 490. Egg, Bendire, 1895, vol. 2, pl. 2, fig. 23.

Damp evergreen thickets; nests among tree roots.

ME.—Uncommon summer resident in northern part. May 21–June 1; summer–Sept. 8. Eggs, June 14–27.

N. H.—Common summer resident above 3000 ft. from White Mts. north; local south to Dublin. May 18–Aug. (? Sept.). Eggs, June 14.

PASSERES.

CLAMATORES.

## TYRANNIDAE.

VT.—Uncommon local summer resident at lower elevations south to Londonderry. May 16–summer.

MASS.—Rare migrant and possible summer resident on Mt. Greylock (June 15, 1907). (May 10) May 15–June 10 (June 15); Aug. 25–Sept. 10 (Nov. 29, 1876, Reading; Dec. 1, 1876, Newton).

R. I.—Rare migrant. May 21; (? July 27) Aug. 7.

CONN.—Uncommon migrant. May 20–early June; Sept. 17.

246. *EMPIDONAX VIRESCENS* (Vieillot) Brewster.**Green-crested flycatcher; Acadian flycatcher.**

Chapman, Handb. birds east. No. Amer., 1895, p. 249. Egg, Bendire, 1895, vol. 2, pl. 2, figs. 26, 27.

*Platyrhynchos virescens* Vieill., Nouv. dict. d'hist. nat., 1818, vol. 27, p. 22. No locality, based on Wilson.

*Muscicapa acadica* Gmel. Audubon, Birds of Amer., 1840, vol. 1, p. 221, pl. 62.

Deep woods near water; nests, pensile, in low trees.

VT.—Rare summer resident: Bennington (nest with young, July 14, 1904); various doubtful records.

MASS.—Very rare summer resident (Hyde Park, 1888; *nee* Springfield).

CONN.—Rare local summer resident (Greenwich, Stamford, Suffield). Eggs, June 2–22.

247. *EMPIDONAX TRAILLI* ALNORUM Brewster.**Alder flycatcher; Traill's flycatcher.**

Brewst., Auk, 1895, vol. 12, p. 161. "UPTON, MAINE." Bird-lore, 1908, vol. 10, plate in no. 3, figs. 2, 3. Egg, Bendire, 1895, vol. 2, pl. 2, fig. 30.

Alder swamps; nests in low bushes.

ME.—Common local summer resident. (May 15) May 18–Sept. 5. Eggs, June 21.

PASSERES.

CLAMATORES.

## TYRANNIDAE.

N. H.—Local summer resident below 2000 ft.; uncommon south of White Mts. (May 11, 16) May 20–Aug. 31. Eggs, June 22–July 8.

VT.—Uncommon local summer resident. (May 17) May 21–Sept. Eggs, June 13–July 30.

MASS.—Local summer resident, rare in eastern part (Essex and Middlesex Cos.), commoner in western. May 16–June 6; summer–Aug. 24. Eggs, June 15–July 3.

CONN.—Local summer resident (Norfolk). (May 6).

248. EMPIDONAX MINIMUS (Baird) Baird.

**Least flycatcher; Chebec.**

Ridgway, Bull. 50, U. S. nat. mus., 1907, pt. 4, p. 561. Birdlore, 1908, vol. 10, plate in no. 3, fig. 1. Egg, Bendire, 1895, vol. 2, pl. 2, fig. 31.

*Tyrannula minima* Baird, Proc. acad. nat. sci. Phila., 1843, p. 284. "NEAR CARLISLE, PENNSYLVANIA."

Orchards, shade trees, and borders of woods; nests in trees.

ME.—Common summer resident. (April 21) April 27–Aug. 31 (Sept. 5). Eggs, June 11.

N. H.—Common summer resident below 2000 ft. April 29–Sept. Eggs, May–June 10; July 10–late July (2d brood).

VT.—Common summer resident of low country. April 29–Sept.

MASS.—Common summer resident. (April 22) April 28–Sept. 21 (Oct. 2). Eggs, May 20–July 1 (2 broods).

R. I.—Common summer resident. (April 22) April 27–fall. Eggs, May 24–June 2.

CONN.—Common summer resident. April 26–Sept.

PASSERES.

OSCINES.

ALAUDIDAE.

249. *OTOCORIS ALPESTRIS* (Linné) Bonaparte.**Horned lark; Shore lark.**

Ridgway, Bull. 50, U. S. nat. mus., 1907, pt. 4, p. 303; Chapman, Bird-life, 1898, pl. 34. Egg, Seebohm, 1896, p. 259, pl. 58, fig. 15.

*Alauda alpestris* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 166. "IN AMERICA SEPTENTRIONALI."

Seashores and open fields; nests on the ground.

ME., N. H.—Common migrant and winter resident, chiefly coastwise. (Sept. 25, Milo, Me.) Oct. 18–April (May 1–7, Moulton, Me.).

VT.—Common migrant. Feb.–Mar.; (Sept. 15)–Nov.

MASS.—Common migrant, less common winter resident, mainly coastwise. (Sept. 27) Oct. 17–April 19 (May 27).

R. I.—Common migrant and winter resident. (Sept. 25)–April 15.

CONN.—Common migrant and winter resident, mainly coastwise. Nov. 18–Mar. 26.

250. *OTOCORIS ALPESTRIS PRATICOLA* Henshaw.**Prairie horned lark.**

Hensh., Auk, 1884, vol. 1, p. 264. "RICHLAND CO., ILL.," but type locality "DENVER, COLORADO" (see Bull. 50, U. S. nat. mus., 1907, pt. 4, p. 311, foot note). Egg, Bendire, 1895, vol. 2, pl. 5, fig. 25.

*Eremophila cornuta* (Wils.). Baird, Brewer, and Ridgway, Hist. No. Amer. birds, 1874, vol. 2, pl. 32, fig. 2 (head of young).

Open fields; nests on the ground.

ME.—Uncommon migrant and summer resident; casual in winter. Feb. 10–Mar. 30; summer–Nov. 26 (Jan. 24, 1899, Pittsfield). Eggs, May 12–31.

## PASSERES.

## OSCINES.

## ALAUDIDAE.

N. H.— Uncommon migrant, and in White Mt. valleys (? and summit of Mt. Washington, 1905) a rare summer resident. Feb. 28–Aug. Eggs and young, May 21, June 22.

Vt.— Uncommon migrant and rare summer resident (at Burlington, Cornwall, Mt. Mansfield, Rygate, St. Albans, Stowe). Feb.–fall. Eggs, Mar.–April 19.

MASS.— Rare migrant and summer resident, occasional winter resident (Longmeadow). Feb. 19–Mar. 22; summer; Oct. 26–Dec. 15 (winter). Eggs, July 10.

R. I.— Rare migrant: Pawtuxet, Nov. 25, 1889.

CONN.— Uncommon migrant and summer resident (Danbury, 1908, Litchfield, 1905, Washington, 1906); “occasional in winter.” Eggs, Mar. 25.

251. *OTOCORIS ALPESTRIS HOYTI* Bishop.**Hoyt's horned lark.**

Bishop, Auk, 1896, vol. 13, p. 130. “CANDO, TOWNER COUNTY, NORTH DAKOTA.”

Open country and coasts; nests on the ground.

CONN.— Rare migrant and (?) winter resident: Guilford, Mar. 7, 1903; West Haven, Jan. 24, 1905 (Bishop).

## CORVIDAE.

252. *CYANOCITTA CRISTATA* (Linné) Strickland.**Blue jay; Corn-bird.**

*Corvus cristatus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 106. “IN AMERICA SEPTENTRIONAL.”

*Garrulus cristatus* Linné. Audubon, Birds of Amer., 1842, vol. 4, p. 110, pl. 231. Egg, Bendire, 1895, vol. 2, pl. 5, figs. 5, 6.

Woodlands; nests in trees.

ME.— Common migrant and resident. Eggs, May–June 4.

PASSERES.

OSCINES.

## CORVIDAE.

N. H.— Common migrant and summer resident; less common in winter south of the White Mts.

VT.— Common migrant and resident. Eggs, May 11.

MASS.— Common migrant and resident. Eggs, April 28–June 15.

R. I.— Common resident. Eggs, May 2–22.

CONN.— Common resident. Eggs, May 5–June 13.

253. *PERISOREUS CANADENSIS* (Linné) Bonaparte.

**Canada jay; Camp robber; Caribou-bird; Gray jay; Grease-bird; Meat-bird; Meat hawk; Moose-bird; Venison hawk; Whiskey-Jack; Whiskey-John; Wiss-ka-chon** (Indian).

*Corvus canadensis* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 158. "CANADA."

*Garrulus canadensis* (Linné). Audubon, Birds of Amer., 1842, vol. 4, p. 121, pl. 234. Egg, Bendire, 1895, vol. 2, pl. 3, figs. 18, 19.

Fir and spruce forests; nests in trees.

ME.— Uncommon resident in northern part, south to Umbagog; elsewhere casual. Eggs, Mar. 16–28.

N. H.— Uncommon resident above 3000 ft. on Sandwich Range north; elsewhere casual.

VT.— Uncommon resident in northern part and on high peaks (Mt. Mansfield); elsewhere casual (Somerset, Aug. 5, 1896).

MASS.— Accidental visitor: Lexington (*nee* Arlington Heights), Oct. 16, 1889; Newtonville, early summer, 1875; Salem, Oct. 25, 1878.

254. *CORVUS CORAX PRINCIPALIS* Ridgway.**Northern raven.**

Ridgw., Manual No. Amer. birds, 1887, p. 361. "NORTHERN NORTH AMERICA, FROM GREENLAND TO ALASKA, SOUTH TO BRITISH COLUMBIA, CANADA, NEW BRUNSWICK."



PASSERES.

OSCINES.

## CORVIDAE.

*Corvus corax* Linné. Audubon, Birds of Amer., 1843, vol. 4, p. 78, pl. 224. Egg, Bendire, 1895, vol. 2, pl. 4, fig. 3.

Vicinity of seacoasts and large rivers; nests on cliffs or rarely in trees.

ME.—Uncommon resident, mainly coastwise. Eggs, April 14.

N. H.—Accidental visitor (in winter?).

VT.—Very rare visitor in northern part: Brandon, 1879.

MASS.—Accidental visitor, no recent records (Williamstown, before 1877).

CONN.—Accidental visitor: South Manchester, Sept. 18, 1890.

255. *CORVUS BRACHYRYNCHOS* C. L. Brehm.**American crow.**

C. L. Brehm, Beitr. z. vögelkunde, 1822, vol. 2, p. 56. AMERICA.

*Corvus americanus* Audubon, Birds of Amer., 1842, vol. 4, p. 87, pl. 225. Egg, Bendire, 1895, vol. 2, pl. 4, fig. 8-12; pl. 5, figs. 21, 22.

Ubiquitous, though avoiding dense forests in the breeding season; nests in coniferous trees.

ME.—Common resident, in winter chiefly confined to coast. Eggs, April 20–May 25 (June 15).

N. H.—Common migrant and summer resident; in winter less common near coast and large rivers, rare elsewhere. Eggs, April 17.

VT.—Common resident. Eggs, April 22.

MASS.—Common migrant and summer resident; common winter resident near coast, but less so inland. Eggs, April 15–June 13.

R. I.—Common migrant, summer and winter resident. Eggs, April 12–23.

CONN.—Common migrant, summer and winter resident. Eggs, April 12–May 9.

PASSERES.

OSCINES

## CORVIDAE.

256. *CORVUS OSSIFRAGUS* Wilson.**Fish crow.**

Wils., Amer. ornith., 1812, vol. 5, p. 27, pl. 37, fig. 2. "NEW JERSEY . . . NEAR . . . PHILADELPHIA." Egg, Bendire, 1895, vol. 2, pl. 4, figs. 16, 17.

Seaboard and near large streams; nests in trees.

VT.—Doubtfully recorded.

MASS.—Rare visitor in southern part (Cambridge, Springfield, Wareham). Mar. 16–July 16.

CONN.—Rare summer resident (Bridgeport, Fairfield, near New London, New Haven, Stratford). Mar.–Nov. 10. Eggs, May 12–16.

## STURNIDAE.

257. *STURNUS VULGARIS* Linné.**Starling.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 167. "EUROPA, AFRICA." Coues, Key to No. Amer. birds, ed. 5, 1903, vol. 1, p. 503, frontispiece. Egg, Seebohm, 1896, p. 230, pl. 54, fig. 13.

Open country and fields; nests in holes of buildings, trees, or cliffs.

ME.—Accidental visitor: Calais, May 4, 1889.

MASS.—Introduced at Springfield in 1897 but all disappeared; also straggler: Agawam, April, 1908; Pittsfield, Oct. 1, 1899.

CONN.—Local resident; apparently spreading from New York colony (New Haven, Dec. 3, 1900; Stamford, 1900; Norwalk, 1900; Weathersfield and Stonington, 1906; Bethel, 1907; New London, 1907). Eggs, April.

PASSERES.

OSCINES.

## ICTERIDAE.

258. *DOLICHONYX ORYZIVORUS* (Linné) Swainson.

**Bobolink; Butter-bird; May-bird; Meadow-wink; Reed-bird;  
Rice-bird; Skunk blackbird.**

Chapman, Handb. birds east. No. Amer., 1895, p. 261; Auk, 1893, vol. 10, pl. 7. Egg, Bendire, 1895, vol. 2, pl. 6, figs. 1, 2.

*Fringilla oryzivora* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 179. "IN CUBA, . . . FEMINAE SOLAE DESCENDUNT IN CAROLINAM."

Grassy meadows; nests on the ground.

ME.—Common migrant and local summer resident. May 1–Sept. 15. Eggs, June 6–15.

N. H.—Common migrant and summer resident at lower elevations. May 1–Sept. 23.

VT.—Common migrant and summer resident. Eggs, June 1.

MASS.—Common migrant and summer resident. (April 27) May 1–Sept. 29. Eggs, June 1–8.

R. I.—Common migrant and summer resident. May 10–Sept. 25. Eggs, May 30–June 3.

CONN.—Common migrant and summer resident. May 7–Sept. 5. Eggs, June 5.

259. *MOLOTHRUS ATER* (Boddaert) Gray.

**Cowbird; Clodhopper; Cow blackbird; Cow bunting; Lazy-bird; Shiney-eye.**

*Oriolus ater* Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 37. Description in Buffon: "AMÉRIQUE."

*Molothrus pecoris* Gmel. Audubon, Birds of Amer., 1842, vol. 4, p. 16, pl. 212. Egg, Bendire, 1895, vol. 2, pl. 6, figs. 3–4.

Open country; lays eggs in nests of other small birds.

ME.—Common migrant and summer resident. Mar. 19–Oct. 13 (Dec. 27). Eggs, May 26–June 2.

N. H.—Uncommon summer resident in southern part; rare winter resident. Mar. 25–Oct. 2 (winter). Eggs, June 1.

PASSERES.

OSCINES.

## ICTERIDAE.

VT.— Uncommon summer resident; very rare in winter. March 23—(Dec. 30).

MASS.— Common migrant and less common summer resident; occasional in winter. (Mar. 1) Mar. 11—Nov. 21; winter. Eggs, May 15—June 15 (July 1).

R. I.— Common migrant and summer resident; rare in winter. Mar. 14—Nov. 3 (Jan. 19). Eggs, May 16—June 2.

CONN.— Common migrant and summer resident; rare winter resident. Mar.—Nov. 4; winter. Eggs, May 9—June 14.

260. *XANTHOCEPHALUS XANTHOCEPHALUS* (Bonaparte) Jordan.

**Yellow-headed blackbird.**

*Icterus xanthocephalus* Bonap., Journ. acad. nat. sci. Phila., 1826, vol. 5, p. 223. Founded on Say: "ENGINEER CANTONMENT," Iowa.

*Icterus icterocephalus* (Linné). Bonap., Amer. ornith., 1825, vol. 1, p. 27, pl. 3, figs. 1, 2. Egg, Bendire, 1895, vol. 2, pl. 6, fig. 10-12.

Open prairies, marshes, and sloughs; nests among reeds.

ME.— Accidental visitor: Spruce Head, Metnic Id., Aug. 17, 1882.

MASS.— Accidental visitor: Eastham, two, Sept. 10, 1877; Monomoy Id., Sept. 8, 1897; Watertown, Oct. 15, 1869.

CONN.— Accidental visitor: Hartford, late July, 1884; New Haven, June, 1878; Stamford, July, 1888.

261. *AGELAIUS PHOENICEUS* (Linné) Vieillot.

**Red-winged blackbird; Fire-wing; Marsh blackbird; Quonka-ree; Redwing; Swamp blackbird.**

Audubon, Birds of Amer., 1842, vol. 4, p. 31, pl. 216. Egg, Bendire, 1895, vol. 2, pl. 6, fig. 13-15.

*Oriolus phoeniceus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 161. "IN AMERICA SEPTENTRIONALI."

PASSERES.

OSCINES.

## ICTERIDAE.

Open country, swamps, and marshes; nests in marsh grass or in bushes near water.

ME.— Common migrant and local summer resident. Mar. 13–Oct. 13 (Nov. 9). Eggs, June 1–July 1.

N. H.— Common migrant and local summer resident in valleys; accidental in winter (Warren). Mar. 3–Nov. 11 (winter).

VT.— Common migrant and summer resident. Mar. 2–Oct.

MASS.— Common migrant and summer resident, wintering rarely in south and east. (Feb. 26) Mar. 6–Oct. 20 (Nov. 28) (winter). Eggs, May 10–June 18.

R. I.— Common migrant and summer resident; casual in winter. Mar. 7–Oct. 19 (Dec. 15, Feb. 10–13). Eggs, May 12–June 16.

CONN.— Common migrant and summer resident; rare winter resident. Mar. 4–Oct. (winter). Eggs, May 30–June 6.

262. *AGELAIUS PHOENICEUS FORTIS* Ridgway.

**Thick-billed redwing; Northern redwing.**

Ridgway, Proc. Washington acad. sci., 1901, vol. 3, p. 153. "OMAHA, NEBRASKA."

Marshes and open country; nests among reeds or bushes.

CONN.— Rare fall migrant and winter resident: North Haven, Nov. 11, 1903, Dec. 25, 1901, Jan. 10, 1902 (Bishop).

263. *STURNELLA MAGNA* (Linné) Swainson.

**Meadowlark; Marsh quail; Old-field lark.**

Chapman, Handb. birds east. No. Amer., 1895, p. 265; Bird-life, 1898, pl. 39. Egg, Bendire, 1895, vol. 2, pl. 6, figs. 20, 21.

*Alda magna* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 167. "IN AMERICA, AFRICA."

Open fields and marshes; nests on the ground.

ME.— Rare migrant and local summer resident in southern

PASSERES.

OSCINES.

## ICTERIDAE.

counties (Rangeley, April 21, 1897); rare in winter (Calais, Jan., 1899; Cape Elizabeth; Westbrook, Jan. 16, 18, 1904). Mar. 15–Nov. 19 (winter). Eggs, June–(July 22).

N. H.—Uncommon summer resident north to White Mt. valleys; rare winter resident on coast. Mar. 26–Oct. 12 (winter).

Vt.—Uncommon local summer resident; rare winter resident (L. Champlain, Middlebury, Pawlet, St. Albans, West Rutland). April 2–Nov. 15 (winter).

MASS.—Common summer resident, and near the coast an uncommon winter resident. Mar. 2–Nov.; winter. Eggs, May 11–June 28.

R. I.—Common summer resident, and in coastal marshes a common winter resident. Eggs, May 6–30.

CONN.—Common summer resident and less common winter resident coastwise. Eggs, June 6.

264. *ICTERUS SPURIUS* (Linné) Bonaparte.**Orchard oriole.**

Chapman, Handb. birds east. No. Amer., 1895, p. 266; Bird-life, 1898, pl. 36. Egg, Bendire, 1895, vol. 2, pl. 7, fig. 3–5.

*Oriolus spurius* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 162. "IN AMERICA SEPTENTRIONALIS."

Shade trees and orchards; nests pensile in trees.

ME.—Accidental visitor: near Auburn; near Calais; Thomaston.

N. H.—Accidental visitor: Rollinsford.

Vt.—Very rare summer resident: Bennington; Brattleboro (with young); Middlebury, June 1, 1883.

MASS.—Rare local summer resident near coast (to Ipswich) and in western valleys. (May 1) May 8–Aug. 15. Eggs, May 28–June 25.

R. I.—Rare summer resident. May 14–July. Eggs, May 30–June 3.

PASSERES.

OSCINES.

## ICTERIDAE.

CONN.— Common summer resident. May 4–Aug. Eggs, June 11–14 (Aug.).

265. *ICTERUS GALBULA* (Linné) Coues.

**Baltimore oriole; Fire-bird; Golden oriole; Golden robin; Hang-bird; Hang-nest; Pea-bird.**

*Coracias galbula* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 108. "IN AMERICA."

*Icterus baltimore* (Linné). Audubon, Birds of Amer., 1842, vol. 4, p. 37, pl. 217. Egg, Bendire, 1895, vol. 2, pl. 7, fig. 6–9.

Orchards and shade trees; nest pensile in trees.

ME.— Common summer resident mainly in southern counties. May 9–Aug. 30 (Sept. 10).

N. H.— Common summer resident at lower elevations. May 6–Sept. 6 (Sept. 20).

VT.— Common summer resident; accidental in winter. May 10–Sept. (Dec. 4, 1884).

MASS.— Common summer resident; accidental in winter. (April 29) May 1–Sept. 9 (Nov. 15, 1885, E. Templeton; Nov. 18, 1888, Hyde Park). Eggs, May 24–July 4.

R. I.— Common summer resident. May 4–Sept. 1. Eggs, May 24–28.

CONN.— Common summer resident. May 1–early Sept. (Nov. 15, 1885, Portland). Eggs, June 6.

266. *ICTERUS BULLOCKI* (Swainson) Bonaparte.

**Bullock's oriole or troupial.**

Audubon, Birds of Amer., 1842, vol. 4, p. 43, pl. 218. Egg, Bendire, 1895, vol. 2, pl. 7, fig. 10–13.

*Xanthornus bullockii* Swains., Phil. mag., 1827, new ser., vol. 1, p. 436. "TABLE LAND. . . . IN MEXICO."

Open country along streams; nests in trees.

ME.— Accidental visitor: Sorrento, about middle of Nov., 1889.

PASSERES.

OSCINES.

ICTERIDAE.

267. *EUPHAGUS CAROLINUS* (Müller) Richmond.**Rusty blackbird; Rusty grackle.**

*Turdus carolinus* Müll., Linné's Vollstand. natursyst., suppl., 1776, p. 140. "CAROLINA."

*Quiscalus ferrugineus* Linné. Audubon, Birds of Amer., 1842, vol. 4, p. 65, pl. 222. Egg, Bendire, 1895, vol. 2, pl. 7, fig. 14-16.

Swamps, marshes, and wet woods; nests in bushes and low trees.

ME.—Common migrant and in northern part a rare summer resident; accidental winter resident (Calais, 1899). Mar. 17–May; summer–Oct. 29 (winter).

N. H.—Common migrant and in northeast part, a rare summer resident (Megalloway R.). Mar. 8–May; summer; Sept. 9–Oct. 16.

VT.—Common migrant.

MASS.—Common migrant, accidental in winter (Brookline, Feb. 8, 1879; Concord, Jan. 22, 1905; Fresh Pond, Feb. 20, 1887; Jamaica Plain, winter 1907–8). Mar. 7–May 18 (May 30); Sept. 12–Nov. 27 (winter).

R. I.—Common migrant. Mar. 13–(? June 5); Sept.–Oct. 23.

CONN.—Common migrant, and rare winter resident. Mar.–April 19; Sept. 11–Nov. (Jan. 16, 29, Feb. 6, Milford; Dec. 25, 1908, New Haven).

268. *QUISCALUS QUISCULA* (Linné) Jordan.**Purple grackle; Crow blackbird.**

Chapman, Handb. birds east. No. Amer., 1895, p. 269. Egg, Bendire, 1895, vol. 2, pl. 7, fig. 21-23.

*Gracula quiscula* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 109. "IN AMERICA SEPTENTRIONALI."

*Quiscalus versicolor* Vieill. Audubon, Birds of Amer., 1842, vol. 4, p. 58, pl. 221.



PASSERES.

OSCINES.

## ICTERIDAE.

Open country, groves, fields; nests in branches of coniferous trees, rarely in cavities or bushes.

N. H.—Accidental visitor: Tilton, Sept. 13, 1902.

VT.—? Summer resident (Perkins).

MASS.—Occasional typical birds breed in southernmost part and coastwise to Cambridge.

R. I.—Common summer resident in southern part, where most birds are typical; casual in winter. Mar. 6–Sept. 21 (Nov., Dec. 31). Eggs, April 25–June 6.

CONN.—Common summer and accidental winter resident. (Feb. 13) Mar. 1–Nov. (winter, East Haven).

269. *QUISCALUS QUISCUA AENEUS* (Ridgway) Stejneger.

**Bronzed grackle; Crow blackbird.**

*Quiscalus aeneus* Ridgw., Proc. acad. nat. sci. Phila., 1869, p. 134. "MISSISSIPPI REGION OF U. S., EAST TO ALLEGHANY MTS." etc. Egg, Bendire, 1895, vol. 2, pl. 7, figs. 26, 27.

Open country, frequenting clumps of coniferous trees in breeding season; nests in branches of trees, or less often in cavities or bushes.

ME.—Common migrant and local summer resident. Mar. 19–Oct. 26. Eggs, May 27–June 21.

N. H.—Uncommon local summer resident; casual in winter. Mar. 12–Nov. (Dec. 24, 1907, Lancaster; Jan.).

VT.—Uncommon local summer resident. April 11–Oct.

MASS.—Common migrant and local summer resident; rare in winter near coast (also Springfield). (Feb. 23, 1909, Boston) Mar. 10–Nov. 22 (winter). Eggs, May 2–June 1.

R. I.—Common migrant and in the northern part a less common summer resident, intergrading with *Q. quiscula*. Mar. 5–Nov. 1. Eggs, May 1–17.

CONN.—Common migrant. Mar. 5–April; Oct.–Nov.

PASSERES.

OSCINES

## FRINGILLIDAE.

270. *HESPERIPHONA VESPERTINA* (W. Cooper) Bonaparte.**Evening grosbeak; Sugar-bird.**

*Fringilla vespertina* Coop., Ann. lyc. nat. hist. New York, 1825, vol. 1, pt. 2, p. 220. "SAULT STE. MARIE, NEAR LAKE SUPERIOR."

*Coccothraustes vespertinus* (Coop.). Chapman, Handb. birds east. No. Amer., 1895, p. 279; Audubon, Birds of Amer., 1841, vol. 3, p. 217, pl. 207.

Wooded districts; nests in trees.

ME.—Accidental visitor: several records in winter of 1889-90, Jan. 10-Mar. 18; Woodfords, Mar. 6, 1909.

N. H.—Accidental visitor: several records in winter of 1889-90, Jan. 6-May 1.

VT.—Accidental visitor: Burlington, about Feb. 20, 1890; Rutland, Feb. 1909.

MASS.—Accidental visitor, first recorded in the 1890 flight (Jan. 1-April); also Wellfleet, one, Dec. 5, 1903; Beverly, flock of five, Mar. 23, 1904; So. Sudbury, Feb. 14, 1909; Townsend, Feb. 7, 1909; Williamstown, Jan. 26, 1909.

CONN.—Accidental visitor, first recorded in 1890 flight (East Hampton, Mar. 2; Gaylordsville, Mar. 10; Portland, Feb., Mar.); also Litchfield, ? Feb. 1, 1905, and Feb. 11, 1905; Bristol, Feb. 27, 1907.

271. *PINICOLA ENUCLEATOR LEUCURA* (Müller) Richmond.**Pine grosbeak; Canadian pine grosbeak; Mope.**

*Loxia leucura* Müller, Linné's Vollstand. natursyst., suppl., 1776, p. 150. "CANADA."

*Pinicola enucleator canadensis* (Cab.). Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 60.

*Pyrrhula enucleator* Temm. Audubon, Ornith. biogr., 1838, vol. 4, p. 414, pl. 358. Egg, Capen, 1886, pl. 7, fig. 13.

PASSERES.

OSCINES.

## FRINGILLIDAE.

Spruce and fir forests; nests in trees.

ME.— Fall and winter visitor of varying abundance, and in northern forests a rare resident. Oct. 25–April 18; summer. Eggs, May 14–27.

N. H.— Fall and winter visitor of varying abundance, and uncommon summer resident above 3000 ft. in White Mts., north. Oct. 18–April 20; summer.

Vt.— Winter visitor of varying abundance. Nov. 15–Mar. 24.

MASS.— Irregular winter visitor of varying abundance. Oct. 24–April 4 (April 24).

R. I.— Irregular winter visitor, sometimes common.

CONN.— Irregular winter visitor, sometimes common. Nov. 24–Mar. 31.

272. *CARPODACUS PURPUREUS* (Gmelin) Gray.

**Purple finch; Red linnet.**

*Fringilla purpurea* Gmelin, Syst. nat., 1788, vol. 1, pt. 2, p. 923.  
“CAROLINA.”

*Erythrospiza purpurea* (Gmel.). Audubon, Birds of Amer., 1841, vol. 3, p. 170, pl. 196. Egg, Capen, 1886, pl. 7, figs. 14, 15.

Orchards, shade trees, and forests; nests in trees.

ME.— Common migrant and summer resident; rare winter resident (Rangeley Lakes, Jan. 15, 1884). Mar. 21–Oct. (winter). Eggs, June 4.

N. H.— Common summer resident below 3000 ft.; rare in winter. April 15–Dec. 22; winter.

Vt.— Common summer resident.

MASS.— Common migrant, less common summer, and occasional winter resident. Mar. 8–Oct. 21; winter. Eggs, May 25–June 19 (!Sept. 10).

R. I.— Common summer and uncommon winter resident. April 7–Oct.; winter. Eggs, May 19–June 8.

PASSERES.

OSCINES.

## FRINGILLIDAE.

CONN.—Common summer and uncommon winter resident. Eggs, June; July 22 (2d brood).

273. *LOXIA CURVIROSTRA MINOR* (Brehm) Ridgway.

**American crossbill; Red crossbill; Screwbill.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 47.

*Crucirostra minor* Brehm, Naumannia, 1853, p. 193. "DER NORDAMERIKANISCHEN FREISTAATEN."

*Loxia curvirostrata* Linné. Audubon, Ornith. biogr., 1835, vol. 2, p. 559, pl. 197. Egg, Capen, 1886, pl. 8, figs. 1, 2.

Coniferous forests; nests in trees.

ME., N. H.—Resident, nomadic and irregularly abundant. Nests from Jan.—Oct.

VT.—Irregular at all seasons, but commonest in winter; breeds occasionally. Eggs, Feb., Mar.

MASS.—Irregularly common winter visitor, and occasional at all seasons; nests rarely (Framingham, 1884; Highland Light, April, 1889; Tyngsborough, Aug., 1875).

R. I.—Irregularly common winter visitor, and occasional at other seasons; not known to breed.

CONN.—Irregular winter visitor, sometimes common. Oct. 7—spring.

274. *LOXIA LEUCOPTERA* Gmelin.

**White-winged crossbill.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 844. "IN SIXU HUDSONIS & NOVEBORACO." Audubon, Birds of Amer., 1841, vol. 3 p. 190, pl. 201.

Coniferous forests; nests in trees.

ME.—Uncommon resident of northern part; elsewhere an irregular visitor. Nest, June.

PASSERES.

OSCINES.

## FRINGILLIDAE.

N. H.— Uncommon resident above 3000 ft. in White Mts. and north; elsewhere an irregular visitor.

VT.— Irregularly common winter visitor, rare in summer in northern part and on higher peaks. Nested at Lunenburg, 1878.

MASS.— Irregular fall and winter visitor, sometimes common. Oct. 22–May 12 (May 25); (June 4, 1900, Lanesboro; June 13, 1866, Newtonville).

R. I.— Rare and irregular winter visitor. Jan. 14–Feb. 24.

CONN.— Rare and irregular winter visitor. Nov. 6–Mar. 7.

275. *ACANTHIS HORNEMANNI EXILIPES* (Coues) Stejneger.

**Hoary redpoll.**

*Aegiothus exilipes* Coues, Proc. acad. nat. sci. Phila., 1861, p. 385. "FORT SIMPSON."

*Fringilla borealis* Temm. Audubon, Ornith. biogr., 1839, vol. 5, p. 87, pl. 400.

Arctic thickets in summer, open tree growth in winter migrations; nests in low trees.

ME.— Rare winter visitor: Westbrook, Jan. 26, 1896; eastern Maine (Boardman).

N. H.— Rare winter visitor: Hampton.

VT.— Doubtfully reported.

MASS.— Rare winter visitor (some ten instances). Nov. 15–Mar. 20.

276. *ACANTHIS LINARIA* (Linné) Bonaparte and Schlegel.

**Redpoll; Lesser redpoll; Mealy redpoll.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 85. Egg, Seebohm, 1896, p. 246, pl. 57, figs. 6, 7.

*Fringilla linaria* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 182. "IN EUROPA." Audubon, Ornith. biogr., 1838, vol. 4, p. 523, pl. 375.

PASSERES.

OSCINES.

## FRINGILLIDAE.

Arctic thickets in summer, open tree growth in winter migrations; nests in low trees.

ME.— Winter visitor of irregular abundance, and rare summer resident (Calais). Oct. 12–April 27 (May 19, 1875; summer).

N. H.— Winter visitor of irregular abundance. Nov. 1–April.

VT.— Winter visitor of irregular abundance; (nest and three eggs said to have been found in late March, 1878, at Taftsville). Nov. 11–May 1 (May 11).

MASS.— Irregularly common fall and winter visitor. (Oct. 16) Oct. 25–April 25.

R. I.— Irregularly common fall and winter visitor. Oct.–Mar. 25.

CONN.— Irregularly common fall and winter visitor. Fall–Mar. 31.

277. *ACANTHIS LINARIA HOLBOELLII* (C. L. Brehm) Dubois.**Holboell's redpoll.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, pp. 80, 89.

*Linaria holboellii* C. L. Brehm, Handb. d. naturgesch. aller vögel Deutschl., 1831, p. 280. "IN DAS MITTLERE DEUTSCHLAND."

*Acanthis holboellii* Bonaparte and Schlegel, Monogr. Loxiens, 1850, p. 50, pl. 53.

Arctic thickets in summer, open tree growth in winter migrations; nests in low trees.

ME.— Rare winter visitor: Gorham, Feb. 3, 1903; North Bridgton, Nov. 25, 1878.

MASS.— Rare winter visitor: Lexington, Mar. 10, 1890; Swampscott, two, Mar. 26, 1883.

278. *ACANTHIS LINARIA ROSTRATA* (Coues) Stejneger.**Greater redpoll.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, pp. 80, 91.

PASSERES.

OSCINES.

## FRINGILLIDAE.

*Acqiothus rostratus* Coues, Proc. acad. nat. sci. Phila., 1861, p. 378. "JACOBSTHAVN, GREENLAND."

*Linota hornemanni* Holb. Dresser, Birds of Europe, 1871-81, vol. 4, pl. [190], lower fig.

Arctic thickets in summer, open tree growth in winter migrations; nests in low trees.

ME.—Winter visitor of irregular abundance. Dec. 12-Mar. 15.  
N. H.—Unsatisfactorily recorded.

MASS.—Winter visitor near coast, irregular and sometimes common. Dec. 27-Feb. 25.

R. I.—Rare winter visitor: East Providence, Mar. 14, 1896.

CONN.—Rare winter visitor: New Haven, Dec. 17, 1878.

279. *ASTRAGALINUS TRISTIS* (Linné) Cabanis.

**American goldfinch; Lettuce-bird; Thistle-bird; Wild canary;  
Yellow-bird.**

*Fringilla tristis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 181.  
"IN AMERICA SEPTENTRIONALI."

*Spinus tristis* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 286; Bird-life, 1898, pl. 52. Egg, Capen, 1886, pl. 8, fig. 3.

Orchards, shade trees, and edges of woods in summer, weedy fields and open country in winter; nests in trees.

ME.—Common migrant and summer resident; uncommon in winter. Eggs, July 29-Aug. 18 (Sept. 5).

N. H.—Common migrant and summer resident; uncommon in winter from White Mts. south.

VT.—Common migrant and summer resident, less common in winter. Eggs, July 26.

MASS.—Common migrant and summer resident, less common winter resident. Eggs, July 21-Aug. 11.

R. I.—Common summer, and less common winter resident. Eggs, July 1-Aug. 7.

CONN.—Common resident. Eggs, July-Aug.

PASSERES.

OSCINES.

## FRINGILLIDAE.

280. *SPINUS PINUS* (Wilson) Stejneger.**Pine siskin; Gray linnet; Pine finch; Pine linnet.**

*Fringilla pinus* Wils., Amer. ornith., 1810, vol. 2, p. 133, pl. 17, fig. 1. "NEIGHBOURHOOD OF PHILADELPHIA."

*Linaria pinus* (Wils.). Audubon, Birds of Amer., 1841, vol. 3, p. 125, pl. 180. Egg, Capen, 1886, pl. 8, fig. 4.

Coniferous forests in summer, open country and coniferous trees in fall and winter; nests in trees.

ME.— Common resident of northern part; elsewhere of irregular occurrence mainly in fall and winter. Eggs, April.

N. H.— Common resident, breeding mainly above 3000 ft. in White Mts. Eggs, mid-April.

VT.— Common winter visitor, less common in summer, breeding irregularly. Nesting, April–May 15.

MASS.— Irregularly common fall and winter visitor, sometimes lingering till late spring and rarely nesting. Sept. 19–May 30 (June 8, July 15, summer). Eggs, May 9–29.

R. I.— Irregularly common fall and winter visitor. Oct.–April 17 (May 31).

CONN.— Irregularly common fall and winter visitor. Oct.–April (May 20).

281. *SPINUS SPINUS* (Linné) Stejneger.**European siskin.**

*Fringilla spinus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 181. "IN EUROPAE JUNIPERETIS."

*Chrysomitris spinus* (Linné). Dresser, Birds of Europe, 1871–81, vol. 3, p. [541], pl. [169]. Egg, Seebohm, 1896, p. 242, pl. 56, fig. 18.

Coniferous woods; nests in trees.

MASS.— Cambridge, Aug. 11, 13, 17, 1904, an apparently wild bird (Brewster).



PASSERES.

OSCINES.

## FRINGILLIDAE.

282. PLECTROPHENAX NIVALIS (Linné) Stejneger.

**Snowflake; Snow bunting; White snowbird.**

*Emberiza nivalis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 176.  
"IN ALPIBUS LAPPONIAE, SPITZBERGAE AD SINUM HUDSONIS  
ALIBIQUE."

*Plectrophanes nivalis* (Linné). Audubon, Birds of Amer., 1841,  
vol. 3, p. 55, pl. 155. Egg, Seebohm, 1896, p. 247, pl. 57, figs. 9, 10.

Open fields, coasts, and barrens; nests on the ground.

ME.—Common migrant and winter resident (? accidental in  
summer, flock, Mt. Katahdin, early Aug. 1869, *vide* Minot). (Oct.  
10) Oct. 15–April 13.

N. H.—Common migrant and winter resident; accidental in  
summer (Webster, June, a cripple). Oct. 18–April 6.

VT.—Common migrant and winter resident. Nov. 17–spring.

MASS.—Common winter resident, especially coastwise. Oct.  
12–April 3 (May).

R. I.—Common winter resident. Oct. 29–Mar. 29.

CONN.—Irregularly common winter resident. Nov. 14–Mar.

283. CALCARIUS LAPPONICUS (Linné) Stejneger.

**Lapland longspur.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 155.

*Fringilla lapponica* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 180.  
"IN LAPPONIA."

*Plectrophanes lapponicus* (Linné). Dresser, Birds of Europe,  
1871–81, vol. 4, pl. [223]. Egg, Seebohm, 1896, p. 248, pl. 57,  
figs. 11, 12.

Open barrens and plains; nests on the ground.

ME.—Rare migrant. Mar. 10–16; Oct. 26–Dec. 29.

N. H.—Rare migrant in late fall: Hampton Beach, Nov. 30,  
1899.

PASSERES.

OSCINES

## FRINGILLIDAE.

Vt.—Rare or accidental migrant: Brandon, Feb. 21, 1879; ? Townshend.

MASS.—Rare spring and uncommon fall migrant and winter resident mainly coastwise. Feb. 22–May 1; Oct. 9–Dec. 11; winter.

R. I.—Uncommon migrant and rare winter resident. (Jan. 6, Feb. 16).

CONN.—Uncommon migrant; may winter. (Jan. 26, 1883, Portland) Feb. 11–Mar. 16; Nov.

284. *CALCARIUS ORNATUS* (Townsend) Stejneger.

**Chestnut-collared longspur.**

*Plectrophaeus ornata* Towns., Journ. acad. nat. sci. Phila., 1837, vol. 7, pt. 2, p. 189. "PRAIRIES OF THE PLATTE RIVER." Audubon, Birds of Amer., 1841, vol. 3, p. 53, pl. 154.

Open plains; nests on the ground.

ME.—Accidental visitor: Scarborough, Aug. 13, 1886.

MASS.—Accidental visitor: Magnolia, July 28, 1876.

285. *POOECETES GRAMINEUS* (Gmelin) Baird.

**Vesper sparrow; Bay-winged bunting; Grass finch; Gray-bird; Ground sparrow; White-tailed field sparrow.**

*Fringilla graminea* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 922. "IN NOVEBORACO."

*Emberiza graminea* (Gmel.). Audubon, Birds of Amer., 1841, vol. 3, p. 65, pl. 159. Egg, Capen, 1886, pl. 8, fig. 9–12.

Grassy fields and uplands; nests on the ground.

ME.—Common migrant and summer resident. (Mar. 30) April 9–Oct. 28. Eggs, late May–June 24; late July–Aug. (2d brood).

N. H.—Common migrant and summer resident at lower elevations. Mar. 29–Oct. (Nov. 14).

PASSERES.

OSCINES.

## FRINGILLIDAE.

VT.—Common migrant and summer resident. Mar. 31–fall. Eggs, June 6.

MASS.—Common migrant and summer resident. (Mar. 17) Mar. 27–Oct. 24 (Nov. 10; Dec. 25, 1889, Somerville). Eggs, May 10–June 10.

R. I.—Common migrant and summer resident. Mar. 19–Oct. 17. Eggs, May 5–21.

CONN.—Common migrant and summer resident; rare winter resident. Late Mar.–Nov. (winter). Eggs, May 22–July 5.

286. *PASSERCULUS PRINCEPS* Maynard.

**Ipswich sparrow; Gray-bird.**

Mayn., Amer. nat., 1872, vol. 6, p. 637. "IPSWICH," Mass.

*Ammodramus princeps* (Mayn.). Dwight, Mem. Nuttall ornith. club, 1895, no. 2, p. 21, plate.

Sand dunes and sea shores; nests on the ground (Sable Id., N. S.).

ME.—Rare migrant coastwise; may winter rarely. Mar. 20–April 8; Oct. 11–Nov. 17 (Jan. 23, 1886, near Pine Point).

N. H.—Rare migrant coastwise. Mar.–April; late Oct.–early Dec. (*nee* Lake Umbagog.)

MASS.—Common migrant and rare winter resident among coastal dunes; casual inland (Fresh Pond, Cambridge). Oct. 12–April 12.

R. I.—Uncommon migrant and winter resident among coastal dunes. Oct. 20–April 11.

CONN.—Uncommon migrant and winter resident coastwise. Oct.–Mar. 24.

287. *PASSERCULUS SANDWICHENSIS SAVANNA* (Wilson) Ridgway.

**Savanna sparrow.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 192.

*Fringilla savanna* Wils., Amer. ornith., 1811, vol. 3, p. 55, pl. 22, fig. 3. "ATLANTIC COAST, FROM SAVANNAH . . . TO . . . NEW

PASSERES.

OSCINES.

## FRINGILLIDAE.

YORK." Audubon, Ornith. biogr., 1839, vol. 5, p. 516, pl. 109. Egg, Capen, 1886, pl. 8, fig. 5-8.

Grassy fields and marshes; nests on the ground.

ME.—Common summer resident, accidental in winter. (Jan. 24, 1897, Seguin Id.) April 6—early Nov. (Nov. 28). Eggs, May 31—June 18; July (2d brood).

N. H.—Common summer resident at lower elevations, uncommon on barren summits of White Mts. (Mar. 31) April 15—Oct. 27. Eggs, June 16—late July.

VT.—Common summer resident. April 7—fall.

MASS.—Common migrant and summer resident; rare winter resident near coast (Ipswich, Longmeadow, Sandwich). (Mar. 22) April 5—Nov. 12 (winter). Eggs, May 21—June 20.

R. I.—Common migrant, less common summer and rare winter resident. April 4—Oct. 23 (winter). Eggs, May 17—29.

CONN.—Common migrant and summer resident. April—Nov.

288. *COTURNICULUS SAVANNARUM AUSTRALIS* (Maynard) A. O. U. comm.

**Grasshopper sparrow; Yellow-winged sparrow.**

*Ammodromus australis* Mayn., Amer. exch. and mart, 1887, vol. 3, p. 33. "BAHAMAS . . . FLORIDA."

*Ammodramus savannarum passerinus* (Wils.). Chapman, Handb. birds east. No. Amer., 1895, p. 293.

*Emberiza passerina* (Wils.). Audubon, Birds of Amer., 1841, vol. 3, pl. 162. Egg, Capen, 1886, pl. 8, fig. 13-15.

Grassy fields and meadows; nests on the ground.

ME.—Rare summer visitor (? or resident) east to Calais. May 24—June 29.

N. H.—Rare summer resident in lowlands south of Lake Winnepesaukee. May 13—?

VT.—Rare summer resident of southern and western valleys. Eggs, July 17.

PASSERES.

OSCINES.

## FRINGILLIDAE.

MASS.—Uncommon local summer resident near coast and in western valleys. (April 25) May 10–Sept. 1 (Dec. 10, 1892, Arlington).

R. I.—Common local summer resident. April 22–Oct. 23. Eggs, May 30–June 5.

CONN.—Common summer resident in southern part and Connecticut valley. Late April–?

289. *AMMODRAMUS HENSLOWII* (Audubon) Gray.

**Henslow's sparrow.**

*Emberiza henslowii* Aud., Ornith. biogr., 1831, vol. 1, p. 360, pl. 77. "OPPOSITE CINCINNATI, IN THE STATE OF KENTUCKY."

*Ammodramus henslowii* (Aud.). Chapman, Handb. birds east. No. Amer., 1895, p. 294. Egg, Capen, 1886, pl. 9, figs. 1, 2.

Meadows and sedgy weed-grown bogs; nests on the ground.

N. H.—Rare local summer resident north to White Mt. valleys (Wonalancet). April 17–Aug.

VT.—Rare local summer resident in southern part (Pownall).

MASS.—Uncommon local summer resident. May 6–Sept. (Nov. 6, Osterville). Eggs, May 25.

R. I.—Rare migrant and summer resident: Cranston, late April, 1874; Warwick, Oct. 6, 1898; Westerly, June 6 and 8, 1900.

CONN.—Rare local summer resident (Eastford, Killingworth, Woodstock, etc.). May–fall. Eggs, –Aug. 6.

290. *AMMODRAMUS CAUDACTUS* (Gmelin) Swainson.

**Sharp-tailed sparrow.**

Chapman, Handb. birds east. No. Amer., 1895, p. 295; Audubon, Birds of Amer., 1841, vol. 3, p. 108, pl. 174. Egg, Capen, 1886, pl. 9, figs. 3, 4.

*Oriolus caudactus* Gmel., Syst. nat., 1788, vol. 1, pt. 1, p. 394. "IN NOVEBORACO."

PASSERES.

OSCINES.

## FRINGILLIDAE.

Salt marshes; nests on the ground.

ME.—Rare summer resident locally in Cumberland Co. June 5–late Oct. (? Nov.).

N. H.—Rare summer resident of the coastal salt marshes. Spring–Oct. 14.

VT.—Doubtfully recorded (Lunenburg).

MASS.—Uncommon summer resident of the coastal salt marshes (Ipswich); rare winter resident (West Barnstable). May 19–Nov. 8 (winter). Eggs, June 8–July 12.

R. I.—Common summer resident of the coastal salt marshes. May 15–Oct. 2. Eggs, May 24–July 14.

CONN.—Common summer and rare winter resident of the coastal salt marshes. April 14–Nov. (winter). Eggs, May 31–June 9.

291. *AMMODRAMUS NELSONI* (Allen) Norton.

**Nelson's sparrow; Nelson's sharp-tailed finch.**

*Ammodramus caudacutus* var. *nelsoni* Allen, Proc. Boston soc. nat. hist., 1875, vol. 17, p. 293. "CALUMET MARSHES NEAR CHICAGO, ILL."

*Ammodramus caudacutus nelsoni* Allen. Dwight, Auk, 1896, vol. 13, p. 273–274, pl. 4, lower fig.

Marshes; nests on the ground.

ME.—Uncommon migrant. May 22–June 5; Oct. 9–25.

MASS.—Uncommon fall migrant; rare winter resident on south coast. Sept. 25–Oct. 15 (winter).

CONN.—Rare spring and uncommon fall migrant on coastal marshes. May; Oct.

292. *AMMODRAMUS NELSONI SUBVIRGATUS* (Dwight) Norton.

**Acadian sharp-tailed sparrow.**

*Ammodramus caudacutus subvirgatus* Dwight, Auk, 1887, vol. 4

PASSERES.

OSCINES.

## FRINGILLIDAE.

p. 233. "HILLSBOROUGH, ALBERT CO., NEW BRUNSWICK."  
Dwight, Auk, 1896, vol. 13, pl. 4, upper fig.

Brackish or freshwater marshes; nests on the ground.

ME.— Uncommon migrant and local summer resident coastwise from Sagadahoc Co., north. May 20—summer; Sept. 10—Oct. 24 (Nov. 15).

MASS.— Uncommon migrant coastwise (once at Springfield). May 20—June 11; Sept. 3—Nov. 5.

R. I.— Uncommon migrant: Charlestown Beach, four, Oct. 15, 1899; Point Judith, April 27, 1887.

CONN.— Rare spring and uncommon fall migrant. Late May—June 9; Oct.

293. *AMMODRAMUS MARITIMUS* (Wilson) Swainson.

**Seaside sparrow or finch; Meadow chippy.**

Audubon, Birds of Amer., 1841, vol. 3, p. 103, pl. 172. Egg, Capen, 1886, pl. 9, figs. 5, 6.

*Fringilla maritima* Wils., Amer. ornith., 1811, vol. 4, p. 68, pl. 34, fig. 2. "LOW, RUSH-COVERED SEA ISLANDS ALONG OUR ATLANTIC COAST."

Salt marshes; nests on the ground.

ME.— Accidental visitor: Shark Id., Aug. 18, 1884.

N. H.— Erroneously recorded for *A. caudacutus*.

MASS.— Rare local summer (Westport) and winter resident on south coast. April 14—(winter). Eggs, July 17.

R. I.— Uncommon summer resident at Point Judith and Middleown marshes. Eggs, July 14.

CONN.— Common summer and possibly rare winter resident of the salt marshes. Late May—Sept. (Dec. 9, 1889, Bridgeport). Eggs, June 8—19.

PASSERES.

OSCINES.

## FRINGILLIDAE.

294. *CHONDESTES GRAMMACUS* (Say) Bonaparte.**Lark sparrow; Quailhead; Road-bird.**

*Fringilla grammaca* Say, Long's Exped. to Rocky Mts., 1823, vol. 1, p. 139. "BELLE FONTAIN, AS WELL AS AT SEVERAL OTHER POINTS ON THIS [MISSOURI] RIVER." Bonap., Amer. ornith., 1825, vol. 1, p. 47, pl. 5, fig. 3.

Grassy fields; nests on the ground or in low trees and bushes.

MASS.—Rare visitor (seven records). April 29; Aug. 12–Nov. 25.

295. *ZONOTRICHIA LEUCOPHRYS* (Forster) Swainson.**White-crowned sparrow.**

*Emberiza leucophrys* Forst., Phil. trans. roy. soc. London, 1772, vol. 62, p. 426. "IN AMERICA BOREALI AD SINUM HUDSONIS."

*Fringilla leucophrys* Gmel. Audubon, Birds of Amer., 1841, vol. 3, p. 157, pl. 192.

Edges of thickets; nests on the ground or in low bushes.

ME.—Uncommon migrant; casual in summer (Androscoggin Lake, June, 1898; Farmington, late June, 1887; Woolastook Valley, Aug.). (May 2) May 5–25; (June, Aug.); Sept. 22–Oct. 22 (Nov. 8).

N. H.—Uncommon migrant. May 6–25; Sept. 21–Oct. 11.

VT.—Uncommon migrant; said to have bred at Rutland; nest built and one egg laid at East Berkshire. May 14 (July, 1906, Mt. Mansfield); Oct. 4–24.

MASS.—Uncommon migrant. May 4–27 (June 6); Sept. 23–Nov. 5.

R. I.—Uncommon migrant; casual in summer (near Providence, July 9). May; Oct.

CONN.—Uncommon migrant; may winter rarely. (Mar. 20) May 10–15; Oct. 12–Dec. 5.



PASSERES.

OSCINES.

## FRINGILLIDAE.

296. *ZONOTRICHIA ALBICOLLIS* (Gmelin) Swainson.**White-throated sparrow; Mountain lark; Peabody-bird.**

*Fringilla albicollis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 921.  
 "IN PENNSYLVANIA."

*Fringilla pennsylvanica* Latham. Audubon, Birds of Amer.,  
 1841, vol. 3, p. 153, pl. 191. Egg, Capen, 1886, pl. 10, figs. 5, 6.  
 Thickets; nests on the ground, rarely in bushes.

ME.—Common migrant, and from Cumberland Co. north, a  
 common summer resident; accidental in winter (Falmouth, winter,  
 1904; Lewiston, Mar. 30, 1904; Lubec, Mar. 24, 1906; Saco  
 Jan. 20, 1886). April 18–Nov. 9 (Nov. 20) (winter). Eggs, May  
 24–June 13.

N. H.—Common migrant, and a common summer resident to  
 5000 ft. in northern half, local in southern. April 23–May 14;  
 summer; Sept.–Oct.

VT.—Common migrant, and except at lower elevations, a less  
 common summer resident; accidental in winter (Brattleboro, Dec.  
 25, 1904, Mar. 31, 1907). April 29–Oct. 24 (winter). Eggs, June  
 21–July 3.

MASS.—Common migrant and at higher elevations of western  
 part, a less common summer resident, sporadic in eastern (Box-  
 ford, Framingham, Wakefield); rare winter resident near coast.  
 April 15–May 26; summer; Sept. 1–Nov. 24; winter. Eggs, June.

R. I.—Common migrant and rare winter resident. April 28–  
 May 10; Oct. 1–Nov. 1; winter.

CONN.—Common migrant and uncommon winter resident;  
 may breed at higher elevations (Litchfield, June 26, 1906). April  
 15–May 20 (June); Sept. 22–Nov.; winter.

297. *SPIZELLA MONTICOLA* (Gmelin) Baird.**Tree sparrow; Winter chip-bird; Winter chippy.**

Chapman, Handb. birds east. No. Amer., 1895, p. 301; Bird-  
 life, 1898, pl. 49.

PASSERES.

OSCINES.

## FRINGILLIDAE.

*Fringilla monticola* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 912.  
"IN AMERICA SEPTENTRIONALI."

Arctic thickets in summer; fields and marshes in winter migration; nests on the ground.

ME.— Common migrant and in southern counties a winter resident. (Early Aug., Fort Kent; Sept.) Oct. 8–April 28 (May 16).

N. H.— Common migrant, and from White Mts. south, uncommon winter resident. Oct. 8–April 24.

VT.— Common migrant and less common winter resident. Oct.–April.

MASS.— Common migrant and less common winter resident. (Sept. 21) Oct. 16–April 28 (May 7, 1893, Lexington).

R. I.— Common winter resident. Nov. 7–April 15.

CONN.— Common migrant and less common winter resident. Oct. 28–April 12.

298. *SPIZELLA PASSERINA* (Bechstein) A. O. U. comm.

**Chipping sparrow; Chip-bird; Chippy; Hair-bird.**

*Fringilla passerina* Bechstein, in Latham, Allgem. uebers. vögel, 1798, vol. 3, pt. 2, p. 544, pl. 120, fig. 1. "CANADA."

*Spizella socialis* (Wils.). Chapman, Handb. birds east. No. Amer., 1895, p. 302. Egg, Capen, 1886, pl. 10, figs. 1, 2.

*Fringilla socialis* Wils. Audubon, Ornith. biogr., 1839, vol. 5, p. 517, pl. 104.

Orchards, fields, and vicinity of dwellings; nests in trees, hedges, or vines.

ME.— Common migrant and summer resident. (April 2) April 9–Oct. 23 (Nov. 7). Eggs, May 23–June 2; early July (2d brood).

N. H.— Common migrant and summer resident at lower elevations. April 10–Oct. 31. Eggs, May 31–June 16; July 29 (2d brood).

VT.— Common migrant and summer resident. April 4–Oct.

PASSERES.

OSCINES.

## FRINGILLIDAE.

MASS.—Common migrant and summer resident. (Mar. 13, 24) April 5–Nov. 25 (Dec. 31, 1869, Watertown). Eggs, May 12–June 26 (2d brood ?).

R. I.—Common migrant and summer resident. April 4–Oct. 17. Eggs, May 19–June 15.

CONN.—Common migrant and summer resident; rare winter resident. April–Nov. (winter).

299. *SPIZELLA BREWERI* Cassin.

**Brewer's sparrow.**

Cass., Proc. acad. nat. sci. Phila., 1856, vol. 8, p. 40. "WESTERN NORTH AMERICA, CALIFORNIA, NEW MEXICO." Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 326.

*Emberiza pallida* Swains. Audubon, Birds of Amer., 1841, vol. 3, pl. 161 (*vide* Ridgway).

Prairies; nests on the ground.

MASS.—Accidental visitor: Watertown, Dec. 15, 1873.

300. *SPIZELLA PUSILLA* (Wilson) Bonaparte.

**Field sparrow; Ground sparrow; Huckleberry-bird.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 318. Egg, Capen, 1886, pl. 10, figs. 3, 4.

*Fringilla pusilla* Wils., Amer. ornith., 1810, vol. 2, p. 121, pl. 16, fig. 2. "PENNSYLVANIA....NORTH AND SOUTH CAROLINA.... GEORGIA." Audubon, Ornith. biogr., 1834, vol. 2, pl. 139.

Bushy fields; nests on the ground or in low bushes.

ME.—Uncommon summer resident in southern counties. April 7–Oct. 1.

N. H.—Locally common summer resident north to White Mt. valleys. April 3–Oct. Eggs, June 12.

VT.—Locally common summer resident. April 4–fall (Dec. 25, 1908, Clarendon). Eggs, July 10.

## PASSERES.

## OSCINES.

## FRINGILLIDAE.

MASS.—Common summer resident and near the coast a rare winter resident. (Mar. 29) April 1–Nov. 13; (winter). Eggs, May 8–June 12.

R. I.—Common summer and rare winter resident. April–Oct. 23; (winter). Eggs, May 19–June 11.

CONN.—Common summer and rare winter resident. (Late Mar.) April 6–Dec. 2 (winter, Hartford, New Haven). Eggs, June 22.

## 301. JUNCO HYEMALIS (Linné) Selater.

**Slate-colored Junco; Black snowbird; Blue snowbird; Junco; Snowbird; White-bill.**

Chapman, Handb. birds east. No. Amer., 1895, p. 304; Bird-life, 1898, pl. 48. Egg, Capen, 1886, pl. 9, figs. 14, 15.

*Fringilla hyemalis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 183. "AMERICA."

Thickets, weedy and brushgrown fields; nests on the ground, rarely in bushes.

ME., N. H.—Common migrant and mainly in northern part, a common summer resident; rare winter resident in southern part. Eggs, May 7–June 29; July 19 (2d brood).

VT.—Common migrant, less common summer and winter resident.

MASS.—Common migrant, less common winter resident; uncommon summer resident in western highlands (Mts. Holyoke, Nonotuck, Tom, Wachusett, Berkshires), sporadic in east (? Boxford, Medford, Wellfleet). Mar. 8–May 21; summer; Sept. 3–Nov. 30; winter. Eggs, –(Aug. 6).

R. I.—Common migrant and winter resident. Oct. 2–April 4 (May 19).

CONN.—Common migrant and winter resident; rare local summer resident (Bear Mt., Salisbury, 2354 ft.). Oct. 8–May 3 (summer).

PASSERES.

OSCINES.

## FRINGILLIDAE.

302. JUNCO HYEMALIS MONTANUS (Ridgway) A. O. U. comm.

**Montana Junco.**

*Junco montanus* Ridgw., Auk, 1898, vol. 15, p. 321. "COLUMBIA FALLS, MONTANA." Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 289.

Edges of thickets; nests on the ground.

MASS.—Accidental visitor: Watertown, Mar. 25, 1874 (this specimen has been recorded as *J. h. shufeldti* and *J. h. oregonus*).

303. MELOSPIZA MELODIA (Wilson) Baird.

**Song sparrow; Bush sparrow; Ground-bird; Ground sparrow.**

*Fringilla melodia* Wils., Amer. ornith., 1810, vol. 2, p. 125, pl. 16, fig. 4. "UNITED STATES." Audubon, Birds of Amer., 1841, vol. 3, pl. 189.

*Melospiza cinerea melodia* (Wils.). Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 354. Egg, Capen, 1886, pl. 9, fig. 10-13.

Swamps, marshes, weedy fields; nests on the ground, in bushes, or even hollow trees.

ME.—Common migrant and summer resident; rare winter resident (Calais, Cape Elizabeth, Gardiner). Mar. 7–Nov. (winter). Eggs, (April) May 20–June 7; July (2d brood).

N. H.—Common migrant and summer resident at lower elevations; rare winter resident near coast. Mar. 9–Dec. (winter). Eggs, May 24–June 22; July 27 (2d brood).

VT.—Common migrant and summer resident; rare winter resident in southern part (Bethel, Brattleboro, Brownsville, Taftsville). Mar. 7–fall (winter).

MASS.—Common migrant and summer resident; less common winter resident at lower elevations. Mar. 10–Nov. 15; winter. Eggs, April 30–July 29 (2d brood).

R. I.—Common migrant and summer resident; less common winter resident. Mar. 1–Nov. 14; winter. Eggs, May 7–10.

CONN.—Common migrant and summer resident; less common winter resident. Eggs, May 24–(Aug. 20).

PASSERES.

OSCINES.

## FRINGILLIDAE.

304. MELOSPIZA LINCOLNII (Audubon) Baird.

**Lincoln's sparrow; Lincoln's finch.**

*Fringilla lincolni* Aud., Ornith. biogr., 1834, vol. 2, p. 535, pl. 193. "LABRADOR."

*Peucaea lincolni* Aud., Birds of Amer., 1841, vol. 3, p. 116, pl. 177.

Edges of dense thickets and tangled growth; nests on the ground.

ME.—Rare spring and uncommon fall migrant; rare summer resident in extreme north (Ft. Kent; ♂ said to have been taken at Boothbay, July 14, and four birds at Bobson's Id., Aug. 9). May 12–June 1; (summer); Sept. 8–Oct. 13.

N. H.—Rare spring and uncommon fall migrant (once reported in summer, Nubanusit Lake, July 28, 1900, Thayer). May 13–26; (? summer); Sept. 7–Oct. 4.

VT.—Rare migrant.

MASS.—Rare spring and uncommon fall migrant. May 7–June 1; Sept. 12–Oct. 14 (Nov. 1).

R. I.—Rare migrant, mainly in fall. (Spring); Sept. 27.

CONN.—Rare spring and uncommon fall migrant. May 23; Sept. 21–Oct. 3.

305. MELOSPIZA GEORGIANA (Latham) Ridgway.

**Swamp sparrow.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 382.

*Fringilla georgiana* Lath., Index ornith., 1790, vol. 1, p. 460. "CUM PRAECEDENTE," = "IN GEORGIAE AMERICANAE INTERIORIBUS."

*Fringilla palustris* Wils. Audubon, Ornith. biogr., 1839, vol. 5, pl. 64. Egg, Capen, 1886, pl. 9, fig. 7–9.

Swamps, marshes, and old fields; nests on the ground.

ME.—Uncommon migrant and local summer resident. (Mar. 20, Bangor) April 6–Nov. 15. Eggs, May 31–June 3.

## PASSERES.

## OSCINES.

## FRINGILLIDAE.

N. H., Vt.—Common migrant and less common local summer resident at lower elevations. April–Oct. 25.

MASS.—Common migrant and local summer resident; rare winter resident near the coast (to Beverly, Fresh Pond, Cambridge). Mar. 27–Nov. 30; winter. Eggs, May 13–July 14.

R. I.—Common migrant and less common local summer resident; may winter locally. April 4–Nov. 14 (Dec. 25). Eggs, May 24–31.

CONN.—Common migrant and less common local summer resident; rare winter resident. April 5–Nov. 20 (winter).

306. *PASSERELLA ILIACA* (Merrem) Swainson.

**Fox sparrow; Fox-colored sparrow.**

*Fringilla iliaca* Merr., Beitr. z. besond. gesch. d. vögel, 1786–87, vol. 2, p. 40–41, pl. 10. "NORD-AMERIKA." Audubon, Ornith. biogr., 1835, vol. 2, p. 58, pl. 108.

Thickets and edges of woods; nests on the ground or in low trees.

ME.—Common migrant; accidental in winter. Mar. 15–April 17 (May 15); Oct. 8–Nov. 28 (Dec. 2, 9, 17, Seguin Id.; Jan. 24, 1909, Cape Elizabeth; Feb.).

N. H.—Common migrant. Mar. 16–April 23; Oct. 5–Nov. 13.

Vt.—Common migrant. Mar. 15–April 5; fall.

MASS.—Common migrant and rare winter resident near the coast. (Feb. 26, 1906, Springfield; Mar. 3) Mar. 9–April 25 (May 2); (Sept. 17, Oct. 2) Oct. 15–Nov. 29 (Dec. 6 and winter).

R. I.—Common migrant and occasional winter resident. Mar. 19–April 20 (May 21); Oct. 13–Nov. 27 (winter).

CONN.—Common migrant, and rare winter resident. Mar. 6–April 19; Oct. 10–Nov. 28 (winter).

307. *PIPILO ERYTHROPHthalmus* (Linné) Vieillot.

**Towhee; Chewink; Ground robin; Marsh robin; Pewink.**

Chapman, Handb. birds east. No. Amer., 1895, p. 310; Bird-life, 1898, pl. 55. Egg, Capen, 1886, pl. 10, fig. 13–15.

PASSERES.

OSCINES.

## FRINGILLIDAE.

*Fringilla erythrophthalma* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 180. "AMERICA."

Thickets and sprout land; nests on the ground, rarely in bushes.

ME.— Uncommon summer resident in southwestern counties. May 6–Oct. 18 (Nov. 21).

N. H.— Common summer resident in southern part, becoming uncommon north to White Mt. valleys. May 1–Oct. 15.

VT.— Uncommon summer resident. May–Sept.

MASS.— Common summer and very rare winter resident. (April 2) April 18–Oct. 22 (Nov. 30, Dec. 4); (winter). Eggs, May 17–June 12.

R. I.— Common summer and very rare winter resident. April 19–Oct. 13 (Jan. 14, 1896, Newport). Eggs, May 21–June 3.

CONN.— Common summer and very rare winter resident. April 25–Nov. 25 (Dec. 28, 1908, New Haven; Jan. 22, 1876, Portland). Eggs, May 21–June 6.

308. *CARDINALIS CARDINALIS* (Linné) Lichtenstein.

**Cardinal; Cardinal grosbeak; Red-bird; Virginia nightingale.**

*Loxia cardinalis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 172. "IN AMERICA SEPTENTRIONALI."

*Pitylus cardinalis* Linné. Audubon, Birds of Amer., 1841, vol. 3, p. 198, pl. 203. Egg, Capen, 1886, pl. 10, figs. 11, 12.

Bushy fields and thickets; nests in bushes.

ME.— Several recorded; probably some are escaped cage birds (♀ from flock of three, Gardiner, Dec. 19, 1895).

VT.— Rare visitor or escape: Brattleboro.

MASS.— Occasional visitor at all seasons and escape; doubtfully stated to have bred at Cambridge (a wild bird and a freed cage bird nested in Brookline, eggs, June 6, 1898).

CONN.— Occasional visitor or escape.



PASSERES.

OSCINES.

## FRINGILLIDAE.

309. *ZAMELODIA LUDOVICIANA* (Linné) Coues.**Rose-breasted grosbeak; Throat-cut.**

*Loxia ludoviciana* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 306.  
"IN LUDOVICIA."

*Ihabia ludoviciana* (Linné). Chapman, Handb. birds east. No. Amer., 1895, p. 312; Bird-life, 1898, pl. 54; Stone, Auk, 1899, vol. 16, pl. 4. Egg, Capen, 1886, pl. 10, figs. 9, 10.

Orchard and shade trees and borders of woods near water; nests in bushes and trees.

ME.—Common summer resident except in eastern part. (April 26) May 4–Aug. 27. Eggs, June 10–30.

N. H.—Common summer resident at lower elevations. May 12–Sept. 18.

VT.—Common summer resident. May–Sept. 14. Eggs, June 2–23.

MASS.—Common summer resident. (April 26) May 2–Oct. 3. Eggs, May 25–June 15.

R. I.—Common summer resident, mainly in northern part. May 4–Oct. 2. Eggs, May 23–June 4.

CONN.—Common summer resident; accidental in winter. May 1–Sept. (Nov. 25; Jan. 18, 1893, Stonington). Eggs, May 24.

310. *GUIRACA CAERULEA* (Linné) Swainson.**Blue grosbeak.**

Chapman, Handb. birds east. No. Amer., 1895, p. 313. McAtee, U. S. dept. agric., biol. surv., bull. 32, pl. 4.

*Loxia caerulea* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 175.  
"AMERICA."

Brush-grown fields and thickets; nests in bushes or weedy growths.

ME.—Recorded from Calais, but in error for Grand Menan, N. B.

## PASSERES.

## OSCINES

## FRINGILLIDAE.

N. H.—Doubtfully recorded from East Derry, May 26, 1894.

Vt.—Doubtfully recorded.

MASS.—Accidental visitor: Brookline, May 29, 1880.

R. I.—Accidental visitor: Drownville, Oct. 12, 1899.

311. *PASSERINA CYANEA* (Linné) Vieillot.

**Indigo bunting; Indigo-bird.**

*Tanagra cyanea* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 315.  
“IN CAROLINA.”

*Spiza cyanea* Wils. Audubon, Birds of Amer., 1841, vol. 3, p. 96, pl. 170. Egg, Capen, 1886, pl. 10, fig. 8.

Brush-grown fields, thickets, and weedy tangles; nests among low bushes, brambles, or tall weeds.

ME.—Common summer resident in southern, and rare in northern counties (Caribou). May 11–Oct. 2. Eggs, June–July 15.

N. H.—Common migrant and summer resident at lower elevations. May 9–Oct. 1.

Vt.—Common migrant and summer resident. May 18–fall. Eggs, May 30.

MASS.—Common migrant and summer resident. May 8–Oct. 29 (Dec. 5, 1871, Watertown). Eggs, June 3–22.

R. I.—Common summer resident, mainly in northern part. May 10–Oct. Eggs, June 1–11.

CONN.—Common migrant and summer resident. May 9–Oct. Eggs, June 16.

312. *SPIZA AMERICANA* (Gmelin) Ridgway.

**Dickcissel; Black-throated bunting; Judas-bird.**

*Emberiza americana* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 872.  
“IN NOVEBORACO.” Audubon, Birds of Amer., 1841, vol. 3, p. 58, pl. 156. Egg, Capen, 1886, pl. 10, fig. 7.

Grassy fields; nests on the ground or in low bushes.

PASSERES.

OSCINES.

## FRINGILLIDAE.

ME.—Accidental visitor: Job's Id., Penobscot Bay, Sept. 29, 1884; St. George; Westbrook, Oct. 10, 1888.

MASS.—Formerly an uncommon summer resident along coast and in western valleys; now unknown. May 15–Oct. 3. Eggs, June 9, 1877 (Medford)—Aug. 1, 1878 (2d brood, Hyde Park).

R. I.—Formerly a rare migrant: Newport, Sept. 25, 1888.

CONN.—Formerly a summer resident, now unknown.

## 313. CALAMOSPIZA MELANOCORYS Stejneger.

**Lark bunting.**

Stejn., Auk, 1885, vol. 2, p. 49. Based on Townsend: "PLAINS OF THE PLATTE RIVER."

*Corydalina bicolor* Towns. Audubon, Birds of Amer., 1841, vol. 3, p. 195, pl. 202.

Prairies; nests on the ground.

MASS.—Accidental visitor: Lynn, Dec. 5, 1877.

## TANGARIDAE.

## 314. PIRANGA LUDOVICIANA (Wilson) Richardson.

**Louisiana tanager; Western tanager.**

*Tanagra ludoviciana* Wils., Amer. ornith., 1811, vol. 3, p. 27, pl. 20, fig. 1. "LOUISIANA."

*Piranga ludoviciana* (Wils.). Audubon, Birds of Amer., 1841, vol. 3, p. 231, pl. 210.

Mountain forests; nests in trees.

ME.—Accidental visitor: near Bangor, about Oct. 1, 1889.

MASS.—Accidental visitor: Salem, Jan. 20, 1878.

CONN.—Accidental visitor: New Haven, Dec. 15, 1892.

## 315. PIRANGA ERYTHROMELAS Vieillot.

**Scarlet tanager; English robin (Berkshires); Fire-bird; Pocket-bird; War-bird.**

*Piranga erythromelas* Vieill., Nouv. dict. d'hist. nat., 1819, vol. 28, p. 293. "AMÉRIQUE."

PASSERES.

OSCINES.

## TANGARIDAE.

*Pyrrhuloxia rubra* (Linné). Audubon, Birds of Amer., 1841, vol. 3, p. 226, pl. 209. Egg, Capen, 1886, pl. 6, figs. 6, 7.

Deciduous woods, especially oaks and chestnuts; nests in trees.

ME.— Rare summer resident. May 12–Oct. 5.

N. H.— Uncommon summer resident, mainly below 2000 ft. south of White Mts. May 13–Sept. 25.

VT.— Uncommon summer resident. May–fall.

MASS.— Common summer resident. (April 30, May 4) May 8–Oct. 16. Eggs, June 2–17.

R. I.— Common summer resident, mainly of northern part. (April 4, 1891, Apponaug) May 4–Oct. 9. Eggs, May 22–June 5.

CONN.— Common summer resident. May 10–Sept. 11. Eggs, May 31–June 17.

316. *PIRANGA RUBRA* (Linné) Vieillot.

**Summer tanager; Summer redbird.**

*Fringilla rubra* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 181. "AMERICA."

*Pyrrhuloxia aestiva* Gmel. Audubon, Birds of Amer., 1841, vol. 3, p. 222, pl. 208.

Open deciduous and mixed woods; nests in trees.

ME.— Accidental visitor: Portland, May 18, 1906; Wiscasset.

N. H., VT.— Doubtfully recorded.

MASS.— Accidental summer visitor. April 21, May, June, July, Aug.

R. I.— Accidental summer visitor: Providence (two seen); Seaconnet, April 27, 1901; Ten Mile River.

CONN.— Rare visitor. April, July.

## HIRUNDINIDAE.

317. *PROGNE SUBIS* (Linné) Baird.

**Purple martin; Black martin; Cape Cod swallow; House martin; Martin.**

PASSERES.

OSCINES.

## HIRUNDINIDAE.

*Hirundo subis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 192. "AD SINUM HUDSONIS."

*Hirundo purpurea* Linné. Audubon, Birds of Amer., 1840, vol. 1, p. 170, pl. 45. Egg, Capen, 1886, pl. 7, fig. 1.

Open country and vicinity of dwellings; nests in cavities or bird houses.

ME.—Till recently a common local summer resident. April 22–Sept. 4 (Sept 30). Eggs, June 5–July 4.

N. H.—Local summer resident north to White Mt. valleys; now uncommon. April 19–Sept.

VT.—Till recently a common local summer resident. April 25–Sept. 20. Eggs, –June 23.

MASS.—Till recently a common local summer resident. April 14–Sept. 30. Eggs, May 30–June 21.

R. I.—Uncommon migrant and summer resident; formerly more common. April 25–Oct. 8. Eggs, May 25–June 4.

CONN.—Uncommon migrant and local summer resident; formerly more common. April 17–fall. Eggs, May 21.

## 318. PETROCHELIDON LUNIFRONS (Say) Cassin.

**Cliff swallow; Eave swallow; Mud swallow; Republican.**

*Hirundo lunifrons* Say, Long's Exped. to Rocky Mts., 1823, vol. 2, p. 47. "ROCKY MOUNTAINS, ABOUT THE SOURCES OF THE ARKANSAS AND PLATTE."

*Hirundo fulva* Vieill. Audubon, Birds of Amer., 1840, vol. 1, p. 177, pl. 47. Egg, Capen, 1886, pl. 6, fig. 11–13.

Open country; nests in colonies under cliffs or eaves of outbuildings.

ME.—Common migrant and summer resident. (April 11, Bangor) April 21–Sept. 6 (Sept. 19). Eggs, June 1–July 2.

N. H.—Common migrant and local summer resident. April 28–Sept. 1.

PASSERES.

OSCINES.

## HIRUNDINIDAE.

VT.—Common migrant and summer resident. Late April–summer.

MASS.—Common migrant and less common local summer resident. April 23–Sept. 4 (Sept. 30). Eggs, May 25–July 20.

R. I.—Uncommon migrant and summer resident. April 28–Sept. 15. Eggs, May 31–July 16.

CONN.—Common migrant and less common local summer resident. (April 15) April 20–Sept. Eggs, May 26–? (2 broods).

319. *HIRUNDO ERYTHROGASTER* Boddaert.

**Barn swallow; Fork-tailed swallow.**

Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 45. Description in Buffon: "CAYENNE."

*Hirundo rustica* Linné. Audubon, Birds of Amer., 1840, vol. 1, p. 181, pl. 48. Egg, Capen, 1886, pl. 6, figs. 8, 9.

Open country; nests in crevices of cliffs, but now generally on rafters of old barns.

ME.—Common migrant and summer resident. April 19–Sept. 19 (Oct. 29, Seguin Id.; Oct. 31, 1907, Portland). Eggs, May 28–June 24.

N. H.—Common migrant and summer resident at lower elevations. April 20–Sept. 21.

VT.—Common migrant and summer resident. April 25–fall.

MASS.—Common migrant, less common summer resident. (April 5, 1893, Waltham) April 17–Sept. 30 (Oct. 7). Eggs, May 18–July 12.

R. I.—Common migrant and summer resident; still breeds in cliffs at Middletown. (April 10)–Sept. 23. Eggs, May 18–30.

CONN.—Common migrant and summer resident. April 21–Sept. Eggs, (!Sept. 3).

320. *IRIDOPROCNE BICOLOR* (Vieillot) Coues.

**Tree swallow; Little martin; White-bellied swallow.**

*Hirundo bicolor* Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 1,

PASSERES.

OSCINES.

## HIRUNDINIDÆ.

p. 61, pl. 31. "ÉTATS-UNIS." Audubon, Birds of Amer., 1840, vol. 1, p. 175, pl. 46. Egg, Capen, 1886, pl. 6, fig. 10.

Open country, especially near water; nests in cavities of dead trees or in bird boxes.

ME.—Common migrant and summer resident. (Mar. 19) April 4–Sept. 21. Eggs, May 30–June 11.

N. H.—Common migrant and less common summer resident at lower elevations. Mar. 23–Sept.

VT.—Common migrant and less common summer resident. Eggs, June 1.

MASS.—Common migrant and less common summer resident. (Mar. 15) Mar. 22–Oct. 17 (Nov. 2). Eggs, May 14–June 15.

R. I.—Common migrant and less common summer resident; casual in winter. (Mar. 12) Mar. 28–Oct. 18. Eggs, May 24–30.

CONN.—Common migrant and less common summer resident. Mar. 25–Oct. 17 (Oct. 22). Eggs, June.

321. *RIPARIA RIPARIA* (Linné) Sharpe and Wyatt.

**Bank swallow; Sand martin or swallow.**

*Hirundo riparia* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 192. "IN EUROPAE COLLIBUS ARENOSIS ABRUPTIS." Audubon, Birds of Amer., 1840, vol. 1, p. 187, pl. 50. Egg, Capen, 1886, pl. 6, fig. 14.

Open country near rivers and lakes; nests in colonies in burrows dug in sandy bluffs.

ME.—Common migrant and summer resident. (April 19) April 25–Sept. 1 (Sept. 19). Eggs, May 29–June 24.

N. H.—Common migrant and local summer resident. May–Aug. 24. Eggs, May 26.

VT.—Common migrant and local summer resident. April 28–summer.

PASSERES.

OSCINES.

## HIRUNDINIDAE.

MASS.—Common migrant and local summer resident. April 24–Sept. 2 (Nov. 2). Eggs, May 28–June 17.

R. I.—Common migrant and local summer resident. April 25–Aug. 29. Eggs, May 26–June 6.

CONN.—Common migrant and local summer resident. April 21–summer.

322. *STELGIDOPTERYX SERRIPENNIS* (Audubon) Baird.**Rough-winged swallow.**

Chapman, Handb. birds east. No. Amer., 1895, p. 322, fig. 90. Egg, Capen, 1886, pl. 6, fig. 15.

*Hirundo serripennis* Aud., Ornith. biogr., 1838, vol. 4, p. 593. "A FEW MILES FROM BAYOU SARA," LA. Aud., Birds of Amer., 1840, vol. 1, pl. 51.

Open country near water; nests in holes in banks, or in cavities of cliffs and walls.

N. H.—Rare summer resident: near Hanover, 1905, 1906.

VT.—Rare summer resident (noted at Bennington; Lunenburg; Norwich, May 6, 1905, April 29, 1906; West Danville, June 11, 1901; White River Junction). April 29–summer. Eggs, May 16.

MASS.—Rare local summer resident in western part (Glendale, young, 1906; Longmeadow, nesting, May, 1906; North Adams, two nesting pairs, June, 1895). May–summer.

CONN.—Rare local summer resident. (April 12) May 9–summer. Eggs, May 26–June 9.

## BOMBYCILLIDAE.

323. *BOMBYCILLA GARRULA* (Linné) Vieillot.**Bohemian waxwing.**

Bonaparte, Amer. ornith., 1828, vol. 3, p. 14, pl. 16, fig. 2. Egg, Ibis, 1861, ser. 1, vol. 3, pl. 4, fig. 1–6.



PASSERES.

OSCINES.

## BOMBYCILLIDAE.

*Lanius garrulus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 95.  
 "IN EUROPA & AMERICA BOREALI; AESTIVAT SUPRA SVECIAM."  
 Evergreen forests; nests in trees.

ME.—Rare and irregular winter visitor. Dec. 6–Feb. 6.

N. H.—Rare winter visitor: Hampton Falls, Feb. 17 and 21, 1897; Nashua, 1880.

VT.—Rare winter visitor: Brattleboro, April, 1901; Burlington (*bis*); St. Albans (*bis*). Nov. 25–April.

MASS.—Rare winter visitor: about six authentic records. Dec. 26–Feb. 18.

CONN.—Accidental winter visitor: East Windsor Hill, near Hartford; ? New Haven, Feb. 11, 1875.

## 324. BOMBYCILLA CEDRORUM Vieillot.

**Cedar waxwing; Apple-bird; Canada robin; Cedar-bird; Cherry-bird.**

Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 1, p. 38, pl. 57. "EX AMÉRIQUE DEPUIS LE CANADA JUSQU'AU MEXIQUE."

*Ampelis cedrorum* (Vieill.). Chapman, Handb. birds east. No. Amer., 1895, p. 323; Bird-life, 1898, pl. 57. Egg, Capen, 1886, pl. 7, figs. 2, 3.

Orchards, shade trees, and woods; nests in trees.

ME.—Common migrant and summer resident; rare in winter. Eggs, June 29.

N. H.—Common migrant, and at lower elevations a common summer resident; rare in winter in southern part. Eggs, July 29.

VT.—Common migrant and summer resident; uncommon in winter in southern part.

MASS.—Common migrant and less common summer resident; irregularly common in winter but generally absent in Nov. and Dec. Eggs, May 30–July.

R. I.—Common migrant and summer resident; irregular in winter. Eggs, June 7–20.

PASSERES.

OSCINES.

## BOMBYCILLIDAE.

CONN.— Common migrant and summer resident; irregular in winter. Eggs, June 20–Aug. 1 (Aug. 27).

## LANIIDAE.

325. *LANIUS BOREALIS* Vieillot.**Northern shrike; Butcher-bird.**

Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 1, p. 80, pl. 50.  
 “CANADA ET SUR-TOUT LA BAIE D’HUDSON.” Audubon, Birds of Amer., 1842, vol. 4, p. 130, pl. 236.

Open country and edges of woods; nests in trees.

ME.— Uncommon winter resident; supposed breeding records refer to *L. l. migrans*. Oct. 15–April 17.

N. H.— Uncommon winter resident at lower elevations. Nov. 12–April.

VT.— Uncommon winter resident. Fall–April 13.

MASS.— Uncommon winter resident. (Oct. 6) Oct. 12–April 16 (May).

R. I.— Uncommon, irregular winter resident. Nov. 28–Mar. 23.

CONN.— Irregular winter resident, sometimes common. Oct. 29–Mar. 24 (April 11).

326. *LANIUS LUDOVICIANUS* Linné.**Loggerhead shrike.**

Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 134. “IN LUDOVICIA.”  
 Ridgway, Bull. 50, U. S. nat. mus., 1904, pt. 3, p. 241. Audubon, Birds of Amer., 1842, vol. 4, pl. 237.

Open country; nests in trees.

CONN.— Accidental visitor: South Norwalk, Feb. 17, 1905 (J. A. Allen).

PASSERES.

OSCINES.

## LANIIDAE.

327. *LANIUS LUDOVICIANUS MIGRANS* W. Palmer.

**Migrant shrike; Butcher-bird; Cricket hawk; Grasshopper hawk; Joree; Mouse hawk.**

W. Palmer, *Auk*, 1898, vol. 15, p. 248. "KINGSTON, ONTARIO."  
Egg, Capen, 1886, pl. 7, figs. 11, 12.

Orchards and open tree growth; nests in trees.

ME.—Uncommon local summer resident. (Mar. 1) Mar 24–Sept. (Nov. 17). Eggs, May 5–25.

N. H.—Uncommon summer resident in valley country; accidental in winter. Mar. 20–Sept. (Jan. 20, Concord). Eggs, April 26–May 5.

VT.—Uncommon local summer resident. Mar. 3–fall. Eggs, April 28–May 23.

MASS.—Rare migrant and in western part, rare summer resident; very rare in winter. Mar. 29–Nov. 27 (Dec. 19, Jan., Feb.). Eggs, May 10 (Williamstown).

R. I.—Rare migrant. (? Spring); Aug. 29–Sept. 2.

CONN.—Rare migrant and winter resident (Portland, Nov. 1876; Saybrook, Jan., 1879); possibly a rare summer resident (Mt. Carmel, May 24, 1873). Spring–(May 24); Aug. 29–winter.

## VIREONIDAE.

328. *VIREOSYLVA OLIVACEA* (Linné) Bonaparte.

**Red-eyed Vireo; Politician; Preacher; Red-eyed greenlet.**

*Muscicapa olivacea* Linné, *Syst. nat.*, ed. 12, 1766, vol. 1, p. 327.  
"IN AMERICA SEPTENTRIONALI."

*Vireo olivaceus* (Linné). DeKay, *Zool. New York*, 1844, vol. 2, p. 124, pl. 36, fig. 79. Egg, Capen, 1886, pl. 7, figs. 4, 5.

Orchards and deciduous shade and forest trees; nest pensile in trees or bushes.

PASSERES.

OSCINES.

## VIREONIDAE.

ME.—Common summer resident. (May 1) May 7–Oct. 2 (Oct. 11). Eggs, –June 29.

N. H.—Common summer resident below 3000 ft. May 8–Sept. 30.

VT.—Common summer resident. May 9–fall.

MASS.—Common summer resident. May 4–Oct. 11 (Oct. 26; Nov. 2, 1870, Watertown). Eggs, May 25–July 20.

R. I.—Common summer resident. (April 29) May–Sept. 24 (Oct. 11). Eggs, May 28–June 6.

CONN.—Common summer resident. May 6–Oct. 2. Eggs, June 1–12.

## 329. VIREOSYLVA PHILADELPHICA Cassin.

**Philadelphia Vireo; Brotherly-love Vireo.**

Cass., Proc. acad. nat. sci. Phila., 1851, vol. 5, p. 153, pl. 10, fig. 2. “NEAR PHILADELPHIA.”

*Vireo philadelphicus* (Cass.). Dwight, Auk, 1897, vol. 14, p. 269, pl. 2. Egg, Brewster, Auk, 1905, vol. 22, pl. 11.

Open deciduous growth, especially of birch and poplar; nest pensile in trees or bushes.

ME.—Uncommon summer resident in northern part (Bucksport, Lake Umbagog). May 21–Oct. 2. Eggs, June 15.

N. H.—Uncommon summer resident north of White Mts., elsewhere rare migrant. May 26–Sept. 29.

VT.—Rare migrant: Breadloaf, Sept. 19, 1900; Morrisville; Pittsford, Sept. 11, 1889. (Reported nesting at Brattleboro, June, 1901.) May; Sept. 11–19.

MASS.—Rare migrant. May (? 10)–17; Sept. 7–27.

R. I.—Rare migrant: Drownville, May 31, 1891; West Greenwich, May 21, 1902.

CONN.—Rare migrant: New Haven, May 19, 1888; Portland, Sept. 21, 1893, Sept. 17, 1894; (? Stratford, “*Vireo gilvus*. . . . under parts. . . . mostly bright yellow”).

PASSERES.

OSCINES.

## VIREONIDAE.

330. *VIREOSYLVA GILVA* (Vieillot) Cassin.**Warbling Vireo.**

*Muscicapa gilva* Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 1, p. 65, pl. 34. "L'ÉTAT DE NEW-YORK."

*Vireo gilvus* (Vieill.). Chapman, Handb. birds east. No. Amer., 1895, p. 329. Dwight, Auk, 1897, vol. 14, pl. 2. Egg, Capen, 1886, pl. 7, fig. 6.

Orchard and shade trees, especially elms; nest pensile in trees.

ME.—Uncommon local summer resident. May 5–Sept. 15 (Oct. 7).

N. H.—Uncommon local summer resident north to White Mt. valleys. May 1–Sept. 17.

VT.—Uncommon local summer resident. May 3–

MASS.—Common summer resident. (April 24, 1908, Auburn-dale) April 29–Sept. 26. Eggs, May 30–June 11.

R. I.—Common summer resident, especially of village shade trees. May 5–Sept. 25. Eggs, May 28–30.

CONN.—Common summer resident. (April 20, 1898, Jewett City) April 29–Oct. 13 (Oct. 25, 1900, Hartford). Eggs, June 2–17.

331. *LANIVIREO FLAVIFRONS* (Vieillot) Lawrence.**Yellow-throated Vireo.**

*Vireo flavifrons* Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 1, p. 85, pl. 54. "ÉTATS-UNIS." Audubon, Birds of Amer., 1842, vol. 4, p. 141, pl. 238. Egg, Capen, 1886, pl. 7, figs. 7, 8.

Orchard and shade trees and edges of woods; nest pensile in trees.

ME.—Rare summer resident in southwestern counties. Mar. 12–Sept.

N. H.—Uncommon summer resident becoming rare north to White Mt. valleys. May 8–Sept. Eggs, June 14.

PASSERES.

OSCINES.

## VIREONIDAE.

VT.— Uncommon summer resident (East Bethel).

MASS.— Common summer resident becoming local in western part. May 1–Sept. 19. Eggs, May 24–June 17.

R. I.— Common summer resident. May 1–Sept. Eggs, May 24–June 2.

CONN.— Common migrant and summer resident. May 6–Sept 17. Eggs, May 21–June 11.

332. *LANIVIREO SOLITARIUS* (Wilson) Allen.

**Blue-headed Vireo; Solitary Vireo.**

*Muscicapa solitaria* Wils., Amer. ornith., 1810, vol. 2, p. 143, pl. 17, fig. 6. "NEAR PHILADELPHIA."

*Vireo solitarius* Vieill. Audubon, Birds of Amer., 1842, vol. 4, p. 144, pl. 239. Egg, Capen, 1886, pl. 7, fig. 9.

Cool mixed woods; nest pensile in trees.

ME.— Uncommon summer resident mainly in northern part. (April 24) April 28–Oct. 6 (Nov.).

N. H.— Uncommon summer resident up to 3000 ft. April 29–Oct. 7. Eggs, May 29–late June.

VT.— Uncommon summer resident.

MASS.— Uncommon migrant and rare summer resident, becoming local in south and east. (April 11, 1908, Stoughton) April 14–Oct. 17. Eggs, May 13–June 26.

R. I.— Uncommon migrant and rare local summer resident (Washington, West Greenwich). April 23–Oct. 22. Eggs, May 15–June 15.

CONN.— Uncommon migrant and rare summer resident (Eastford, near New Haven, Norfolk, West Simsbury). Late April–Oct. 13. Eggs, May 27–June 23.

333. *VIREO GRISEUS* (Boddaert) A. O. U. comm.

**White-eyed Vireo; White-eye.**

*Tanagra grisea* Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 45. Based on Buffon.

PASSERES.

OSCINES.

## VIREONIDAE.

*Vireo noveboracensis* (Gmel.). DeKay, Zool. New York, 1844, vol. 2, p. 122, pl. 36, fig. 78. Egg, Capen, 1886, pl. 7, fig. 10.

Bushy thickets; nest pensile in bushes.

ME.—Doubtfully recorded.

N. H.—Several unsatisfactory records: Chesterfield, Manchester.

VT.—Accidental visitor: Brattleboro; ? St. Johnsbury, various records (Clark); Townshend, 1893.

MASS.—Rare local summer resident near coast and in southern valleys (Sheffield). May 2–Sept. 27 (Oct. 30). Eggs, May 22–June 18.

R. I.—Uncommon local summer resident. May 7–Sept. 18. Eggs, June 2, 3.

CONN.—Common summer resident in southern part and Connecticut valley. May 2–Sept. Eggs, June 9–12.

334. *VIREO BELLII* Audubon**Bell's Vireo.**

Aud., Birds of Amer., 1844, vol. 7, p. 333, pl. 485. "SHORES AND PRAIRIES OF THE UPPER MISSOURI."

Bushy thickets; nest pensile in bushes or low trees.

N. H.—Accidental visitor: Durham, Nov. 19, 1897.

## MNIOTILTIDAE.

335. *MNIOTILTA VARIA* (Linné) Vieillot.**Black and white warbler; Black and white creeper.**

Chapman, Warblers No. Amer., 1907, p. 39, pl. 2, figs. 4, 5. Egg, Capen, 1886, pl. 3, figs. 2, 3.

*Motacilla varia* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 333. "IN JAMAICA, DOMINICA."

Deciduous and mixed woods; nests on the ground.

## PASSERES.

## OSCINES.

## MNIOTILTIDAE.

ME.— Common migrant and local summer resident. (April 27) May 3–Oct. 5. Eggs, May 25–June 21.

N. H.— Common migrant and less common summer resident at lower altitudes. April 27–Sept. 29. Eggs, June 24.

VT.— Common migrant and summer resident. May 5–fall. Eggs, June 20.

MASS.— Common migrant and less common summer resident. (April 18) April 28–Oct. 29 (Dec. 5). Eggs, May 17–June 22.

R. I.— Common migrant and less common summer resident. (April 15) May 1–Oct. 7. Eggs, May 21, 23.

CONN.— Common migrant and less common summer resident. (April 20) April 25–Oct. 10. Eggs, May 20–June 4.

336. *PROTONOTARIA CITREA* (Boddaert) Baird.**Prothonotary warbler; Golden swamp warbler.**

Chapman, Warblers No. Amer., 1907, p. 54, pl. 3, fig. 3–4.

*Motacilla citrea* Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 44. Description in Buffon: "LOUISIANE."

Bushy and tree-grown swamps; nests generally in cavities of stumps or dead trees.

ME.— Accidental visitor: Calais, Oct. 30, 1862.

VT.— Doubtfully recorded.

MASS.— Accidental visitor. May 1–Sept. 15.

R. I.— Accidental visitor: Lonsdale, April 29, 1892, April 19, 1893; South Kingston, April 20 or 21, 1884.

337. *HELMITHEROS VERMIVORUS* (Gmelin) Bonaparte.**Worm-eating warbler.**

*Motacilla vermicora* [= *vermicora*] Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 951. "IN PENNSYLVANIA."

*Hclinaia vermicora* Lath. Audubon, Birds of Amer., 1841, vol. 2, p. 86, pl. 105. Egg, Capen, 1886, pl. 3, fig. 6.



PASSERES.

OSCINES.

## MNIOTILTIDAE.

Dry wooded slopes and ravines with undergrowth; nests on the ground.

ME., N. H.—Unsatisfactorily recorded.

VT.—Accidental visitor: St. Albans, 1891.

MASS.—Accidental visitor: Cambridge, Sept. 19, 1881; Easthampton; Salem Willows, April 14–16, 1902; Taunton, May 9, 1890.

CONN.—Rare summer resident. May 10–Aug. 22. Eggs, (New Haven, Saybrook) May 25–June 19.

338. *HELMINTHOPHILA PINUS* (Linné) Ridgway.

**Blue-winged warbler; Blue-winged yellow warbler.**

Chapman, Warblers No. Amer., 1907, p. 65, pl. 5, fig. 1–2. Egg, Capen, 1886, pl. 3, fig. 7.

*Certhia pinus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 187. "IN AMERICA SEPTENTRIONALI."

Low woods and sprout land; nests on the ground.

VT.—Unsatisfactorily recorded.

MASS.—Very rare visitor. May 12–17 (? 29).

R. I.—Very rare summer resident: nest at Gloucester, May 30, 1890.

CONN.—Locally common summer resident in southern part and Connecticut valley. (May 2) May 9–late Aug. Eggs, May 20–June 20.

339. *HELMINTHOPHILA CHRYSOPTERA* (Linné) Ridgway.

**Golden-winged warbler.**

Chapman, Warblers No. Amer., 1907, p. 60, pl. 5, fig. 5–6. Egg, Capen, 1886, pl. 3, fig. 8.

*Motacilla chrysoptera* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 333. "IN PENNSYLVANIA."

Low bushy growths and sprout lands; nests on the ground.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

N. H.— Rare in spring and early summer in southernmost part; may breed. May, June.

Vt.— Rare migrant; ? “rare summer resident” (Perkins). Spring; (Oct. 26, 1907, Woodstock).

MASS.— Uncommon summer resident, mainly in eastern part. (May 4) May 7–Sept. 1 (Oct. 8). Eggs, June 1–20.

R. I.— Rare summer resident.

CONN.— Uncommon summer resident. (May 3) May 12–Aug. 24. Eggs, May 29–June 4.

340. HELMINTHOPHILA RUBRICAPILLA (Wilson) Faxon.

**Nashville warbler.**

*Sylvia rubricapilla* Wils., Amer. ornith., 1811, vol. 3, p. 120, pl. 27, fig. 3; vol. 6, p. xv. “TENNESSEE...NOT FAR FROM NASHVILLE.”

*Helinaia rubricapilla* (Wils.). Audubon, Birds of Amer., 1841, vol. 2, p. 103, pl. 113. Egg, Capen, 1886, pl. 3, figs. 9, 10.

Open woodland and thickets bordering swamps; nests on the ground.

ME.— Common migrant and summer resident. (April 30) May 4–Sept. 27 (Oct. 4). Eggs, June 3–July 10.

N. H.— Common migrant and summer resident up to 4500 ft. (April 30) May 4–Sept. 13. Eggs, May 25–June 8.

Vt.— Common migrant and summer resident. May 3–fall.

MASS.— Common migrant and less common summer resident, accidental in winter. (April 20, 28) May 2–Sept. 25 (Oct. 14; Jan. 1890, Swampscott). Eggs, May 21–June 21.

R. I.— Common migrant and uncommon summer resident. May 4–Sept. 17. Eggs, May 22–June 1.

CONN.— Common migrant and uncommon summer resident. (April 22, 1889, Bridgeport) May 5–Sept. Eggs, May 30–June 8.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

## 341. HELMINTHOPHILA CELATA (Say) Ridgway.

**Orange-crowned warbler.**

Chapman, Warblers No. Amer., 1907, p. 86, pl. 7, figs. 3, 4.

*Sylvia celatus* Say, Long's Exped. to Rocky Mts., 1823, vol. 1, p. 169. "ENGINEER CANTONMENT," IOWA.

Wooded and bushy country; nests on or near the ground.

ME.—Unsatisfactorily accredited (Brunswick).

N. H.—Very rare migrant: Hollis, May 16, 1876 (*see* Isles of Shoals).

MASS.—Very rare spring and late fall migrant; accidental in winter. May 15 (Springfield); Sept. 30–Nov. 29 (Jan. 1, 1875, Lynn).

R. I.—Very rare spring and late fall migrant: Cranston, Dec. 3, 1874; East Providence, May 9, 1891.

CONN.—Rare migrant. May 8, 1888 (East Hartford); Oct.–Nov.

## 342. HELMINTHOPHILA PEREGRINA (Wilson) Ridgway.

**Tennessee warbler.**

Chapman, Warblers No. Amer., 1907, p. 83, pl. 7, figs. 1, 2.

*Sylvia peregrina* Wils., Amer. ornith., 1811, vol. 3, p. 83, pl. 25, fig. 2. "BANKS OF THE CUMBERLAND RIVER, . . . TENNESSEE."

Dense thickets and edges of woods; nests near the ground.

ME.—Rare migrant and in northern part, a rare summer resident. May 15–26; summer. Eggs, June 4.

N. H.—Rare migrant and from White Mts. north, a rare local summer resident (Franconia, Lake Umbagog). May 20–Sept. 9.

VT.—Rare migrant and possible summer resident (Mt. Holly, two pairs, June, 1897, above 2000 ft.).

MASS.—Rare migrant and possibly a very rare summer resident (Mt. Greylock, July 15 and 16, 1888; ? Springfield, very doubtful). May 10–30; (July 15, 16); Aug.–Sept. 10.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

R. I.— Rare migrant. May 18; Sept. 18.

CONN.— Rare migrant. May 19–27 (June 8); Sept. 24–(? Nov., Clark).

343. *COMPSOTHTLYPIS AMERICANA USNEAE* Brewster.

**Northern Parula warbler; Blue yellow-backed warbler; Usnea warbler.**

Brewst., Auk, 1896, vol. 13, p. 44. "LAKE UMBAGOG, MAINE." Chapman, Warblers No. Amer., 1907, p. 104, pl. 8, fig. 1–3. Egg, Capen, 1886, pl. 3, figs. 4, 5.

Forests and open tree growth; nests in pendulous masses of Usnea lichen.

ME.— Common migrant and uncommon local summer resident. May 5–Sept. 30. Eggs, May 31–June 17.

N. H.— Common migrant and less common summer resident below 3000 ft. (April 30) May 3–Oct. 1. Eggs, May 31–June 12.

VT.— Common migrant and less common summer resident. May 2–fall.

MASS.— Common migrant and less common local summer resident. (April 25) May 1–Oct. 10 (Nov. 19, 1881, Cambridge). Eggs, May 30–July 5.

R. I.— Common migrant and uncommon local summer resident. April 30–Oct. 7. Eggs, June 2–5.

CONN.— Common migrant and uncommon local summer resident. May 2–Oct. 15 (Oct. 20). Eggs, (May 18) May 29–June 22 (Aug.).

344. *DENDROICA TIGRINA* (Gmelin) Baird.

**Cape May warbler.**

Chapman, Handb. birds east. No. Amer., 1895, p. 352. Egg, Capen, 1886, pl. 4, fig. 15.

*Motacilla tigrina* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 985. "CANADA."

PASSERES.

OSCINES.

## MNIOTILTIDAE.

*Sylvicola maritima* Wils. Audubon, Birds of Amer., 1841, vol. 2, p. 44, pl. 85.

Northern coniferous forests in summer, and open tree growth in migration; nests in trees.

ME.— Rare migrant and in northern part a rare summer resident. May 6–24; summer–Aug. 22.

N. H.— Rare migrant and north of White Mts. a rare summer resident (Franconia, June 3). May 10–Sept.

VT.— Rare migrant and (?) summer resident (Mt. Killington, Ross).

MASS.— Rare migrant. (May 2) May 10–25; Aug. 18–Sept. 15 (Oct. 9).

R. I.— Rare migrant. May 13–20; Sept.

CONN.— Rare migrant. May 8–26; Aug. 28.

345. *DENDROICA AESTIVA* (Gmelin) Baird.

**Yellow warbler; Cotton wren; Garden warbler; Golden warbler; Spy-about; Summer yellow-bird; Wild canary; Yellow-belly; Yellow-bird.**

*Motacilla aestiva* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 996.  
“IN GUJANA, AESTATE IN CANADA.”

*Sylvicola aestiva* (Gmel.). Audubon, Birds of Amer., 1841, vol. 2, p. 50, pl. 88. Egg, Capen, 1886, pl. 3, fig. 11–14.

Orchards, shade trees, and bushy swamps; nests in trees and bushes.

ME.— Common summer resident. (April 30) May 3–Sept. 10. Eggs, May 29–June 30 (July).

N. H.— Uncommon local summer resident of valleys and bottomlands; rare from White Mts. north. May 4–Sept. 11. Eggs, June 7.

VT.— Common summer resident of valleys and bottomlands. May 4–Sept.

MASS.— Common migrant and summer resident. (April 23)

## PASSERES.

## OSCINES.

## MNIOTILTIDAE.

May 1–Sept. 30 (Nov. 28, 1891, Cambridge). Eggs, May 20–June 14.

R. I.—Common summer resident. (April 17, 29) May 1–Sept. Eggs, May 25–June 4.

CONN.—Common migrant and summer resident. (April 30) May 1–Sept. Eggs, May 20–June 30.

346. *DENDROICA CAERULESCENS* (Gmelin) Baird.**Black-throated blue warbler.**

Chapman, Warblers No. Amer., 1907, p. 133, pl. 9, fig. 1–3. Egg, Capen, 1886, pl. 4, fig. 1.

*Motacilla caerulescens* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 960. "IN INSULA S. DOMINICI."

Thickets and forest undergrowth; nests near the ground.

ME.—Common migrant, and less common summer resident. May 4–late Oct. Eggs, June 10.

N. H.—Common migrant, and from White Mts. north a less common summer resident, becoming rare and local south. May 10–Oct. 10. Eggs, June 19.

VT.—Uncommon summer resident in northern part. May 5–fall.

MASS.—Common migrant and at higher elevations of western part (Granville, Russell, and north) a less common summer resident. May 5–June 3; summer; Aug. 26–Oct. 11 (Nov. 3, 19).

R. I.—Uncommon migrant. May 14–22; fall–Oct. 17.

CONN.—Common migrant and rare summer resident (Eastford, Litchfield). (April 29) May 10–summer; Sept.–Oct. 19. Eggs, June 6–13.

347. *DENDROICA CORONATA* (Linné) Gray.**Myrtle warbler; Golden-crowned warbler; Willow warbler; Yellow-rumped warbler.**

Chapman, Warblers No. Amer., 1907, p. 141, pl. 10, fig. 1–3. Egg, Capen, 1886, pl. 4, fig. 2–4.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

*Motacilla coronata* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 333.  
 "PENNSYLVANIA."

Coniferous woods, and in migration, open tree growth; nests in trees.

ME.—Common migrant and south to Scarborough a less common summer resident; rare winter resident (Cape Elizabeth). April 15–Oct. 30 (winter). Eggs, May 30–June 11.

N. H.—Common migrant and summer resident, local south of White Mts. (Mar. 5, Mar. 20) April 23–May 28; summer; Sept.–Oct. 23 (Nov. 30). Eggs, May 31–June 7.

VT.—Common migrant, and uncommon summer resident (south to Brattleboro). April 12–fall.

MASS.—Common migrant, and in higher parts of western counties an uncommon summer resident (to Mt. Wachusett); near the coast a regular local winter resident in small numbers. April 13–May 30; summer; Aug. 21–Nov. 26; winter.

R. I.—Common migrant and less common winter resident. Sept. 7–May 10.

CONN.—Common migrant and uncommon winter resident. April 8–May 20; Sept. 20–Nov. 10; winter.

348. *DENDROICA AUDUBONI* (Townsend) Baird.

**Audubon's warbler.**

Chapman, Warblers No. Amer., 1907, p. 147, pl. 10, fig. 4–6.

*Sylvia auduboni* Towns., Journ. acad. nat. sci. Phila., 1837, vol. 7, p. 191. "FORESTS OF THE COLUMBIA RIVER."

Coniferous woods; nests in trees.

MASS.—Accidental visitor: Cambridge, Nov. 15, 1876.

CONN.—Doubtfully recorded: New Haven, May 6 and 8, 1893 (A. H. Verrill).

349. *DENDROICA MAGNOLIA* (Wilson) A. O. U. comm.

**Magnolia warbler; Black and yellow warbler; Spotted warbler.**

*Sylvia magnolia* Wilson, Amer. ornith., 1811, vol. 3, p. 63, pl.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

23, fig. 2. "BANKS OF THE LITTLE MIAMI NEAR ITS JUNCTION WITH THE OHIO"; and "NOT FAR FROM FORT ADAMS ON THE MISSISSIPPI."

*Dendroica maculosa* (Gmel.). Chapman, Warblers No. Amer., 1907, p. 121, pl. 11, fig. 1-3. Egg, Capen, 1886, pl. 4, figs. 12, 14.

Coniferous and mixed thickets near streams in summer, woodlands generally in migration; nests in low trees.

ME.—Common migrant and in northern part a less common summer resident. May 6-Sept. 26. Eggs, May 30-June 16.

N. H.—Common migrant; uncommon local summer resident south of White Mts., commoner north, to 3000 ft. May 9-Oct. 3. Eggs, May 24-June 20.

VT.—Common migrant, less common local summer resident. May 14-fall.

MASS.—Common migrant and in northwestern counties a less common local summer resident. (May 4) May 7-30; summer; Aug. 25-Oct. 9. Eggs, June 15.

R. I.—Common migrant. May 14-25; Sept.-Oct. 8.

CONN.—Common migrant. May 9-June 1; Sept. 15-Oct. 1.

350. *DENDROICA CERULEA* (Wilson) Baird.**Cerulean warbler.**

Chapman, Warblers No. Amer., 1907, p. 170, pl. 9, fig. 4-6.

*Sylvia caerulea* Wils., Amer. ornith., 1810, vol. 2, p. 141, pl. 17, fig. 5. "PENNSYLVANIA."

High tree tops; nests in trees.

MASS.—Accidental visitor: Cobasset, 1874.

R. I.—Accidental visitor: Lonsdale, May 14, 1893; Pawtucket, May 22, 1879; Providence, May 22, 1878.

CONN.—Accidental visitor: Seymour, May 10, 1888; Stratford, April, 1841; Suffield, June 12, 1875.



PASSERES.

OSCINES.

## MNIOTILTIDAE.

## 351. DENDROICA PENNSYLVANICA (Linné) Baird.

**Chestnut-sided warbler.**

Chapman, Warblers No. Amer., 1907, p. 187, pl. 12, fig. 4-6.  
Egg, Capen, 1886, pl. 4, figs. 11, 12.

*Motacilla pensylvanica* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 333. "IN PENNSYLVANIA."

Brushy thickets, clearings, and sproutlands; nests in low bushes.

ME.—Common migrant and summer resident. May 6-Sept. 8.  
Eggs, June 4-July 4.

N. H.—Common migrant and summer resident, mainly at lower elevations. May 6-Sept. 25. Eggs, May 26-June 6.

VT.—Common migrant and summer resident. Eggs, June 6.

MASS.—Common migrant and summer resident. (April 28, 1908, Stoughton) May 2-Sept. 20. Eggs, May 26-June 17.

R. I.—Common migrant and summer resident. May 1-Sept. 7.  
Eggs, May 22-June 4.

CONN.—Common migrant and summer resident. May 4-Sept. 8.  
Eggs, May 23-June 15.

## 352. DENDROICA STRIATA (Forster) Baird.

**Black-poll warbler; Autumnal warbler.**

Chapman, Warblers No. Amer., 1907, p. 196, pl. 2, fig. 1-3.  
Egg, Capen, 1886, pl. 4, figs. 7, 8.

*Muscicapa striata* Forst., Phil. trans. roy. soc. London, 1772, vol. 62, pp. 406, 428. "SEVERN RIVER. . . AD SINUM HUDSONIS."

Spruce and fir forests in summer, and tree growth generally in migrations; nests in low trees.

ME.—Common migrant, and in northern part, a common summer resident. May 9-June 9; summer-Oct. 10. Eggs, June 15-early July.

N. H.—Common migrant and from White Mts. north, a com-

PASSERES.

OSCINES.

## MNIOTILTIDAE.

mon summer resident mainly above 3000 ft. May 8–Oct. 10. Eggs, June 28.

VT.— Common migrant, and on higher mountains an uncommon summer resident (Killington Peak, Mt. Mansfield, Stratton Mt.).

MASS.— Common migrant; may breed rarely (young at North Adams, August). (April 15) May 8–June 14; (Aug. 17, 1886, Concord) Sept. 1–Nov. 7.

R. I.— Common migrant. May 10–31 (June 17); Sept. 3–Oct. 23.

CONN.— Common migrant. May 11–June 5; Sept. 10–Oct. 26.

353. *DENDROICA BLACKBURNIAE* (Gmelin) Baird.

**Blackburnian warbler; Hemlock warbler.**

Chapman, Warblers No. Amer., 1907, p. 175, pl. 3, figs. 1, 2. Egg, Capen, 1886, pl. 4, figs. 5, 6.

*Motacilla blackburniae* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 977. "IN NOVEBORACO."

High coniferous and mixed woods, especially hemlock and white pine; nests in tree tops.

ME.— Rare migrant in southern counties, elsewhere an uncommon local summer resident. (May 4) May 6–Sept. 17. Eggs, June 5–15.

N. H.— Uncommon local summer resident south of White Mts., thence commoner up to 3000 ft. (May 3) May 8–Sept. 10. Eggs, June 4–15.

VT.— Uncommon local summer resident. (May 2) May 7–summer.

MASS.— Rare migrant and summer resident in eastern, commoner in northwestern part. (May 2) May 6–30; summer; Sept. 7–23 (Oct. 10). Eggs, June 5–26.

R. I.— Uncommon migrant. May 14; Sept.–Oct.

CONN.— Common migrant and rare summer resident. May 5–27; summer; Sept. 15–Oct. 3 (Nov. 8, 1907, New Haven).

PASSERES.

OSCINES.

## MNIOTILTIDAE.

354. *DENDROICA CASTANEA* (Wilson) Baird.**Bay-breasted warbler.**

Chapman, Warblers No. Amer., 1907, p. 192, pl. 12, fig. 1-3.  
Egg, Capen, 1886, pl. 4, figs. 9, 10.

*Sylvia castanea* Wils., Amer. ornith., 1810, vol. 2, p. 97, pl. 14,  
fig. 4. "PENNSYLVANIA."

Spruce and fir forests in summer, tree tops generally in migration;  
nests in trees.

ME.— Rare migrant and in northern part a rare summer resident.  
(May 8) May 14–Aug. Eggs, June 8–15.

N. H.— Uncommon migrant and from White Mts. north an  
uncommon summer resident from 2–4000 ft. May 14–Sept.  
Eggs, June 3.

VT.— Rare migrant.

MASS.— Usually rare, but sometimes common migrant. May 8–  
May 28 (June 19; July 2, Springfield); Aug. 23–Oct. 8.

R. I.— Uncommon migrant. May 9– ; Sept.

CONN.— Migrant, irregularly rare or common. May 15–30;  
Sept. 30–Oct. 17.

355. *DENDROICA DOMINICA* (Linné) Baird.**Yellow-throated warbler.**

Chapman, Warblers No. Amer., 1907, p. 180, pl. 13, figs. 1, 2.

*Motacilla dominica* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 334.  
"JAMAICA, DOMINICA."

Tree growth bordering streams or bayous; nests in trees.

VT.— Recorded from Lunenburg, 1878.

MASS.— Accidental visitor: Dedham, Nov. 4, 1866; Springfield,  
May, 1892.

CONN.— Accidental visitor: Connecticut valley, "several";  
New Haven.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

356. *DENDROICA VIRENS* (Gmelin) Baird.**Black-throated green warbler.**

Chapman, Warblers No. Amer., 1907, p. 157, pl. 14, fig. 1-3.  
Egg, Capen, 1886, pl. 3, fig. 15.

*Motacilla virens* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 985.  
"PENSILVANIA."

Forest trees, particularly white pines; nests in coniferous trees (once "on the ground," Brewster).

ME.—Common migrant and summer resident. April 28–Oct. 13. Eggs, May 30–July 1.

N. H.—Common migrant and summer resident mainly below 3000 ft. (April 26) May 1–Oct. 25. Eggs, June 2.

VT.—Common migrant and summer resident. May 2–fall.

MASS.—Common migrant and summer resident. (April 20) May 1–Oct. 19 (Nov. 3). Eggs, May 30–June 18.

R. I.—Common migrant and summer resident. (April 25) May 1–Oct. 13. Eggs, June 10.

CONN.—Common migrant and uncommon local summer resident. (April 27) May 1–23; summer; Sept. 15–Oct. 20. Eggs, May 21–June 18.

357. *DENDROICA VIGORSII* (Audubon) Stejneger.**Pine warbler; Pine-creeping warbler.**

*Sylvia vigorsii* Aud., Ornith. biogr., 1832, vol. 1, p. 153, pl. 30.  
"A SMALL ISLAND OF PERKIOMING CREEK, . . . PENNSYLVANIA."

*Sylvicola pinus* Lath. Audubon, Birds of Amer., 1841, vol. 2, p. 37, pl. 82. Egg, Capen, 1886, pl. 5, figs. 2, 3.

Coniferous woods, especially pitch-pine groves; nests in trees.

ME.—Common migrant and uncommon local summer resident mainly in southern counties. April 5–Oct. 21. Eggs, June–July 6.

N. H.—Uncommon local summer resident north to White Mts. April 6–Sept. 30.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

Vt.— Uncommon local summer resident in southern part. April 17–fall.

Mass.— Common migrant and summer resident; accidental in winter. (Mar. 29) April 1–Nov. 25 (winter). Eggs, May 8–June 16; July (2d brood).

R. I.— Common summer resident of pitch-pine groves. April 9–Oct. 29 (Nov. 15). Eggs, May 25.

Conn.— Common migrant and summer resident. April 14–Sept. 29.

358. *DENDROICA PALMARUM* (Gmelin) Baird.**Palm warbler; Redpoll warbler.**

Chapman, Warblers No. Amer., 1907, p. 213, pl. 16, figs. 4, 5.

*Motacilla palmarum* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 951.

“IN INSULA S. DOMINICI.”

Open tree or bushy growth, and edges of fields; nests on the ground.

N. H.— Uncommon fall migrant (a doubtful spring record, Manchester, May 9, 1900). Sept. 7–16.

Vt.— Rare fall migrant.

Mass.— Rare fall migrant. Sept. 7–Oct. 28 (Dec. 6, 1902, Belmont).

R. I.— Rare fall migrant: Westerly, Sept. 21, 1896.

Conn.— Rare fall migrant; once recorded in winter (Jan. 1880, at New Haven). Sept. 19–Oct. 10 (Jan.).

359. *DENDROICA PALMARUM HYPOCHRYSEA* Ridgway.**Yellow palm warbler; Twitch-tail warbler; Yellow redpoll warbler.**

Ridgw., Bull. Nuttall ornith. club, 1876, vol. 1, p. 85. “CAMBRIDGE, MASS.”

*Sylvicola petechia* Lath. Audubon, Birds of Amer., 1841, vol. 2, p. 55, pl. 90.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

Swamps, edges of woods, or open country; nests on or near the ground.

ME.—Common spring and uncommon fall migrant and local summer resident. April 11–May 20; summer; Sept. 17–Oct. 20 (Oct. 27, Nov. 2). Eggs, May 27–June 26.

N. H.—Common spring and uncommon fall migrant. April 13–May 11; Sept. 16–Oct. 9.

VT.—Uncommon migrant.

MASS.—Common spring and uncommon fall migrant; said to have wintered. (Mar. 28, 1908, Hyde Park; April 5) April 13–May 21; Sept. 5–Oct. 26.

R. I.—Common spring and uncommon fall migrant. April 5–22; Sept. 15–26.

CONN.—Common spring and uncommon fall migrant. April 9–May 5; Oct. 18–Nov. 4.

360. *DENDROICA DISCOLOR* (Vieillot) Baird.

**Prairie warbler.**

Chapman, Handb. birds east. No. Amer., 1895, p. 364. Egg, Capen, 1886, pl. 5, fig. 1.

*Sylvia discolor* Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 2, p. 37, pl. 98. “ÉTATS-UNIS ET LES GRANDS ÎLES ANTILLES.”

*Sylvicola discolor* (Vieill.). Audubon, Birds of Amer., 1841, vol. 2, p. 68, pl. 97.

Dry bush-grown fields and sproutlands; nests in bushes.

N. H.—Rare local summer resident in southern part (Hollis, Manchester).

VT.—Accidental visitor: Uxbridge, Dec. 4, 1884.

MASS.—Uncommon summer resident, mainly near coast; accidental in winter (South Yarmouth, Jan. 2, 1909). May 2–Sept. 25 (Jan. 2). Eggs, May 28–June 18.

R. I.—Common local summer resident. May 4–Sept. (Dec. 4, 1884, Providence). Eggs, May 27–June 14.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

CONN.—Uncommon summer resident. (April 27) May 1–Sept. 15. Eggs, May 27–June 25.

361. *SEIURUS AUROCAPILLUS* (Linné) Swainson.

**Oven-bird; Golden-crowned thrush.**

Chapman, Handb. birds east. No. Amer., 1895, p. 365; Allen, Auk, 1894, vol. 11, pl. 2. Egg, Capen, 1886, pl. 5, figs. 4, 5.

*Motacilla aurocapilla* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 334. "IN PENNSYLVANIA."

Forests and groves; nests on the ground.

ME.—Common migrant and summer resident. (April 23) May 5–Sept. 28. Eggs, May 30–June 10 (July 29).

N. H.—Common migrant and summer resident below 3000 ft. May 3–Oct. 2. Eggs, June 2–July 5.

VT.—Common migrant and summer resident. May 5–fall. Eggs, June 7.

MASS.—Common migrant and summer resident. (April 30) May 2–Sept. 27 (Dec. 30, 1881, Watertown). Eggs, May 17–June 16 (July 8).

R. I.—Common migrant and summer resident. May 2–Sept. 28. Eggs, May 19–30.

CONN.—Common migrant and summer resident (April 26) May 3–Sept. Eggs, May 20–July 10.

362. *SEIURUS NOVEBORACENSIS* (Gmelin) Bonaparte.

**Water-thrush; New York Accentor; Northern water-thrush; Wagtail warbler; Water wagtail.**

Ridgway, Bull. 50, U. S. nat. mus., 1902, pt. 2, p. 642. Egg, Capen, 1886, pl. 5, figs. 6, 7.

*Motacilla noveboracensis* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 958. "IN LOUISIANA, & NOVEBORACI SEPIBUS."

*Turdus aquaticus* Wils. Audubon, Ornith. biogr., 1839, vol. 5, p. 284, pl. 433, fig. 7.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

Swampy woods and borders of streams; nests on the ground or among tree roots.

ME.— Common migrant and in northern part a less common local summer resident. (April 28) May 6–Sept. 25. Eggs, May 28–June 23.

N. H.— Common migrant and less common summer resident, local at lower elevations south of White Mts. (April 30) May 3–Sept. 26.

Vt.— Common migrant, and mainly in northern part, a less common summer resident.

MASS.— Common migrant and rare local summer resident (near Boston, Lancaster). (April 25) May 2–June 5; summer; Aug. 1–Oct. 16. Eggs, May 21.

R. I.— Common migrant and rare local summer resident (Washington Co.). May 7–(summer); Aug. 12–Sept. 15. Eggs, May 19, 20.

CONN.— Common migrant. (May 1) May 5–27; Aug.–Oct. 18.

363. *SEIURUS NOVEBORACENSIS NOTABILIS* (Ridgway) Ridgway.

**Grinnell's water-thrush.**

*Siurus naevius notabilis* Ridgw., Proc. U. S. nat. mus., 1880, vol. 3, p. 12. "BLACK HILLS, WYOMING."

Vicinity of streams; nests on or near the ground.

CONN.— Rare migrant: East Haven, Sept. 21, 1904; Hamden, Sept. 7, 1904; Orange, May 20, 1905 (Bishop).

364. *SEIURUS MOTACILLA* (Vieillot) Bonaparte.

**Louisiana water-thrush; Large-billed water-thrush.**

Ridgway, Bull. 50, U. S. nat. mus., 1902, pt. 2, p. 639.

*Turdus motacilla* Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 2, p. 9, pl. 65. "KENTUCKY."

*Turdus ludovicianus* Audubon, Ornith. biogr., 1831, vol. 1, p. 99, pl. 19. Egg, Capen, 1886, pl. 5, fig. 8.



PASSERES.

OSCINES.

## MNIOTILTIDAE.

Wooded borders of streams; nests on the ground or among tree roots.

ME.— Accidental visitor: Norway, May, 1865 (*see* Waterville, 1865).

N. H.— Doubtfully recorded (Dublin).

VT.— Doubtfully recorded (Brattleboro, "fall").

MASS.— Rare visitor and summer resident (Berkshire Co. and Connecticut valley). April 13–July 28.

R. I.— Rare summer resident. May 2. Eggs, May 15–28.

CONN.— Uncommon summer resident in southern part. (Feb. 15, 1882, ? Saybrook) April 13–July. Eggs, May 6–June 10.

## 365. OPORORNIS FORMOSA (Wilson) Baird.

**Kentucky warbler.**

Chapman, Warblers No. Amer., 1907, p. 235, pl. 18, figs. 1, 2.

*Sylvia formosa* Wils., Amer. ornith., 1811, vol. 3, p. 85, pl. 25, fig. 3. "KENTUCKY AND TENNESSEE."

Undergrowth of swampy woods; nests on or near the ground.

VT.— Accidental visitor: Lunenburg, May 30, 1905; also a former doubtful record (Taftsville, May 19, 1884).

MASS.— Accidental visitor: Wellesley, May 14, 1907; also doubtfully recorded from Lynn.

R. I.— Accidental visitor: Newport, spring of 1890.

CONN.— Rare summer visitor (Lyme; Suffield, Aug. 16, 1876; West Stratford, May 30, 1888) and summer resident (Greenwich, with nestlings, July 10, 1892).

## 366. OPORORNIS AGILIS (Wilson) Baird.

**Connecticut warbler.**

Chapman, Warblers No. Amer., 1907, p. 241, pl. 18, figs. 3, 4.

*Sylvia agilis* Wils., Amer. ornith., 1812, vol. 5, p. 64, pl. 39, fig. 4. "CONNECTICUT."

PASSERES.

OSCINES.

## MNIOTILTIDAE.

Tamarack swamps, and in migration, damp woods and thickets; nests on the ground.

ME.— Rare fall migrant. Sept. 5–20 (*nee* Aug. 30, 1878, Cape Elizabeth).

N. H.— Rare fall migrant. Sept. 6–Oct. 11.

VT.— Rare (? spring and) fall migrant. (? Brattleboro, May; ? Rutland, April 24, 1879); Sept. 20–30.

MASS.— Uncommon fall migrant (? accidental in spring). (May 1890, West Newton, *vide* Maynard); Sept. 7–Oct. 9.

R. I.— Uncommon fall migrant. Sept. 24–Oct. 2 (Nov. 12).

CONN.— Accidental spring and uncommon fall migrant. (Spring, Saybrook; *nee* New Haven); Sept. 15–Oct. 10.

367. OPORORNIS PHILADELPHIA (Wilson) Ridgway.

**Mourning warbler.**

*Sylvia philadelphia* Wils., Amer. ornith., 1810, vol. 2, p. 101, pl. 14, fig. 6. "WITHIN A FEW MILES OF PHILADELPHIA."

*Trichas philadelphia* (Wils.). De Kay, Zool. New York, 1844, vol. 2, p. 81, pl. 54, fig. 122. Egg, Capen, 1886, pl. 5, figs. 12, 13.

Tangles of bushes, vines, and briars; nests on or near the ground.

ME.— Uncommon migrant, and in northern part an uncommon summer resident. (May 16) May 24–fall. Eggs, June 10–16.

N. H.— Uncommon migrant and from White Mts. north an uncommon summer resident. May 19–Sept. 13. Eggs, June 8.

VT.— Uncommon migrant and summer resident south to Londonderry. May 20–Sept. 20.

MASS.— Rare migrant and in Berkshire Co., a rare summer resident. (May 14) May 21–June 13; summer; Sept. 12–30.

R. I.— Rare migrant. May 21–June 5; Sept. 4.

CONN.— Rare migrant. May 12–27; fall.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

368. *GEOTILYPIS TRICHAS* (Linné) Cabanis.

**Maryland yellowthroat; Black-masked ground warbler; Yellowthroat.**

*Turdus trichas* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 293.  
"IN AMERICA SEPTENTRIONALI."

*Geothlypis trichas brachidactyla* (Swains.). Chapman, Warblers No. Amer., 1907, p. 251, pl. 20, fig. 3-5. Egg, Capen, 1886, pl. 5, fig. 9-11.

Weed patches and thickets near water; nests on or near the ground.

ME.—Common migrant and summer resident. May 6–Oct. 13 (Oct. 22). Eggs, May 28–June 12.

N. H.—Common migrant and summer resident at lower elevations. May 6–Oct. 7. Eggs, June 3–21.

VT.—Common migrant and summer resident. Eggs, June 2.

MASS.—Common migrant and summer resident; accidental in winter. (April 30) May 2–Oct. 21 (winter). Eggs, May 24–June 17.

R. I.—Common migrant and summer resident. (April 22)–Oct. 17. Eggs, May 25–June 5.

CONN.—Common migrant and summer resident. (April 29) May 4–Nov. 1. Eggs, May 22–June 18.

369. *ICTERIA VIRENS* (Linné) Baird.

**Yellow-breasted chat; Ghost-bird (Indian); Yellow chat.**

Audubon, Birds of Amer., 1842, vol. 4, p. 162, pl. 244. Egg, Capen, 1886, pl. 5, figs. 14, 15.

*Turdus virens* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 171.  
"AMERICA."

Bushy thickets; nests in low bushes.

ME.—Accidental visitor (five instances). June 6–Oct. 5.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

N. H.—Doubtfully recorded.

VT.—Very rare summer resident in extreme south: Pownall, June 16, 1896.

MASS.—Rare local summer resident near coast and in western valleys. May 10–Sept. 10 (Oct. 26, 1907, Lynnfield). Eggs, May 18–June 18.

R. I.—Locally common summer resident in southern part. May 3–Aug. 5. Eggs, May 31–June 13.

CONN.—Locally common summer resident. May 5–Aug. Eggs, May 22–July 7.

370. *WILSONIA CITRINA* (Boddaert) A. O. U. comm.

**Hooded warbler.**

*Muscicapa citrina* Bodd., Tabl. d. planch. enlum. d'hist. nat., 1783, p. 41. Based on Buffon.

*Wilsonia mitrata* (Gmel.). Chapman, Warblers No. Amer., 1907, p. 269, pl. 21, fig. 1–3. Egg, Capen, 1886, pl. 6, fig. 1.

Undergrowth of woods; nests in bushes.

ME.—Accidental visitor: Falmouth, Sept. 10, 1904.

VT.—Doubtfully recorded.

MASS.—Rare visitor; possibly breeding in southwestern valleys. May 8–Oct. 15 (June 25, 1879, Brookline; June 25, 1888, Provincetown).

R. I.—Accidental visitor: Kingston.

CONN.—Uncommon summer resident of laurel undergrowth. May 2–Sept. Eggs, May 25–June 24.

371. *WILSONIA PUSILLA* (Wilson) Bonaparte.

**Wilson's warbler; Black-capped flycatching warbler; Wilson's blackcap.**

Chapman, Warblers No. Amer., 1907, p. 274, pl. 22, figs. 1, 2. Egg, Capen, 1886, pl. 6, fig. 2.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

*Muscicapa pusilla* Wils., Amer. ornith., 1811, vol. 3, p. 103, pl. 26, fig. 4. "SOUTHERN STATES, AND . . . NEW JERSEY AND DELAWARE."

Bushy growth near streams or in swamps; nests on the ground.

ME.—Uncommon migrant and summer resident (south to Pittsfield). May 7–June 5; summer; Aug. 18–Sept. 11 (Oct.). Eggs, May 31–June 19.

N. H.—Uncommon migrant and north of White Mts. a rare local summer resident (Lancaster). May 9–29; (summer); Aug. 15–Oct. 3. Eggs, June 6.

VT.—Uncommon migrant; (? summer). May 17; Aug. 24.

MASS.—Uncommon migrant. May 7–30 (June 1); Aug. 23–Sept. 27 (Nov. 20, 1898, Belmont).

R. I.—Uncommon migrant. May 15–22; Sept. (Nov. 30).

CONN.—Uncommon migrant. May 10–25; Sept. 27.

372. *WILSONIA CANADENSIS* (Linné) Coues.**Canadian warbler; Canada flycatcher.**

Chapman, Warblers No. Amer., 1907, p. 280, pl. 22, fig. 3–5. Egg, Capen, 1886, pl. 6, figs. 3, 4.

*Muscicapa canadensis* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 327. "CANADA."

Wet woods or thickets along streams; nests on or near the ground.

ME.—Common migrant and summer resident. (April 25, Bangor; May 7) May 10–Sept. 12. Eggs, May 30.

N. H.—Common migrant and less common local summer resident below 3000 ft. May 13–Sept. 13. Eggs, June 9–13.

VT.—Common migrant and less common local summer resident. May 15–fall.

MASS.—Common migrant and less common summer resident mainly in western part. May 9–Sept. 29. Eggs, June 9–26.

R. I.—Common migrant and uncommon local summer resident. May 11–fall.

PASSERES.

OSCINES.

## MNIOTILTIDAE.

CONN.—Common migrant and rare local summer resident (Portland). (May 6) May 12–fall. Eggs, May 30.

373. *SETOPHAGA RUTICILLA* (Linné) Swainson.

**American redstart.**

Chapman, Warblers No. Amer., 1907, p. 287, pl. 23, fig. 1–3. Egg, Capen, 1886, pl. 6, fig. 5.

*Motacilla ruticilla* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 186. "AMERICA."

Orchards, shade and forest trees; nests in trees.

ME.—Common migrant and summer resident. May 4–Oct. 1 (Oct. 13). Eggs, June 2–30.

N. H.—Common migrant and summer resident below 3000 ft. May 4–Oct. 5. Eggs, May 30–June 13.

VT.—Common migrant and summer resident. May 8–fall.

MASS.—Common migrant and summer resident. (April 13, 23, 30) May 4–Oct. 12 (Nov. 1, 1908, Arnold arboretum). Eggs, May 29–June 21.

R. I.—Common migrant and summer resident. May 1–Sept. 8. Eggs, June 1–13.

CONN.—Common migrant and summer resident. May 3–Oct. 16 (Oct. 26). Eggs, May 20–June 27.

## MOTACILLIDAE.

374. *ANTHUS RUBESCENS* (Tunstall) A. O. U. comm.

**American pipit; Brown lark; Titlark; Wagtail.**

*Alauda rubescens* Tunstall, Ornith. Britannica, 1771, p. 2. No locality; based on Red lark from London and North America (*vide* Oberholser).

*Anthus pensilvanicus* (Lath.). Ridgway, Bull. 50, U. S. nat. mus., 1904, pt. 3, p. 12.

*Anthus pipiens* Audubon, Ornith. biogr., 1832, vol. 1, pl. 80.

## PASSERES.

## OSCINES.

## MOTACILLIDAE.

Barren grounds, fields and sea coasts; nests on the ground.

ME.— Rare spring and common fall migrant. May 2–18 (June 1, 1907, Lubec); Sept. 13–Oct. 25 (Dec. 13, 1901, Seguin Id.).

N. H.— Rare spring and common fall migrant. April 3–May 10; Sept. 15–Nov.

VT.— Rare spring and uncommon fall migrant.

MASS.— Uncommon spring and common fall migrant; accidental in winter (Boston Harbor, Feb. 25, 1882; Newburyport, Jan. 4, 1878). April 18–May 20 (June 8); (Aug. 29, 1888, Monomoy Id.) Sept. 10–Nov. 24 (winter).

R. I.— Rare spring and common fall migrant. Mar. 29–May 15; Sept. 16–Nov. 28.

CONN.— Rare spring and common fall migrant. April–May; Sept. 15–Nov. 14 (Dec. 25, 1908, South Norwalk).

## MIMIDAE.

375. *MIMUS POLYGLOTTOS* (Linné) Bonaparte.

**Mockingbird.**

Chapman, Handb. birds east. No. Amer., 1895, p. 377; Bird-life, 1898, pl. 66. Egg, Capén, 1886, pl. 1, fig. 8–10.

*Turdus polyglottos* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 169. "VIRGINIA."

Thickets and dense tree clumps; nests in bushes or trees.

ME.— Occasional visitor at all seasons or escape; said to have bred at Leeds about 1885.

N. H.— Accidental visitor or escape: Hampton; Milford.

VT.— Accidental visitor or escape; pair nested at Lunenburg, 1884.

MASS.— Occasional visitor at all seasons or escape; rare summer resident (Arlington, Groton, Marshfield, ? Provincetown, Springfield). Eggs, June 13–20.

R. I.— Occasional visitor at all seasons (or escape).

## PASSERES.

## OSCINES.

## MIMIDAE.

CONN.—Rare summer resident (more frequent formerly); occasional visitor (or escape) at all seasons. Eggs, June 20 (Jewett City).

376. *DUMETELLA CAROLINENSIS* (Linné) A. O. U. comm.

**Catbird; Pussy.**

*Muscicapa carolinensis* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 328. "CAROLINA."

*Orpheus carolinensis* (Linné). Audubon, Birds of Amer., 1841, vol. 2, p. 195, pl. 140. Egg, Capen, 1886, pl. 1, fig. 11.

Bushy thickets near water; nests in bushes or among vines.

ME.—Common summer resident, mainly of southern counties. May 17–Oct. 4. Eggs, June 2–17; July (2d brood).

N. H.—Common summer resident north to White Mt. valleys; accidental in early winter (Concord, Dec. 3, 1901; Wilton, Dec. 25, 1904). May 6–Oct. 14 (Dec. 25). Eggs, June 6–July 28 (? 2d brood).

VT.—Common summer resident.

MASS.—Common summer resident; accidental in winter. (April 13, 1908, Westwood) April 29–Oct. 16 (Dec.; winter 1908–09, Forest Hills). Eggs, May 22–June 20.

R. I.—Common summer and casual winter resident. (April 24)–Oct. 13 (Dec. 28–Mar. 4, Mar. 19). Eggs, May 25–June 10.

CONN.—Common summer resident. May 3–Nov. 4. Eggs, May 17–23; July 26 (2d brood).

377. *TOXOSTOMA RUFUM* (Linné) Cabanis.

**Brown thrasher; Brown thrush; Mavis; Mocking thrush; Planting-bird; Song thrush; Thrasher.**

*Turdus rufus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 169. "IN AMERICA SEPTENTRIONALI & MERIDIONALI."

*Orpheus rufus* (Linné). Audubon. Birds of Amer., 1841, vol. 3, p. 9, pl. 141. Egg, Capen, 1886, pl. 1, figs. 12, 13.



PASSERES.

OSCINES.

## MIMIDAE.

Thickets and sproutlands; nests in bushes or rarely on the ground.

ME.— Common summer resident in southwestern counties. (? April 8) April 28–Oct. 14.

N. H.— Uncommon summer resident north to White Mt. valleys. (April 16)–Oct. 10.

VT.— Uncommon summer resident at lower elevations.

MASS.— Common summer and accidental winter resident (Arlington). (April 16) April 20–Oct. 26 (Nov. 4; winter). Eggs, May 10–June 21.

R. I.— Common summer resident; casual in winter. (April 14)–Oct. 13 (Jan. 30). Eggs, May 19–June 4.

CONN.— Common summer resident. April 22–Oct. 2.

## TROGLODYTIDAE.

378. *THRYOTHORUS LUDOVICIANUS* (Latham) Bonaparte.

**Carolina wren; Great Carolina wren; Mocking wren.**

*Sylvia ludoviciana* Lath., Index ornith., 1790, vol. 2, p. 548. "LOUISIANA."

*Troglodytes ludovicianus* Bonap. Audubon, Birds of Amer., 1841, vol. 2, p. 116, pl. 117. Egg, Capen, 1886, pl. 2, fig. 11.

Dense undergrowth and tangles; nests in cavities of trees or crevices of buildings.

ME.— Accidental visitor (Falmouth, Oct. 3, 1908; Waterville) and accidental summer resident (Norway Lake, eggs, June, 1883, *vide* Knight).

N. H.— Accidental visitor: Alstead, July 6, 1903; Rye Beach, Aug. 7, 1880; Webster, about 1900.

VT.— Doubtfully recorded.

MASS.— Occasional visitor at all seasons and very rare summer resident (Naushon Id., July, 1901).

PASSERES.

OSCINES.

## TROGLODYTIDAE.

R. I.— Occasional visitor in summer and fall, and rare summer resident (at Middletown). April 9–Nov. 28. Eggs, April 15.

CONN.— Occasional visitor at all seasons and rare resident (near Bridgeport, Chester, South Norwalk). Eggs, July 16.

379. *THRYOMANES BEWICKII* (Audubon) Ridgway.**Bewick's wren.**

Ridgway, Bull. 50, U. S. nat. mus., 1904, pt. 3, p. 552.

*Troglodytes bewickii* Aud., Ornith. biogr., 1831, vol. 1, p. 96, pl. 18. "ABOUT FIVE MILES FROM ST. FRANCISVILLE, . . . LOUISIANA."

Brush piles, orchards, and vicinity of outbuildings; nests in cavities of trees or buildings.

ME.— Doubtfully recorded (Seguin Id., Oct. 6).

N. H.— Accidental visitor: Alton, April 25, 1890.

380. *TROGLODYTES AEDON* Vieillot.**House wren.**

Vieill., Hist. nat. ois. Amér. Sept., 1807, vol. 2, p. 52, pl. 107. "CANADA, . . . NOUVELLE-ÉCOSSE." Audubon, Birds of Amer., 1841, vol. 2, p. 125, pl. 120. Egg, Capen, 1886, pl. 2, figs. 12, 13.

Brush piles, orchards, and vicinity of outbuildings; nests in hollow limbs, bird boxes, or any convenient crevice.

ME.— Uncommon local summer resident (to Caribou). May 8–Aug.

N. H.— Uncommon summer resident becoming rare among White Mt. valleys. May 18–Sept. 24.

VT.— Uncommon summer resident. May 11–fall.

MASS.— Locally common summer resident. (April 24) May 1–Oct. 30. Eggs, May 25–June 5.

R. I.— Locally common summer resident; formerly more common. April 26–Sept. Eggs, May 21–27.

CONN.— Common summer resident. April 25–Oct. 30.

PASSERES.

OSCINES.

## TROGLODYTIDAE.

381. *NANNUS HIEMALIS* (Vieillot) A. O. U. comm.**Winter wren.**

*Troglodytes hiemalis* Vieill., Nouv. dict. d'hist. nat., 1819, vol. 34, p. 514. "NOUVELLE-ÉCOSSE ET DANS L'ÉTAT DE NEW YORK." Audubon, Birds of Amer., 1841, vol. 2, p. 128, pl. 121. Egg, Capen, 1886, pl. 2, fig. 14.

Damp woods or thickets near water, and in migration, tangles, brush heaps, and brier patches; nests in crevices of upturned roots.

ME.— Common migrant and in northern part a common summer resident; very rare winter resident (Bangor). (April 1) April 9–Nov. 17 (winter). Eggs, May 20–June 3; Aug. 8 (2d brood).

N. H.— Common migrant and less common summer resident, local south of White Mts.; rare winter resident in southeast part. April 5–Nov. 15 (winter). Eggs, May 20–July 23 (2d brood).

VT.— Uncommon migrant, less common summer resident in mountainous parts south to Londonderry; may winter. April 7–Nov. 24 (Dec. 30, Jan. 1). Eggs, June 4.

MASS.— Uncommon migrant, less common summer resident (Berkshire Co., and locally east, Lynn, Winchendon); rare winter resident near coast. April 4–29; summer; Sept. 2–Nov. 25; winter.

R. I.— Uncommon migrant and rare winter resident; rare local summer resident (Washington Co., nest with young, May 24, 1908). April; (summer); Sept.–Nov. 28; winter.

CONN.— Uncommon migrant and less common winter resident. Oct. 2–April.

382. *CISTOTHORUS STELLARIS* (Lichtenstein) Cabanis.**Short-billed marsh wren.**

*Troglodytes stellaris* Licht., in Naumann's Vögel Deutschl., 1823, vol. 3, tab. ad p. 724. "CAROLINA."

PASSERES.

OSCINES.

## TROGLODYTIDAE.

*Troglodytes brevirostris* Nutt. Audubon, Birds of Amer., 1841, vol. 2, p. 138, pl. 124. Egg, Capen, 1886, pl. 3, fig. 1.

Wet grassy meadows; nest usually built among tall grass stems near the ground.

ME.—Rare local summer resident (Bangor, Glenburn). May 4–Oct. 10.

N. H.—Rare local summer resident in southern part; rare visitor in White Mt. region.

MASS.—Very local summer resident, sometimes common. May 6–Oct. 9. Eggs, May 25–June 18; July 7 (? 2d brood).

R. I.—Rare local summer resident (Newport). Eggs, June 6.

CONN.—Rare local summer resident. Spring–Sept. 17.

## 383. TELMATODYTES PALUSTRIS (Wilson) Henry.

**Long-billed marsh wren.**

Ridgway, Bull. 50, U. S. nat. mus., 1904, pt. 3, p. 489. Egg, Capen, 1886, pl. 2, fig. 15.

*Certhia palustris* Wils., Amer. ornith., 1810, vol. 2, p. 58, pl. 12, fig. 4. "PENNSYLVANIA...ALONG THE TIDE WATER OF OUR RIVERS."

Cat-tail swamps; nests among the flags.

VT.—Rare visitor: Brattleboro, May, 1902; Middlebury; Rutland, May 27, 1907.

MASS.—Common local summer resident in western valleys and coastwise (Essex Co.); rare winter resident (Fresh Pond). (May 2) May 15–Nov. 30 (winter). Eggs, June 5–29 (Aug. 1).

R. I.—Common local summer resident. May–Oct. 6. Eggs, May 31–June 16.

CONN.—Common summer and rare winter resident. May 5–(winter). Eggs, June 7–July 28

PASSERES.

OSCINES.

## CERTHIIDAE.

384. *CERTHIA FAMILIARIS AMERICANA* (Bonaparte) Ridgway.**Brown creeper; Tree creeper.**

*Certhia americana* Bonap., Geogr. and comparative list birds Europe and No. Amer., 1838, p. 11. "WESTERN AND NORTHERN PARTS" OF NO. AMERICA.

*Certhia familiaris* Linné. Audubon, Birds of Amer., 1841, p. 109, pl. 115. Egg, Capen, 1886, pl. 2, fig. 10.

Cool mixed or coniferous woods in summer, tree growth generally at other seasons; nests behind loose flake of bark.

ME.—Common migrant, less common summer and winter resident. Eggs, May 31–June 23.

N. H.—Resident, breeding locally south of White Mts., and more commonly north. Eggs, May.

VT.—Common migrant and winter resident; uncommon local summer resident.

MASS.—Common migrant and winter resident; less common summer resident in western part, but rare and local in eastern swamps (to Taunton). Sept. 12–May 8; summer. Eggs, May 6–23.

R. I.—Common migrant and winter resident. Sept. 2–April 26.

CONN.—Common migrant and winter resident; rare summer resident. Late Sept.–late April (summer).

## SITTIDAE.

385. *SITTA CAROLINENSIS* Latham.**White-breasted nuthatch; Sapsucker.**

Lath., Index ornith., 1790, vol. 1, p. 262. "IN AMERICA, JAMAICA; EUROPAEA MINOR." Audubon, Birds of Amer., 1842, vol. 4, p. 175, pl. 247. Egg, Capen, 1886, pl. 2, fig. 8.

Orchards, shade and forest trees; nests in holes in trees or stumps.

PASSERES.

OSCINES.

## SITTIDAE.

ME., N. H., VT.—Common migrant, uncommon summer and winter resident. Eggs, May 1–15 (Me.).

MASS.—Common migrant and winter resident; uncommon summer resident. Eggs, April 3–May 1.

R. I.—Common migrant and winter resident; uncommon summer resident. Sept. 14–April 22; summer. Eggs, April 20–May 22.

CONN.—Common migrant and winter resident, less common summer resident. Eggs, May 8–June 7.

386. *SITTA CANADENSIS* Linné.

**Red-breasted nuthatch; Canada nuthatch; Red-bellied nuthatch.**

Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 177. "CANADA." Audubon, Birds of Amer., 1842, vol. 4, p. 179, pl. 248. Egg, Capen, 1886, pl. 2, fig. 9.

Coniferous woods, or, on migrations, tree growth generally; nests in holes excavated in dead trees.

ME.—Common resident south to Cape Elizabeth; elsewhere a periodically common winter resident. Eggs, May 6–June 2 (early July).

N. H.—Common resident from White Mts. north, elsewhere a rare summer and irregular winter resident. Eggs, May 20.

VT.—Uncommon winter visitor and local summer resident. Eggs, May 5–30.

MASS.—Irregular fall and winter visitor; in western part at higher levels an uncommon, and elsewhere a rare and irregular summer resident. Aug. 15–May 20; summer. Eggs, May 18–June 10.

R. I.—Uncommon and irregular fall and winter visitor. Oct. 5–winter.

CONN.—Uncommon and irregular fall and winter visitor. (Aug. 28) Sept. 24–April 19.

PASSERES.

OSCINES.

## PARIDAE.

387. *BAEOLOPIUS BICOLOR* (Linné) Cabanis.**Tufted titmouse.**

*Parus bicolor* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 340.  
 "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer.,  
 1841, vol. 2, p. 143, pl. 125.

Woodlands generally; nests in holes of trees and stumps.

ME.—Accidental visitor: near Orono.

N. H., VT.—Doubtfully recorded.

CONN.—Rare visitor: near Hartford; Lyme, Feb. 27, 1872,  
 Jan. 1874; New Haven (Linsley).

388. *PENTHESTES ATRICAPILLUS* (Linné) Ridgway.**Chickadee; Black-capped chickadee or titmouse.**

*Parus atricapillus* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 341.  
 "CANADA." Audubon, Birds of Amer., 1841, vol. 2, p. 146, pl.  
 126. Egg, Capen, 1886, pl. 2, figs. 4, 5.

Tree growth generally; nests near water in low dead stumps.

ME.—Common resident. Eggs, May 10–June 18.

N. H.—Common resident, breeding below 3000 ft.

VT.—Common resident. Eggs, May 10.

MASS.—Common winter and less common summer resident.  
 Eggs, May 4–June 25.

R. I.—Common winter and less common summer resident.  
 Eggs, May 9–18.

CONN.—Common resident. Eggs, May 10–June 4.

389. *PENTHESTES HUDSONICUS* (Forster) Ridgway.**Hudsonian chickadee.**

*Parus hudsonicus* Forst., Phil. trans. roy. soc. London, 1772,  
 vol. 62, pp. 408, 430. "SEVERN RIVER. . . . AD SINUM HUDSONIS."  
 Audubon, Birds of Amer., 1841, vol. 2, p. 155, pl. 128. Egg,  
 Capen, 1886, pl. 2, figs. 6, 7.

PASSERES.

OSCINES.

## PARIDAE.

Spruce and fir forests; nests in holes in low trees or stumps.

ME.—Rare resident in northern and eastern counties; elsewhere an uncommon or rare winter visitor.

N. H.—Uncommon resident mainly above 3000 ft. on White Mts., north; elsewhere a rare winter visitor.

VT.—Uncommon resident in northern part and on higher mountains (Mt. Stratton, Sutton); elsewhere a rare winter visitor.

MASS.—Rare late fall and winter visitor; once in Plymouth Co., June 5, 1896. Oct. 18–April 5 (June 5).

R. I.—Accidental late fall visitor: Smithfield, Nov. 1, 1880.

CONN.—Accidental late fall visitor: New Haven, Nov. 13, 1875.

## SYLVIIDAE.

390. *REGULUS SATRAPA* Lichtenstein.

**Golden-crowned kinglet; Golden-crested wren.**

Licht., Verzeichn. d. doubl. d. Mus. Berlin, 1823, p. 35. "AM[ERICA] SEPT[ENTRIONALI]." Audubon, Birds of Amer., 1841, vol. 2, p. 165, pl. 132. Egg, Capen, 1886, pl. 2, figs. 1, 2.

Spruce and fir forests in summer, tree growth generally at other times; nests in conifers.

ME.—Common migrant and less common winter resident; common summer resident in northern part. Eggs, June 15–24.

N. H.—Common migrant, less common winter resident; common summer resident from White Mts. north, local to the south. Eggs, May 24.

VT.—Common migrant and winter resident; less common local summer resident. Sept. 18–May 7; summer.

MASS.—Common migrant and winter resident; uncommon summer resident in western part but rare and local in eastern (Lynn, Plymouth Co., Winchendon). Sept.–April; summer. Eggs, May–June 29.

R. I.—Common migrant and winter resident. Sept. 25–April 2.



PASSERES.

OSCINES.

## SYLVHIDAE.

CONN.—Common migrant and winter resident. Oct. 13–May.

391. *REGULUS CALENDULA* (Linné) Lichtenstein.**Ruby-crowned kinglet; Ruby-crested wren.**

Audubon, *Birds of Amer.*, 1841, vol. 2, p. 168, pl. 133.

*Motacilla calendula* Linné, *Syst. nat.*, ed. 12, 1766, vol. 1, p. 337.

“PENNSYLVANIA.”

Spruce and fir forests, and in migrations, tree growth generally; nests in conifers.

ME.—Common migrant and probably a rare summer resident in northern part (Ft. Kent, with young; near Bangor, nest without eggs). April 14–May 16 (May 23); (summer); Sept. 25–Oct. 22.

N. H.—Common migrant. April 14–May 11; Sept. 9–Oct. 20.

VT.—Common migrant. April 11; fall.

MASS.—Common migrant; casual in winter (at Highland Light, Jan. 15, 1892; Fresh Pond, Cambridge, Dec. 23, 1899). April 5–May 13; Sept. 23–Nov. 20 (winter).

R. I.—Common migrant. April–May 6; Oct.

CONN.—Common migrant. April 11–(May 16); Oct. 13–Nov. 24.

392. *POLIOPTILA CAERULEA* (Linné) Selater.**Blue-gray gnatcatcher.**

*Motacilla caerulea* Linné, *Syst. nat.*, ed. 12, 1766, vol. 1, p. 337.

“PENNSYLVANIA.”

*Culicivora caerulea* Lath. Audubon, *Birds of Amer.*, 1840, vol. 1, p. 244, pl. 70. Egg, Capen, 1886, pl. 2, fig. 3.

Woodlands; nests in trees.

ME.—Accidental visitor: Cape Elizabeth, Aug. 29, 1880; April 18, 1896.

N. H., VT.—Doubtfully recorded.

MASS.—Accidental visitor (? may once have bred near Taunton). Aug. 27–Dec. 18.

PASSERES.

OSCINES.

## SYLVIIDAE.

R. I.—Accidental visitor (several instances). May 23, June 24.

CONN.—Rare summer visitor (five instances, *see* Wauregan).  
April 6–Sept. 1.

## TURDIDAE.

393. *HYLOCICHLA MUSTELINA* (Gmelin) Ridgway.

**Wood thrush; Mountain thrush (Vt.); Swamp angel; Swamp thrush; Wood robin.**

*Turdus mustelinus* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 817.  
"IN NOVEBORACO." Seebohm and Sharpe, Monogr. of Turdidae,  
1898, pt. 4, pl. 43. Chapman, Handb. birds east. No. Amer.,  
1895, p. 395, fig. Egg, Capen, 1886, pl. 1, fig. 2.

Bushy growth near edges of moist shady woods; nests in bushes  
or trees.

ME.—Rare summer resident in southwestern part (to Franklin  
Co.). Spring–Sept. 6.

N. H.—Rare summer resident north to White Mt. valleys.  
May 15–Sept.

VT.—Uncommon summer resident at lower elevations (to Mt.  
Mansfield and Willoughby Lake). May 10–fall.

MASS.—Common summer resident, mainly at low elevations.  
(April 26) May 1–Sept. 28 (Oct. 10; Nov. 18, 1897, Wellesley  
Hills). Eggs, May 25–June 15 (June 26).

R. I.—Common summer resident. May 4–Sept. Eggs, May  
21–June 6.

CONN.—Common summer resident. May 1–Oct. Eggs, May  
27–June 21.

394. *HYLOCICHLA FUSCESCENS* (Stephens) Ridgway.

**Wilson's thrush; Cheeury; Nightingale; Swamp robin; Tawny  
thrush; Veery.**

*Turdus fuscescens* Steph., Shaw's Gen. zool., 1817, vol. 10, pt.

PASSERES.

OSCINES.

## TURDIDAE.

1, p. 182. "TO THE NORTH OF PENNSYLVANIA." Seebohm and Sharpe, Monogr. Turdidae, 1898, pt. 4, pl. 44. Egg, Capen, 1886, pl. 1, fig. 7.

Swamps and undergrowth along streams; nests on or near the ground, rarely in trees.

ME.—Common summer resident. (? April 17, 26) May 1–Sept. 20 (Oct. 1). Eggs, June 1–18 (July 1).

N. H.—Common summer resident below 1500 ft. May 10–Sept. Eggs, June 9.

VT.—Common summer resident. May 11–Sept. Eggs, May 30–June 28 (July 26).

MASS.—Common summer resident. (April 30) May 1–Sept. 27 (Oct. 14, 1871, Fresh Pond, Cambridge). Eggs, May 20–June 30.

R. I.—Common summer resident. (April 29)–Sept. 25. Eggs, May 26–31.

CONN.—Common summer resident. May 1–Sept. 1.

395. *HYLOCICHLA FUSCESCENS SALICICOLA* Ridgway.

**Willow thrush; Newfoundland thrush.**

Ridgw., Proc. U. S. nat. mus., 1882, vol. 4, p. 374. "ROCKY MOUNTAIN REGION. . . . FORT GARLAND, COLO.," etc.

Swamps and undergrowth bordering streams; nests on or near the ground.

MASS.—Rare migrant: Lanesboro, Sept. 27, 1900; Lowell (these are the race "*fuliginosa*" Howe, considered by the A. O. U. committee inseparable from *salicicola* and by Ridgway as synonymous with *H. fuscescens*).

R. I.—Rare migrant: Bristol, Sept. 24, 1899; Newport, Sept. 25, 1885.

CONN.—Rare migrant. May 5, 6, 14, 16; Sept. 23 (= *salicicola*, Bishop).

PASSERES.

OSCINES.

## TURDIDAE.

396. *HYLOCICHLA ALICIAE* (Baird) Ridgway.**Gray-cheeked thrush; Alice's thrush; Arctic thrush.**

*Turdus aliciae* Baird, U. S. war dept., expl. and surv. Miss. Riv. to Pac. Ocean, 1858, pt. 2, birds, p. 217. "ILLINOIS, . . . UPPER MISSOURI." Seebohm and Sharpe, Monogr. Turdidae, 1898, pt. 4, pl. 45, fig. 1.

Dense fir thickets in summer, moist shaded spots on migration; nests in low trees.

ME.—Uncommon migrant. May 20–25; Sept. 20.

N. H.—Rare migrant: Dublin, Oct. 2, 1899.

VT.—Rare migrant. May 18.

MASS.—Uncommon migrant. (May 13) May 17–June 3 (June 18); Sept. 10–Oct. 9.

R. I.—Uncommon migrant. May 20; Sept. 23–Oct. 7.

CONN.—Uncommon migrant.

397. *HYLOCICHLA ALICIAE BICKNELLI* Ridgway.**Bicknell's thrush.**

Ridgw., Proc. U. S. nat. mus., 1882, vol. 4, p. 377. "SLIDE MOUNTAIN, ULSTER COUNTY, NEW YORK." Bull. 50, U. S. nat. mus., 1907, pt. 4, p. 62. (Unfigured).

Fir and spruce thickets in summer, moist shady spots on migration; nests in low trees.

ME.—Common local summer resident on higher mountain tops (Mt. Abraham, Ktaadu, etc.); Maine breeding birds may be true *aliciae* (Knight). May–Sept. 20.

N. H.—Common summer resident above 3000 ft. to tree limit among White Mts., south to Sandwich Range. May 25–Sept. 15. Eggs, June 22.

VT.—Uncommon summer resident on higher mountains: Jay Peak, Mt. Annanance, Mt. Killington, Mt. Mansfield, Pico Peak, etc. May–Sept. Eggs, June 22.

MASS.—Uncommon migrant, and on summit of Mt. Greylock

## PASSERES.

## OSCINES.

## TURDIDAE.

(3505 ft.) a rare summer resident. (May 2)–May 30; summer; Sept. 18–Oct. 25.

R. I.—Uncommon migrant. Oct. 9.

CONN.—Uncommon migrant.

398. *HYLOCICHLA USTULATA SWAINSONII* (Cabanis) Ridgway.

**Olive-backed thrush; Flycatching thrush; Mosquito thrush; Swainson's thrush; Swamp robin.**

*Turdus swainsonii* Cab., Fauna Peruana, 1845–46, p. 188. "PERU"; specimen described from NEW JERSEY (= type locality). Seebohm and Sharpe, Monogr. Turdidae, 1898, pt. 4, pl. 45, fig. 3.

*Turdus ustulatus swainsonii* (Cab.). Chapman, Handb. birds east. No. Amer., 1895, p. 398. Egg, Capen, 1886, pl. 1, figs. 5, 6.

Cool shady thickets; nests in low trees or bushes.

ME.—Common migrant and in northern part a common summer resident. (May 6) May 9–26; summer–Oct. 23. Eggs, May 29–June 9 (early July).

N. II.—Common migrant, and from White Mts. north a common summer resident, local to the south. May 16–Oct. Eggs, June 16–July 11.

VT.—Common migrant and less common summer resident mainly in northern part and at higher levels south to Londonderry.

MASS.—Common migrant and in higher parts of Berkshire Co., an uncommon summer resident. May 3–June 7; summer; Sept. 9–Oct. 21.

R. I.—Common migrant. May 8–; Sept. 27–Oct. 2.

CONN.—Common migrant. May 10–26; Sept. 4–Oct. 13.

399. *HYLOCICHLA GUTTATA PALLASII* (Cabanis) Faxon and Hoffmann.

**Hermit thrush; Ground swamp robin (Me.); Rufous-tailed thrush; Solitary thrush; Swamp robin or thrush; Woodchuck.**

*Turdus pallasi* Cab., Arch. f. naturgesch., 1847, vol. 13, pt. 1,

PASSERES.

OSCINES.

## TURDIDAE.

p. 205. Locality not given; based on Cab., in Tschudi's Fauna Peruana. Seebohm and Sharpe, Monogr. Turdidae, 1898, pt. 4, pl. 46, fig. 2.

*Turdus aonalaschkae pallasii* (Cab.). Chapman, Handb. birds east. No. Amer., 1895, p. 400. Egg, Capen, 1886, pl. 1, figs. 3, 4.

Undergrowth of dry woods; nests on the ground.

ME.—Common migrant and summer resident; may winter rarely. (April 5) April 13–Nov. 26 (Dec. 25, 1904, Cape Elizabeth). Eggs, (May 1) May 17–June 10; July 14–Aug. 10 (2d brood).

N. H.—Common migrant and summer resident below 3000 ft. (April 5) April 10–Nov. 24. Eggs, –July 8.

VT.—Common migrant and summer resident. (April 4) April 10–fall.

MASS.—Common migrant, less common summer resident in western, local in eastern part (Cape Cod, Martha's Vineyard); rare winter resident near coast. (April 1) April 8–Nov. 30 (winter). Eggs, June 14.

R. I.—Common migrant and rare winter resident; rare local summer resident (Kent Co.). April 10–(summer); Oct.–Nov. 20 (Dec. 3, winter). Eggs, June 6.

CONN.—Common migrant, uncommon local summer resident (Bear Mt., Hartford Co., Norfolk, Salisbury); rare winter resident. (April 3) April 12–May; summer; Oct.–Nov. 20 (winter).

400. *PLANESTICUS MIGRATORIUS* (Linné) Coues.

**American robin.**

*Turdus migratorius* Linné, Syst. nat., ed. 12, 1766, vol. 1, p. 292. "IN AMERICA SEPTENTRIONALI." Audubon, Birds of Amer., 1841, vol. 3, p. 14, pl. 142. Egg, Capen, 1886, pl. 1, fig. 1.

Ubiquitous, but avoiding woods in breeding season; nests in trees, on ledges of buildings, or on the ground.

ME., N. H., VT.—Common migrant and summer resident; rare winter resident mainly in southern part. Mar. 3–Nov.; win-

## PASSERES.

## OSCINES.

## TURDIDAE.

ter. Eggs, (Me.) April 28–June 15; July–early Aug. (2d and 3d broods).

MASS.— Common migrant and summer resident; less common winter resident chiefly near coast. Mar. 8–Nov. 23; winter. Eggs, April 25–July 25 (2d brood) (! Oct. 29, 1903, Brandon).

R. I.— Common migrant, less common summer and winter resident. Mar. 11–Nov. 28; winter. Eggs, April 17–June 7; July.

CONN.— Common migrant and summer resident; less common winter resident. Feb. 18–Nov.; winter. Eggs, April 27–July 2.

401. *IXOREUS NAEVIUS* (Gmelin) Richmond.

**Varied thrush.**

*Turdus naevius* Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 817.  
“IN SINU AMERICAE NATCA.”

*Orpheus meruloides* (Swains.). Swainson and Richardson, Fauna Boreali-Amer., 1831, vol. 2, p. 187, pl. 38.

Evergreen forests; nests in trees.

MASS.— Accidental visitor: Ipswich, Dec. 1864.

402. *SIALIA SIALIS* (Linné) Haldeman.

**Bluebird.**

*Motacilla sialis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 187.  
“IN BERMUDIS & AMERICA CALIDIORE.”

*Sialia wilsonii* Swains. Audubon, Birds of Amer., 1841, vol. 2, p. 171, pl. 134. Egg, Capen, 1886, pl. 1, figs. 14, 15.

Orchards and open country; nests in hollow limbs or in bird houses.

ME.— Common migrant and summer resident. (Feb. 28) Mar. 10–Nov. 14. Eggs, April 21–May 15; June 24–July 6 (2d brood).

N. H.— Common migrant and less common summer resident at lower elevations. Mar. 2–Oct. 25. Eggs, May 1–June 25.

PASSERES.

OSCINES.

## TURDIDAE.

VT.—Common migrant and summer resident. (Feb. 23) Mar. 17–Oct. 16.

MASS.—Common migrant, less common summer resident; near coast, a rare winter resident (Melrose Highlands, Roxbury, Wellesley, Cape Cod). (Feb. 23) Mar. 1–Nov. 20 (Dec. 10, 29); winter. Eggs, April 15–June 5.

R. I.—Common migrant, less common summer resident, and occasional winter resident. Mar. 2–Nov. 27; winter. Eggs, (Mar. 1, Westerly) April 7–20.

CONN.—Common migrant and summer resident; uncommon winter resident. Mar. 5–Nov. 13; winter. Eggs, April 15–June 14.

## SPECIES INTRODUCED OR ERRONEOUSLY ACCREDITED.

COLYMBUS CRISTATUS Linné. **Crested grebe**.—An Old World species erroneously recorded by early writers.

AETHIA CRISTATELLA (Pallas). **Crested auklet**.—Doubtfully recorded from Chatham, Mass., winter of 1884–85.

CEPPHUS MANDTI (Lichtenstein). **Mandt's guillemot**.—The supposed Maine specimen proves to be *C. grylle*, and the Massachusetts record is too doubtful for acceptance.

PUFFINUS PUFFINUS (Brünnich). **Manx shearwater**.

PUFFINUS "CINEREUS" (= KUHLI Gould). **Mediterranean shearwater**.—European species listed from New England by several of the older writers but probably in error for *P. gravis* or *P. borealis*.

THALASSIDROMA PELAGICA (Linné). **Stormy petrel**.—Given on authority of Boardman as a rare summer visitor "off the coast of Eastern Maine," but the record is not allowed by Knight.

PELAGODROMA MARINA (Latham). **White-faced petrel**.—A specimen taken in lat. 40° 34' 18" N., long. 66° 09' W., Sept. 2, 1885, was included in Dr. J. A. Allen's list of Massachusetts birds (1886), but the locality is outside the limits covered by the present catalogue.

PHALACROCORAX GRACULUS (Linné). **Green cormorant**.—This European species is accredited to Massachusetts by W. B. O. Peabody, who doubtless confused it with the American *P. auritus*.



MERGELLUS ALBELLUS (Linné). **Smew**.—Accredited by Samuels to Massachusetts, but apparently without good reason

CHENALOPEX AEGYPTIACUS (Linné) Stephens.

**Egyptian goose.**

Salvadori, Cat. birds British mus., 1895, vol. 27, p. 169; Gould, Birds of Europe, 1837, vol. 5, pl. [353]. Egg, Seebohm, 1896, p. 36, pl. 12, fig. 1.

*Anas aegyptiaca* Linné, Syst. nat., ed. 12, 1766, vol. 1, pt. 1, p. 197. "IN AEGYPTO?"

Rivers and marshes; nests on the ground.

MASS.—The several instances of capture may relate to either accidental visitors or escapes.

ARDEA OCCIDENTALIS Audubon. **Great white heron**.—Recorded from Massachusetts in error for *Herodias egretta*.

SCOLOPAX RUSTICOLA Linné. **European woodcock**.—Doubtful reference to this species at Newport, R. I.; ? Maine.

TOTANUS NEBULARIUS (Gunnerus). **Greenshank**.—This Old World species was accredited to Connecticut by Linsley who states that a specimen was killed at Stratford in fall of 1842.

OREORTYX PICTUS (Douglas) Baird.

**Mountain partridge; Plumed partridge.**

*Ortyx picta* Dougl., Trans. Linn. soc. London, 1829, vol. 16, p. 143. "INTERIOR OF CALIFORNIA."

*Ortix plumifera* Gould. Audubon, Birds of Amer., vol. 5, p. 69, pl. 291.

Thickets and open woods; nests on the ground.

MASS.—Introduced unsuccessfully about 1890.

LOPHORTYX CALIFORNICUS (Shaw) Bonaparte.

**California partridge or quail.**

*Tetrao californicus* Shaw, Naturalist's miscellany, 1797 (?), vol. 9, pl. 345 and text. "CALIFORNIA."

*Callipepla californica* (Shaw). Gould, Monogr. of Odontophorinae No. Amer., 1850, p. [57], pl. [16]. Egg, Bendire, 1892, vol. 1, pl. 1, fig. 8-10.

Bushy lowlands; nests on the ground.

MASS.—Unsuccessfully introduced in Berkshire Co. about 1890.

CONN.—Introduced: Bridgeport, Dec. 1840.

LOPHORTYX GAMBELII Gambel.

**Gambel's partridge.**

Gambel (ex Nuttall), Proc. acad. nat. sci. Phila., 1843, p. 260. "WEST OF CALIFORNIA."

*Callipepla gambelii* (Nuttall). Cassin and Stephens, Illustr. birds of Calif., Texas, 1853, no. 2, p. 45, pl. 9. Egg, Bendire, 1892, vol. 1, pl. 1, fig. 11-14.

Bushy growth; nests on the ground.

MASS.— Unsuccessfully introduced in Berkshire and Bristol Cos. in 1893; Martha's Vineyard in 1890; Winchendon in 1891.

TETRAO UROGALLUS Linné.

**Capercaillie; Muscovian black gamecock; Wood grouse.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 159. "IN EUROPAE FRIGIDAE PINETIS PALUDOSIS." Ogilvie-Grant, Cat. birds British mus., 1893, vol. 22, p. 62. Dresser, Birds of Europe, 1871-81, vol. 5, pl. [489], p. [490]. Egg, Seebohm, 1896, p. 274, pl. 59, fig. 11.

Mountainous forests; nests on the ground.

VT.— Introduced about 1897 at Middlebury, Rutland, and elsewhere, unsuccessfully.

TYMPANUCHUS AMERICANUS (Reichenbach) Ridgway.

**Prairie hen; Prairie chicken.**

*Cupidonia americanus* Reich., Syst. avium, 1852, p. xxix. No locality.

*Tetrao cupido* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 93, pl. 296. Egg, Bendire, 1892, vol. 1, pl. 2, fig. 18-20.

Grassy prairies; nests on the ground.

N. H.— Unsuccessfully introduced at Blue Mt. Park about 1893.

MASS.— Unsuccessfully introduced: Barnstable and Berkshire Cos. in the '80's; between Concord and Belmont in 1884 or 1885; Martha's Vineyard, 1902; Naushon Id.; Springfield, 1893.

R. I.— Unsuccessfully introduced about 1897.

PEDIOECETES PHASIANELLUS (Linné) Elliot.

**Sharp tailed grouse.**

Baird, Brewer, and Ridgway, No. Amer. birds. 1874, vol. 3, p. 434, pl. 9, fig. 3 (head).

*Tetrao phasianellus* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 160. "CANADA."

*Pediacaetes phasianellus* (Linné). Elliot, Monogr. of Tetraoninae, 1865, pl. 15. Egg, Bendire, 1892, vol. 1, pl. 3, fig. 3-5.

Thickets and clearings; nests on the ground.

VT.— Unsuccessfully introduced about 1897.

MASS.— Unsuccessfully introduced near Westport about 1892.

PEDIOECETES PHASIANELLUS CAMPESTRIS Ridgway.

**Prairie sharp-tailed grouse.**

Ridgway, Proc. biol. soc. Washington, 1881, vol. 2, p. 93. "ILLINOIS, AND . . . ROSEBUD CREEK, MONTANA TERR."

*Tetrao phasianellus* Linné. Audubon, Birds of Amer., 1842, vol. 5, p. 110, pl. 298. Egg, Bendire, 1892, vol. 1, pl. 3, figs. 9, 10.

Woods and thickets; nests on the ground.

MASS.— Unsuccessfully introduced at Springfield, 1892.

COTURNIX COTURNIX (Linné) Lichtenstein.

**European quail; Migratory quail.**

Ogilvie-Grant, Cat. birds British mus., 1893, vol. 22, p. 235.

*Tetrao coturnix* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 161. "EUROPA, ASIA, AFRICA."

*Coturnix communis* Bonmarterre. Dresser, Birds of Europe, 1871-81, vol. 7, pl. [176]. Egg, Seebohm, 1896, p. 277, pl. 59, figs. 1, 3.

Meadows and fields; nests on the ground.

ME. — Introduced unsuccessfully in 1880; 2600 liberated, but failed to establish themselves. Nest at Falmouth, June.

VT.— Unsuccessfully introduced at Rutland, June, 1877.

MASS.— Unsuccessfully introduced at various places (No. Falmouth, Springfield, Wareham) between 1870 and 1890.

CONN.— Unsuccessfully introduced: Lakeville, 1878.

PERDIX PERDIX (Linné).

**European partridge; Hungarian partridge; Partridge.**

*Tetrao perdix* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 166. "IN EUROPAE AGRIS."

*Perdix cinerea* Latham. Dresser, Birds of Europe, 1871-81, vol. 7, p. [131], pls. [474, 475].

Open country and thickets; nests on the ground.

CONN.— Introduced in 1908.

PHASIANUS COLCHICUS Linné.

**English pheasant; Pheasant.**

Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 158. "AFRICA, ASIA." Ogilvie-Grant, Cat. birds British mus., 1893, vol. 22, p. 321. Dresser, Birds of Europe, 1871-81, vol. 7, pl. [169]. Egg, Seebohm, 1896, p. 275, pl. 59, fig. 9.

Bushy woods; nests on the ground, rarely in trees.

N. H.— Introduced from England at Wolfboro in 18th century.

MASS.— Unsuccessfully introduced in Berksh re Co.

*PHASIANUS TORQUATUS* Gmelin.

**Ring-necked pheasant; Denny pheasant; Ring pheasant.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 742. Refers to Latham's Synopsis: "CHINA. . . . ABOUT THE CASPIAN SEA, . . . GREAT TARTARY, AND IN THE SOUTH OF THE MONGOLIAN DESERT." Ogilvie-Grant, Cat. birds British mus., 1893, vol. 22, p. 332. Elliot, Monogr. Phasianidae, 1872, vol. 2, pl. 5.

Thickets; nests on the ground.

VT.— Introduced successfully at Shelburne.

MASS.— Introduced at Winchester and elsewhere, since 1894, and now well established at many places, mainly in eastern part.

CONN.— Successfully introduced (Fisher's Id.).

*COLUMBA DOMESTICA* Gmelin.

**Rock dove; Domestic pigeon; Dove; Pigeon.**

Gmel., Syst. nat., 1788, vol. 1, pt. 2, p. 769. No locality.

*Columba livia* Linné. Dresser, Birds of Europe, 1871-81, vol. 7, p. [11], pl. [457] (upper fig., wild bird). Tegetmeier, Pigeons: their structure, varieties, habits, and management, London, 1868, 190 pp., 16 pls. Egg, Seebohm, 1896, p. 158, pl. 47, fig. 9.

Sea cliffs, nesting in crevices of rocks; now domesticated about buildings.

ME.-CONN.— Introduced under domestication but now common in a feral state in towns and cities.

*BUTEO BOREALIS HARLANI* (Audubon). **Harlan's hawk.** Accredited to Massachusetts by Peabody on Nuttall's authority, probably in error.

"*BUTEO COOPERI*" Cassin. **Cooper's hen-hawk; California hawk.** A record for Cambridge, Mass., is believed to have been based on a very light-colored example of *B. lineatus*.

*SURNIA ULULA* (Linné). **European hawk owl.**— Specimens of this and the American race are said to have been taken at Houlton, Me., in 1877, but probably all were *S. u. caparoch*.

*DRYOBATES VILLOSUS LEUCOMELAS* (Boddaert). **Northern hairy woodpecker.**— Under this name Chadbourne lists the Hairy woodpecker from Williamstown, Mass., probably in error for *D. villosus*.

*SPHYRAPICUS VARIUS NUCHALIS* Baird. **Red-naped sapsucker.**— The supposed New Hampshire and Massachusetts specimens prove to be *S. varius varius*.

*AGYRTRIA VIRIDISSIMA* (Lesson). **Linne's hummingbird.** The authenticity of the supposed Cambridge specimen is open to doubt.

*ALAUDA ARVENSIS* Linné.

**Skylark; Laverock.**

Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 165. "IN EUROPAE APRICIS." Ridgway, *Bull.* 50, U. S. nat. mus., 1907, pt. 4, p. 292; Dresser, *Birds of Europe*, 1871–81, vol. 4, pl. [231]. Egg, Bendire, 1895, vol. 2, pl. 5, fig. 23.

Open country, fields and meadows; nests on the ground.

MASS.— Unsuccessfully introduced from Europe at Cambridge, prior to 1880.

*MUNIA ORYZIVORA* (Linné). **Java sparrow.**— A specimen was taken at Belmont, Mass., Nov. 2, 1884, but this and others are doubtless escaped cage birds.

*MUNIA ATRICAPILLA* (Vieillot). **Rufous and black weaver finch.**— A specimen of this Indian species, (doubtless an escape), was captured near Brunswick, Maine, in March, 1879 (recorded as *Amadina rubronigra*).

*STURNELLA NEGLECTA* Audubon. **Western meadowlark.**— Doubtfully recorded from Taunton, Mass.

*MEGAQUISCALUS MAJOR* (Vieillot). **Boat-tailed grackle.**— Formerly accredited to Maine, Massachusetts, and Connecticut but probably the Bronzed grackle was the species observed.

*SERINUS BUTYRACEA* (Linné) Bonaparte.

**Yellow finch.**

*Fringilla butyracea* Linné, *Syst. nat.*, ed. 10, 1758, vol. 1, p. 181. "MADERA."

*Serinus flaviventris* (Swains.). Sharpe, *Cat. birds British Mus.*, 1888, vol. 12, p. 353.

A South African species.

MASS.— Accidental visitor or escape: South Scituate, Feb. 1879, with flock of Redpolls.

*SERINUS SERINUS* (Linné) Sharpe.

**Serin finch.**

Sharpe, *Cat. birds British mus.*, 1888, vol. 12, p. 369.

*Fringilla serinus* Linné, Syst. nat., ed. 12, 1766, vol. 1, pt. 1, p. 320. "IN EUROPA AUSTRALI."

*Serinus hortulanus* Koeh. Dresser, Birds of Europe, 1871-81, vol. 3, [pl. 170]. Egg, Seebohm, 1896, p. 240, pl. 56, fig. 16.

Open country, orchards and gardens; nests in trees.

MASS.—Probably escaped cagebird; Springfield, Nov. 1866.

PASSER DOMESTICUS (Linné) Koeh.

**House sparrow; English sparrow; European house sparrow.**

Ridgway, Birds No. and Middle Amer., Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 143. Dresser, Birds of Europe, 1876, vol. 3, p. [587], pl. [176], fig. 1. Egg, Seebohm, 1896, p. 238, pl. 56, fig. 10-12.

*Fringilla domestica* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 183. "IN EUROPA PROPE RURA."

Villages and cities in vicinity of dwellings; nests in trees, vines, or any cavity.

ME.—Common resident; introduced at Portland in 1854 and 1858, at Lewiston in 1874, at Bangor about 1875.

N. H.—Resident, common about large cities, less common in White Mt. valleys.

VT.—Common resident about towns and cities; introduced at St. Johnsbury between 1874 and 1876.

MASS.—Common resident about towns and cities; introduced at Boston in 1858 and 1868, Brookline, 1868, Charlestown, 1869, Cottage City about 1882, Somerville, 1871.

R. I.—Common resident; introduced at Peacedale in 1858, Providence in 1866 or before, Newport, 1874.

CONN.—Common resident; introduced at New Haven in 1867, Norwich and Meriden between 1865 and 1870, Portland, 1869.

ACANTHIS BREWSTERII (Ridgway).

**Brewster's linnet.**

Brewster, Mem. Nuttall ornith. club, 1906, no. 4, pl. 6.

*Aegiothus (flavirostris var.) brewsterii* Ridgw., Amer. nat., 1872, vol. 6, p. 433. "WALTHAM, MASSACHUSETTS."

Status uncertain; probably a hybrid between *Acanthis linaria* and *Spinus pinus*.

MASS.—Only known specimen taken at Waltham, Nov. 1, 1870.

CARDUELIS CARDUELIS (Linné) Schäffer.

**European goldfinch.**

Ridgway, Bull. 50, U. S. nat. mus., 1901, pt. 1, p. 94. Egg, Seebohm, 1896, p. 241, pl. 56, fig. 17.

*Fringilla carduelis* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 180. "IN EUROPAE JUNIPERETIS."

*Carduelis elegans* Steph. Dresser, Birds of Europe, 1871-81, vol. 3, pl. [166].

Open country, orchards, and deciduous woods; nests in trees.

MASS.—Introduced in eastern part, but not well established; bred near Worcester, July 11, 1890.

CONN.—Escape, or straggler from lots introduced in neighboring country: New Haven, May 9, 1892.

RYNCHOPHANES MACOWNII (Lawrence). **McCown's longspur.**—Doubtfully recorded for Massachusetts on the basis of a market specimen said to have been taken at Ipswich, Jan. 7, 1877.

COTURNICULUS BAIRDII (Audubon). **Baird's sparrow.**—The first known specimens of *Passerculus princeps* were recorded under this name.

PEUCAEA AESTIVALIS BACHMANII (Audubon). **Bachman's finch.**—Recorded from Berlin, Mass., in error for *Ammodramus henslowii*.

PASSERINA CIRIS (Linné) Vieillot.

**Painted bunting; Nonpareil.**

*Emberiza ciris* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 179. "AMERICA."  
*Spiza ciris* (Wils.). Audubon, Birds of Amer., 1841, vol. 3, p. 93, pl. 169.

Thickets and hedges; nests in bushes or low trees.

ME.—"Escape," Portland, June 10, 1901.

MASS.—Four instances; believed to be escaped cape birds.

R. I.—Accidental visitor or escape: Scituate, summer, 1882.

PAROARIA CUCULLATA (Latham). **Crested cardinal.**—A South American species, once taken at Newport, R. I., Sept., 1885, but doubtless an escaped cagebird.

GUBERNATRIX CRISTATA (Vieillot).—A South American finch, of which a specimen supposed to be an escape, was taken at Providence, R. I., July 7, 1888.

VIREO ATRICAPILLUS Woodhouse. **Black-capped Vireo.**—Very doubtfully recorded from Lanesboro, Mass.

HELINALA SWAINSONII (Audubon). **Swainson's warbler.**—Erroneously accredited to Massachusetts by Audubon.

HELMINTHOPHILA LAWRENCEI (Herrick) Ridgway.

**Lawrence's warbler.**

Ridgway, Bull. 50, U. S. nat. mus., 1902, pt. 2, p. 152.

*Helminthophaga laurencii* Herriek, Proc. acad. nat. sci. Phila., 1874, p. 220, pl. 15. "NEAR CHATHAM, NEW JERSEY."

Probably a hybrid form between *H. pinus* and *H. chrysoptera*, which it resembles in habits.

MASS.— One recorded as seen at Dedham, May 27, 1908.

CONN.— Summer resident: Bridgeport, New Haven, Portland, Stamford. May 12–July. Eggs, June 14.

HELMINTHOPHILA LEUCOBRONCHIALIS (Brewster) Ridgway.

**Brewster's warbler; White-throated warbler.**

*Helminthophaga leucobronchialis* Brewst., Bull. Nuttall ornith. club, 1876, vol. 1, p. 1, pl. 1. "NEWTONVILLE, MASS."

Probably a hybrid form between *H. pinus* and *H. chrysoptera*, which it resembles in habits.

MASS.— Rare summer resident in eastern part (Hudson, Lexington, Newtonville, Jamaica Plain). May 14–summer. Eggs, June 8–18.

CONN.— Rare summer resident. May 6–Aug. 8. Eggs, June 24.

"MUSCICAPA MINUTA" Wilson = WILSONIA MICROCEPHALA (Ridgway). **Small-headed flycatcher.** An hypothetical species, described by Wilson and accredited to Massachusetts by Nuttall, Brewer, and Peabody.

PENTHESTES ATRICAPILLUS SEPTENTRIONALIS (Harris). **Long-tailed chickadee.** Erroneously recorded from Rhode Island; the specimen proves to be *P. a. atricapillus*.

SYLVIA ATRICAPILLA (Linné) Scopoli.

**European blackcap warbler; Blackcap.**

Dresser, Birds of Europe, 1871–81, vol. 2, p. [421], pl. [66]. Egg, Seebohm, 1896, p. 201, pl. 52, fig. 17–19.

*Motacilla atricapilla* Linné, Syst. nat., ed. 10, 1758, vol. 1, p. 187. "IN EUROPA."

Groves and orchards; nests in bushes.

MASS.— Unsuccessfully introduced at Cambridge prior to 1877.

TURDUS PILARIS Linné. **Fieldfare.**— A European species, once taken near Stamford, Conn., Mar. 1888, but this probably an escaped cagebird.

PLANESTICUS MERULA (Linné). **Blackbird.**— A male, probably an escape, spent the winter of 1908–09 near the Boston public garden.

SAXICOLA OENANTHE LEUCORHOA (Gmelin). **Greenland wheatear.**— The supposed Maine records in reality refer to birds taken across the line in New Brunswick.













AMNH LIBRARY



100115417

A. L. S. (74)